

LEARN MATHEMATICS

1. If matrix $P = \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}$, then the matrix P^{50} is equal to

- (1) $\begin{bmatrix} 1 & 0 \\ 50 & 1 \end{bmatrix}$
- (2) $\begin{bmatrix} 1 & 0 \\ 25 & 1 \end{bmatrix}$
- (3) $\begin{bmatrix} 1 & 0 \\ 75 & 1 \end{bmatrix}$
- (4) $\begin{bmatrix} 1 & 1 \\ 25 & 0 \end{bmatrix}$

Ans. (2)

Sol. $P^2 = \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}$

$$P^3 = \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}$$

$$P^3 = \begin{bmatrix} 1 & 0 \\ \frac{3}{2} & 1 \end{bmatrix}$$

Similarly

$$P^{50} = \begin{bmatrix} 1 & 0 \\ \frac{50}{2} & 1 \end{bmatrix}$$

$$P^{50} = \begin{bmatrix} 1 & 0 \\ 25 & 1 \end{bmatrix}$$

2. If the first sample A of 100 items has the mean 15 and standard deviation 3 and second sample B has 150 items. If the combined mean and standard deviation of items of both the sample is 15.6 and $\sqrt{13.44}$. Then the standard deviation of items of sample B is :

Ans. (4)

Sol. Combined mean = 15.6

$$\therefore 15.6 = \frac{100 \times 15 + 150 \times \bar{x}_B}{250}$$

$$\Rightarrow \bar{x}_B = 16 \quad (\text{mean of sample B})$$

$$\text{Combined standard deviation} = \sqrt{13.44}$$

$$\Rightarrow \text{combined variance } (\sigma^2) = 13.44$$

$$\sigma^2 = \frac{\sum x_i^2}{n} - (\bar{x})^2$$

$$13.44 = \frac{\sum x_i^2}{250} - 243.36$$

Activate Windows
Go to Settings to activate Windows.

3. The value of $x \in \left[\frac{\pi}{4}, \frac{\pi}{4} \right]$ for which

$$\begin{vmatrix} \sin x & \cos x & \cos x \\ \cos x & \sin x & \cos x \\ \cos x & \cos x & \sin x \end{vmatrix} = 0$$

(1) $-\frac{\pi}{4}$

(2) $-\frac{\pi}{8}$

(3) $\frac{\pi}{4}$

(4) $\frac{\pi}{8}$

Ans. (3)

Sol. $\begin{vmatrix} \sin x & \cos x & \cos x \\ \cos x & \sin x & \cos x \\ \cos x & \cos x & \sin x \end{vmatrix} = 0$

$R_1 \rightarrow R_1 + R_2 + R_3$

$$\Rightarrow \begin{vmatrix} \sin x + 2\cos x & \sin x + 2\cos x & \sin x + 2\cos x \\ \cos x & \sin x & \cos x \\ \cos x & \cos x & \sin x \end{vmatrix} = 0$$

$$(\sin x + 2\cos x) \begin{vmatrix} 1 & 1 & 1 \\ \cos x & \sin x & \cos x \\ \cos x & \cos x & \sin x \end{vmatrix} = 0$$

$C_2 \rightarrow C_2 - C_1, C_3 \rightarrow C_3 - C_1$

$$(\sin x + 2\cos x) \begin{vmatrix} 1 & 0 & 0 \\ \cos x & \sin x - \cos x & 0 \\ \cos x & 0 & \sin x - \cos x \end{vmatrix} = 0$$

$(\sin x + 2\cos x) (\sin x - \cos x)^2 = 0$

$\sin x = \cos x$ or $\sin x = -2\cos x$

$\tan x = 1$ or $\tan x = -2$

$\therefore x \in \left[-\frac{\pi}{4}, \frac{\pi}{4} \right]$

$x = \frac{\pi}{4}$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

Activate Windows
Go to Settings to activate Windows.

4. If \vec{a} and \vec{b} are two vectors such that $|\vec{a} \times \vec{b}| = 8$, $|\vec{a}| = 2$, $|\vec{b}| = 5$, then the value of $|\vec{a} \cdot \vec{b}|$ is :
(1) 6 (2) 3 (3) 12 (4) 9

Ans. (1)

Sol. $|\vec{a} \times \vec{b}| = |\vec{a}||\vec{b}| \sin \theta$
 $8 = 2 \times 5 \times \sin \theta$
 $\sin \theta = \frac{4}{5} \Rightarrow \cos \theta = \pm \frac{3}{5} \Rightarrow |\cos \theta| = \frac{3}{5}$
 $|\vec{a} \cdot \vec{b}| = |\vec{a}||\vec{b}||\cos \theta| = 2 \times 5 \times \frac{3}{5} = 6$

5. If function $f(x) : A \rightarrow B$, and $g(x) : B \rightarrow C$ are defined such that $(g(f(x)))^{-1}$ exist then $f(x)$ and $g(x)$ are :
(1) one-one and onto (2) many-one and onto
(3) one-one and into (4) many-one and into

Ans. (1)

Sol. Clearly $g(f(x))$ is one-one onto so $f(x)$ and $g(x)$ both are one-one and onto

6. If $a + b + c = 1$, $ab + bc + ca = 2$ and $abc = 3$, then the value of $a^4 + b^4 + c^4$ is:

Ans. 13

Sol. $(a + b + c)^2 = 1$
 $\Rightarrow a^2 + b^2 + c^2 + 2(ab + bc + ca) = 1$
 $\Rightarrow a^2 + b^2 + c^2 = -3 \dots(i)$
 $\Rightarrow ab + bc + ca = 2 \dots(ii)$
Squaring of equation (ii),
 $\Rightarrow a^2b^2 + b^2c^2 + c^2a^2 + 2(ab^2c + bc^2a + ca^2b) = 4$
 $\Rightarrow a^2b^2 + b^2c^2 + c^2a^2 + 2abc(a + b + c) = 4$
 $\Rightarrow a^2b^2 + b^2c^2 + c^2a^2 + 6 = 4$
 $\Rightarrow a^2b^2 + b^2c^2 + c^2a^2 = -2 \dots(iii)$
Squaring of equation (i),
 $\Rightarrow a^4 + b^4 + c^4 + 2(a^2b^2 + b^2c^2 + c^2a^2) = 9$
 $\Rightarrow a^4 + b^4 + c^4 - 4 = 9$
 $\Rightarrow a^4 + b^4 + c^4 = 13$

7. Which of the following value is just greater than $\left(1 + \frac{1}{10^{100}}\right)^{10^{100}}$
(1) 2 (2) 3 (3) 4 (4) 5

Ans. (2)

Sol. Let $10^{100} = n$
 $\left(1 + \frac{1}{n}\right)^n \approx (1 + \frac{1}{n})^n \approx (1 + \frac{1}{n})^n \approx (1 + \frac{1}{n})^n \approx (1 + \frac{1}{n})^n$

Activate Windows
Go to Settings to activate Windows.

Squaring of equation (1),
 $\Rightarrow a^4 + b^4 + c^4 + 2(a^2b^2 + b^2c^2 + c^2a^2) = 9$
 $\Rightarrow a^4 + b^4 + c^4 - 4 = 9$
 $\Rightarrow a^4 + b^4 + c^4 = 13$

7. Which of the following value is just greater than $\left(1 + \frac{1}{10^{100}}\right)^{10^{100}}$
(1) 2 (2) 3 (3) 4 (4) 5

Ans. (2)
Sol.

Let $10^{100} = n$
So, $\left(1 + \frac{1}{n}\right)^n = {}^nC_0 + {}^nC_1 \left(\frac{1}{n}\right) + {}^nC_2 \left(\frac{1}{n}\right)^2 + {}^nC_3 \left(\frac{1}{n}\right)^3 + \dots$
 $= 1 + 1 + \frac{n(n-1)}{2n^2} + \frac{n(n-1)(n-2)}{6n^3} + \dots$
 $\Rightarrow \left(1 + \frac{1}{n}\right)^n > 2$
Also $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e < 3$.

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 3

| JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

8. If a rectangle is inscribed in an equilateral triangle of side 2 2 , then the square of maximum area of rectangle is :

Ans. (3)

Sol.

A


Activate Windows
Go to Settings to activate Windows.

| JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

8. If a rectangle is inscribed in an equilateral triangle of side $2\sqrt{3}$, then the square of maximum area of rectangle is :

Ans. (3)

Sol.



Area of rectangle = $x \cdot h$ (i)

from $\triangle BDE$

$h = BE \tan 60^\circ$

$h = \frac{2\sqrt{3} - x}{2} \cdot \sqrt{3}$ (ii)

So area, $A = \frac{3}{2} (2\sqrt{3}x - x^2)$

For maxima $\frac{dA}{dx} = \frac{3}{2} (2\sqrt{3} - 2x) = 0$

$\Rightarrow x = \sqrt{3}$

From (ii) $h = \frac{3}{2}$

Area = $x \cdot h = \frac{9}{2}$

$(\text{Area})^2 = \frac{81}{4}$

9. If the coefficients of x^7 and x^8 in the expansion of $\left(2 + \frac{x}{3}\right)^n$ are equal then the value of n is :

(1) 53

(2) 54

(3) 55

(4) 56

Activate Windows
Go to Settings to activate Windows.

$(Area)^2 = 3$

9. If the coefficients of x^7 and x^8 in the expansion of $\left(2 + \frac{x}{3}\right)^n$ are equal then the value of n is :

- (1) 53 (2) 54 (3) 55 (4) 56

Ans. (3)

Sol. $\left(2 + \frac{x}{3}\right)^n = \sum_{r=0}^n {}^nC_r 2^{n-r} \left(\frac{x}{3}\right)^r$

Coefficient of $x^7 = {}^nC_7 2^{n-7} \cdot \left(\frac{1}{3}\right)^7$

Coefficient of $x^8 = {}^nC_8 2^{n-8} \cdot \left(\frac{1}{3}\right)^8$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 4

| JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

$\therefore {}^nC_7 \frac{2^{n-7}}{3^7} = {}^nC_8 \frac{2^{n-8}}{3^8}$

$\Rightarrow {}^nC_7 \cdot 6 = {}^nC_8$

$\Rightarrow \frac{6 \cdot n!}{7! \cdot (n-7)!} = \frac{n!}{8! \cdot (n-8)!}$

$\Rightarrow 48 = n - 7 \Rightarrow n = 55$

Activate Windows
Go to Settings to activate Windows.

10. If $f(x) = \begin{cases} \frac{P(x)}{x-2} & ;x \neq 2 \\ 7 & ;x = 2 \end{cases}$ and $P(x)$ is a polynomial such that $P''(x)$ is constant and $P(3) = 9$. If $f(x)$ is continuous

at $x = 2$ then find the value of $P(5)$

Ans. (39)

Sol. $P(x) = K(x-2)(x-\beta)$

$$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} \frac{K(x-2)(x-\beta)}{(x-2)}$$

$$\Rightarrow K(2-\beta) = 7 \dots(1)$$

$$\text{and } P(3) = K(3-2)(3-\beta) = 9$$

$$K(3-\beta) = 9 \dots\dots(2)$$

Divide equation (1) by (2)

$$\frac{2-\beta}{3-\beta} = \frac{7}{9} \Rightarrow \beta = \frac{-3}{2}$$

So, $K = 2$

$$\text{Then } P(x) = 2(x-2) \left(x + \frac{3}{2} \right)$$

$$P(5) = 2 \times (5-2) \times \left(5 + \frac{3}{2} \right) = 39$$

11. The number of real solutions of the equation $x^2 - |x| - 12 = 0$ is :

- (1) 0 (2) 1 (3) 3 (4) 2

Ans. (4)

Sol. $|x|^2 - |x| - 12 = 0$

$$|x| = 4, -3 \text{ (not possible)}$$

$$\Rightarrow |x| = 4 \Rightarrow x = \pm 4$$

\therefore Number of real solutions = 2

| JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

12. A coin is tossed n times. If the probability of getting at least one head is greater than 0.9, then the minimum value of n is :

- (1) 3 (2) 5 (3) 4 (4) 2

Ans. (3)

Sol. $1 - \binom{1}{2}^n > 0.9$

$\Rightarrow 0.1 > \left(\frac{1}{2}\right)^n \Rightarrow n = 4$

13. Negation of the statement :

“We will play football only if ground is not wet and there is no sunlight” is:

- (1) We will play football if ground is wet and there is no sunlight.
 (2) We will play football if ground is wet and there is sunlight.
 (3) There is no sunlight and ground is not wet and we will not play football.
 (4) There is sunlight or ground is wet and we will play football.

Ans. (4)

Sol. p: We will play football.

q: Ground is not wet.

r: There is no sunlight.

\therefore Given statement is $p \rightarrow (q \wedge r)$

\therefore Negation is $p \wedge \sim (q \wedge r)$

$p \wedge (\sim q \vee \sim r)$

14. If ${}^n C_0 + 3^n C_1 + 5^n C_2 + 7^n C_3 + \dots$ till $(n + 1)$ terms, is equal to $2^{100} \cdot 100$, then the value of $2 \binom{n-1}{2}$

(where [.] represents G.I.F.)

- (1) $(n - 1) \cdot 2^n$ (2) $(n + 1) \cdot 2^n$ (3) $n \times 2^{n+1}$ (4) $n \times 2^{n-1}$

Ans. (2)

Sol. ${}^n C_0 + 3^n C_1 + 5^n C_2 + 7^n C_3 + \dots$ (n + 1) terms = $\sum_{r=0}^n (2r + 1) \cdot 2^n C_r$

$= 2 \sum_{r=0}^n r^n C_r + \sum_{r=0}^n {}^n C_r$

$= 2n \cdot 2^{n-1} + 2^n = (n + 1) \cdot 2^n$

Activate Windows
Go to Settings to activate Windows.

r: There is no sunlight.
 \therefore Given statement is $p \rightarrow (q \wedge r)$
 \therefore Negation is $p \wedge \sim (q \wedge r)$
 $p \wedge (\sim q \vee \sim r)$

14. If ${}^n C_0 + 3^n C_1 + 5^n C_2 + 7^n C_3 + \dots$ till $(n + 1)$ terms, is equal to $2^{100} \cdot 100$, then the value of $2 \left[\frac{n-1}{2} \right]$ (where $[.]$ represents G.I.F.)
 (1) $(n - 1) \cdot 2^n$ (2) $(n + 1) \cdot 2^n$ (3) $n \times 2^{n+1}$ (4) $n \times 2^{n-1}$

Ans. (2)

Sol. ${}^n C_0 + 3^n C_1 + 5^n C_2 + 7^n C_3 + \dots$ (n + 1) terms = $\sum_{r=0}^n (2r + 1) \cdot {}^n C_r$
 $= 2 \sum_{r=0}^n r {}^n C_r + \sum_{r=0}^n {}^n C_r$
 $= 2n \cdot 2^{n-1} + 2^n = (n + 1) \cdot 2^n$

15. Evaluate $\int_{-1}^1 \log(x + \sqrt{x^2 + 1}) dx$
 (1) 0 (2) 1 (3) 2 (4) 3

Ans. (1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE # 6

Activate Windows
Go to Settings to activate Windows.

| JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol. $I = \int_{-1}^1 \log(x + \sqrt{x^2 + 1}) dx$

$$f(x) = \log(\sqrt{x^2 + 1} + x)$$

$$f(-x) = \log(\sqrt{x^2 + 1} - x)$$

$$= -f(x)$$

So $f(x)$ is an odd function.

$$\Rightarrow I = 0$$

16. If ${}^n P_r = {}^n P_{r+1}$ and ${}^n C_r = {}^n C_{r-1}$, then the value of n is :

Ans. (3)

Sol. ${}^n C_r = {}^n C_{r-1} \Rightarrow \frac{n-r+1}{r} = 1 \Rightarrow n+1 = 2r \dots\dots (1)$

$$\text{and } {}^n P_r = {}^n P_{r+1} \Rightarrow \frac{n!}{(n-r)!} = \frac{n!}{(n-r-1)!}$$

$$\Rightarrow n-r = 1 \dots\dots (2)$$

$$\text{Solving (1) \& (2) } n+1 = 2(n-1) \Rightarrow n = 3$$

17. Value of $\sum_{n=8}^{100} (-1)^n \left[\frac{n}{2} \right]$ is : (Where $[.]$ represent greatest integer function)

Ans. (4)

Sol. $\sum_{n=8}^{100} (-1)^n \left[\frac{n}{2} \right]$

$$\Rightarrow \underbrace{[4] + [-4.5]}_{-1} + \underbrace{[5] + [-5.5]}_{-1} + \dots\dots + [49] + [-49.5] + [50]$$

$$\Rightarrow -1 \times 46 + 50 = 4$$

Activate Windows
Go to Settings to activate Windows.

18. Evaluate $\cot\left(\frac{\pi}{24}\right)$

- (1) $\sqrt{6} - \sqrt{3} + \sqrt{2} - 2$ (2) $\sqrt{6} + \sqrt{3} - \sqrt{2} + 2$ (3) $\sqrt{6} + \sqrt{3} + \sqrt{2} - 2$ (4) $\sqrt{6} + \sqrt{3} + \sqrt{2} + 2$

Ans. (4)

Sol. $\cot\theta = \frac{\cos\theta}{\sin\theta} = \frac{2\cos^2\theta}{2\sin\theta\cos\theta}$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [Facebook](https://www.facebook.com/ResonanceEdu) | [Twitter](https://twitter.com/ResonanceEdu) | [YouTube](https://www.youtube.com/resowatch) | [Blog](http://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 7



JEE MAIN-2021 | DATE : 25-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

$$= \frac{1 + \cos 2\theta}{\sin 2\theta}$$

$$\therefore \cot \frac{\pi}{24} = \frac{1 + \cos \frac{\pi}{12}}{\sin \frac{\pi}{12}} \quad \left(\sin \frac{\pi}{12} = \frac{3-1}{2\sqrt{2}} \right) \& \quad \left(\cos \frac{\pi}{12} = \frac{3+1}{2\sqrt{2}} \right)$$

$$\Rightarrow \frac{1 + \frac{3+1}{2\sqrt{2}}}{\frac{3-1}{2\sqrt{2}}} = \frac{2\sqrt{2} + 3+1}{3-1}$$

$$\Rightarrow \frac{2\sqrt{2} + 3+1}{2}$$

Activate Windows
Go to Settings to activate Windows.

$$\Rightarrow \frac{2\sqrt{6} + 2\sqrt{2} + 3 + \sqrt{3} + \sqrt{3} + 1}{2} = \sqrt{6} + \sqrt{2} + \sqrt{3} + 2$$

19. If two lines $L_1 \equiv \frac{x-k}{1} = \frac{y-2}{2} = \frac{z-3}{3}$ and $L_2 \equiv \frac{x+1}{3} = \frac{y+2}{2} = \frac{z+3}{1}$ are coplanar. Then the value of k is :
- (1) -1 (2) 1 (3) 2 (4) -2

Ans. (2)

Sol.
$$\begin{vmatrix} k+1 & 2+2 & 3+3 \\ 1 & 2 & 3 \\ 3 & 2 & 1 \end{vmatrix} = 0$$

$$\Rightarrow (k+1)(-4) - 4(-8) + 6(2-6) = 0$$

$$\Rightarrow (k+1)(-4) = -8$$

$$k = 1.$$

20. If $y = f(x)$ is the solution of differential equation $xdy = (y + x^3 \cos x)dx$ and $f(\pi) = 0$ then $f\left(\frac{\pi}{2}\right)$ is:

- (1) $\frac{\pi^2}{4} + \frac{\pi}{6}$ (2) $\frac{\pi^2}{4} + \frac{\pi}{2}$ (3) $\frac{\pi^2}{6} + \frac{\pi}{4}$ (4) $\frac{\pi^2}{6} + \frac{\pi}{6}$

Ans. (2)

Sol.
$$\frac{xdy - ydx}{x^2} = x \cos x dx$$

$$\Rightarrow \int d\left(\frac{y}{x}\right) = \int x \cos x dx$$

$$\Rightarrow \frac{y}{x} = x \sin x + \cos x + c$$

$$\Rightarrow 0 = 0 - 1 + c \Rightarrow c = 1$$

$$\Rightarrow y = x^2 \sin x + x \cos x + x$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \frac{\pi^2}{4} + 0 + \frac{\pi}{2} = \frac{\pi^2}{4} + \frac{\pi}{2}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

Activate Windows
Go to Settings to activate Windows.

21. If three vector $\hat{i} + \hat{k}, \hat{b}\hat{i} + \hat{b}\hat{j} + \hat{c}\hat{k}$ and $a\hat{i} + a\hat{j} + b\hat{k}$ are coplanar then
 (1) $a^2 = bc$ (2) $b^2 = ac$ (3) $c^2 = ab$ (4) $ab = c$

Ans. (2)

Sol.
$$\begin{vmatrix} 1 & 0 & 1 \\ b & b & c \\ a & a & b \end{vmatrix} = 0$$

$1(b^2 - ac) + 1(ab - ab) = 0 \Rightarrow b^2 = ac$

22. If \vec{x} & \vec{y} are two vectors such that $|\vec{x}| = |\vec{y}|$ & $|\vec{x} - \vec{y}| = n|\vec{x} + \vec{y}|$, then the angle between \vec{x} & \vec{y}

- (1) $\cos^{-1}\left(\frac{1-n}{1+n}\right)$ (2) $\cos^{-1}\left(\frac{n^2+1}{1-n^2}\right)$ (3) $\cos^{-1}\left(\frac{n+1}{n-2}\right)$ (4) $\cos^{-1}\left(\frac{1-n^2}{n^2+1}\right)$

Ans. (4)

Sol. $|\vec{x} - \vec{y}| = n|\vec{x} + \vec{y}|$

$x^2 + y^2 - 2xy \cos\theta = n^2(x^2 + y^2 + 2xy \cos\theta)$

$x^2(1+n^2) - 2xy \cos\theta = n^2 x^2(1+n^2) + 2n^2 xy \cos\theta$

$2(1-n^2) = 2\cos\theta(n^2 + 1)$

$\cos\theta = \frac{1-n^2}{n^2+1}$

$\theta = \cos^{-1}\left(\frac{1-n^2}{n^2+1}\right)$

23. If combined equation of line $y = p(x)$ and $y = q(x)$ can be written as $(y - p(x))(y - q(x)) = 0$ then angle bisector of $x^2 - 4xy - 5y^2 = 0$ is :

- (1) $x^2 + 3xy + y^2 = 0$ (2) $x^2 + 3xy - y^2 = 0$
 (3) $x^2 - 3xy + y^2 = 0$ (4) $x^2 - 3xy - y^2 = 0$

Ans. (2)

Sol. Equation of angle bisector of homogeneous equation of pair of straight line $ax^2 + 2hxy + by^2$ is

$$\frac{x^2 - y^2}{a - b} = \frac{xy}{h}$$

Activate Windows
Go to Settings to activate Windows.

$$x^2(1+n^2-2\cos\theta) - 2xy(1+n^2-2\cos\theta) + y^2(1+n^2-2\cos\theta)$$

$$2(1-n^2) = 2\cos\theta (n^2 + 1)$$

$$\cos\theta = \frac{1-n^2}{n^2+1}$$

$$\theta = \cos^{-1}\left(\frac{1-n^2}{n^2+1}\right)$$

23. If combined equation of line $y = p(x)$ and $y = q(x)$ can be written as $(y - p(x))(y - q(x)) = 0$ then angle bisector of $x^2 - 4xy - 5y^2 = 0$ is :

- (1) $x^2 + 3xy + y^2 = 0$
- (2) $x^2 + 3xy - y^2 = 0$
- (3) $x^2 - 3xy + y^2 = 0$
- (4) $x^2 - 3xy - y^2 = 0$

Ans. (2)

Sol. Equation of angle bisector of homogeneous equation of pair of straight line $ax^2 + 2hxy + by^2$ is

$$\frac{x^2 - y^2}{a - b} = \frac{xy}{h}$$

for $x^2 - 4xy - 5y^2 = 0$

$$a = 1, h = -2, b = -5$$

so, equation of angle bisector is

$$\frac{x^2 - y^2}{1 - (-5)} = \frac{xy}{-2}$$

$$x^2 - y^2 + 3xy = 0$$

so, combined equation of angle bisector is $x^2 + 3xy - y^2 = 0$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | WhatsApp : 7340010333 | Facebook : facebook.com/ResonanceEdu | Twitter : twitter.com/ResonanceEdu | YouTube : www.youtube.com/resowatch | Blog : blog.resonance.ac.in

Activate Windows
Go to Settings to activate Windows.

24. Equation of a circle is $\text{Re}(z^2) + z(\text{Im}(z))^2 + z\text{Re}(z) = 0$ where $z = x + iy$ and a line passes through the vertex of parabola $x^2 - 6x + y + 13 = 0$ and the centre of circle, then y intercept of the line is :
 (1) -2 (2) -1 (3) 2 (4) 1

Ans. (2)

Sol. $z = (x + iy)$

So, $z^2 = x^2 - y^2 + i2xy$

Now $x^2 - y^2 + 2xy + 2x = 0$

$x^2 + y^2 + 2x = 0 \Rightarrow$ centre = $(-1, 0)$ and $x^2 - 6x + y + 13 = 0$

$(x-3)^2 = -(y+4)$

Vertex $(3, -4)$

\therefore Equation of line is $(y-0) = \frac{-4-0}{3+1}(x+1) \Rightarrow 4y = -4(x+1)$

$x + y + 1 = 0 \Rightarrow \frac{x}{-1} + \frac{y}{-1} = 1$

25. If $f(x) = \begin{cases} 5x+1 & ; x < 2 \\ \int_0^x (5+|1-t|)dt & ; x \geq 2 \end{cases}$

- (1) $f(x)$ is differentiable $\forall x \in \mathbb{R}$
- (2) $f(x)$ is continuous at $x = 2$ but not differentiable at $x = 2$
- (3) $f(x)$ is continuous at $x = 2$ but not differentiable at $x = 1$.
- (4) $f(x)$ is neither continuous nor differentiable at $x = 2$.

Ans. (2)

Sol. LHL = $\lim_{x \rightarrow 2^-} (5x + 1) = 11$

RHL = $\lim_{x \rightarrow 2^+} \int_0^x (5 + |1-t|)dt = \int_0^1 (5 + (1-t))dt + \int_1^2 (5 - (1-t))dt = 11$

$f(2) = 11$

So, $f(x)$ is continuous at $x = 2$

LHD at $x = 2$ is $\left. \frac{d}{dx} (5x + 1) \right|_{x=2} = 5$

RHD at $x = 2$ is $\left. \frac{d}{dx} \int_0^x (5 + |1-t|)dt \right|_{x=2} = 6$

LHD \neq RHD, So function is not differentiable at $x = 2$.

Activate Windows
Go to Settings to activate Windows.