


PART : PHYSICS

1. In the given arrangement, block A of mass m moving with speed 9 m/s , strikes with block B of mass $2m$ elastically. Then block B strikes with block C of mass $2m$ perfect in-elastically. Find the final speed of block C?



- (1) 3 m/s (2) 4 m/s (3) 5 m/s (4) 6 m/s

Ans. (1)

Sol. $9m = mv_A + 2mv_B$

$$9 = v_A + 2v_B \quad \dots(1)$$

$$9 = v_B - v_A \quad \dots(2)$$

Thus $v_B = 6\text{ m/s}$

Collision between B & C

$$2mv_B = 4mv_C$$


$$v_C = 3\text{ m/s}$$

2. A particle is thrown upward at $t = 0$. It attains maximum height of h . It is found at height $h/3$ at $t = t_1$ & $t = t_2$. Find t_1/t_2 :

- (1) $\frac{\sqrt{3}-1}{\sqrt{3}+1}$ (2) $\frac{1}{3}$ (3) $\frac{\sqrt{6}-\sqrt{5}}{\sqrt{6}+\sqrt{5}}$ (4) $\frac{1}{2}$

Ans. (3)

Sol.



Activate Windows
Go to Settings to activate Windows.

3. In YDSE, if the light used is shifted from orange to blue then which of the following is correct.

- (1) Fringe width first increases then decreases (2) Fringe width increase
(3) Fringe width decreases (4) Fringe width first decreases then increases


Ans. (3)

Sol. $\beta = \frac{\lambda D}{d}$

$$\lambda_{\text{blue}} < \lambda_{\text{orange}}$$

$$\beta_{\text{blue}} < \beta_{\text{orange}}$$

4. In the circuit shown in the figure, S_1 remains closed for a long time and S_2 remains open. Now S_2 is closed and S_1 is opened. Find out the di/dt just after that moment.



- (1) $-\frac{4\epsilon}{L}$ (2) $-\frac{6\epsilon}{L}$ (3) $-\frac{2\epsilon}{L}$ (4) $-\frac{\epsilon}{L}$

Ans. (2)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029


Toll Free : 1800 258 5555 [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) twitter.com/ResonanceEdu www.youtube.com/resowatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 2

Activate Windows
Go to Settings to activate Windows.

Sol. Before S_2 is closed and S_1 is opened current in the left part of the circuit = $\frac{\epsilon}{R}$. Now when S_2 closed S_1 opened, current through the inductor cannot change suddenly, current $\frac{\epsilon}{R}$ will continue to move in the inductor.



Applying KVL in loop 1.


$$L \frac{di}{dt} + \frac{\epsilon}{R} (2R) + 4\epsilon = 0$$

$$\frac{di}{dt} = -\frac{6\epsilon}{L}$$

5. Two identical tennis balls of mass m and charge q are hinged by a common support with the help of a string of length ' ℓ '. If the system is in equilibrium, then find the distance between the balls? Ignore gravitational interactions between balls. (θ is very small)

- (1) $\left(\frac{kq^2 \ell}{mg}\right)^{1/3}$ (2) $\left(\frac{3kq^2 \ell}{mg}\right)^{1/3}$ (3) $\left(\frac{2kq^2 \ell}{mg}\right)^{1/3}$ (4) $\left(\frac{3kq^2 \ell}{2mg}\right)^{1/3}$


Ans. (3)
Sol.



Force due to charge = $\frac{kq^2}{d^2}$

$$\tan\theta = \frac{kq^2}{mg d^2} \approx \sin\theta$$

Activate Windows
Go to Settings to activate Windows.



Ans. (3)
Sol.

$$\text{Force due to charge} = \frac{kq^2}{d^2}$$

$$\tan\theta = \frac{kq^2}{mg} \approx \sin\theta$$

$$\frac{kq^2}{mgd^2} = \frac{d}{2l}$$

$$d^3 = \frac{2kq^2 l}{mg} \Rightarrow d = \left(\frac{2kq^2 l}{mg} \right)^{1/3}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222


To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal | PAGE # 3

Activate Windows
Go to Settings to activate Windows.

6. For the given semicircle with centre O. choose the correct relation? If A, B, C & D are points on the semicircle such that $|\overline{AB}| = |\overline{BC}| = |\overline{CD}|$.



- (1) $2|\overline{BO}| = 2|\overline{AO}| = |\overline{AD}|$ (2) $|\overline{BC}| = 2|\overline{AO}| = |\overline{AD}|$ (3) $2|\overline{BO}| = |\overline{AO}| = |\overline{AD}|$ (4) $|\overline{BO}| = |\overline{AO}| = |\overline{AD}|$

Ans. (1)

Sol. For the given semicircle

$$|\overline{AO}| = |\overline{BO}| = |\overline{CO}| = |\overline{DO}| = \text{radius of circle}$$

$$|\overline{AD}| = \text{Diameter of circle}$$

$$|\overline{AD}| = 2|\overline{AO}| = 2|\overline{BO}|$$

7. A monoatomic gas is kept in a 1 litre container at pressure 1 atm. If average energy per molecule is 2×10^{-9} J, find no. of molecules in the container :

- (1) 0.75×10^{-6} (2) 0.75×10^9 (3) 0.5×10^{11} (4) 0.75×10^{11}

Ans. (4)

Sol. Total energy = $\frac{f}{2}nRT = \frac{3}{2}PV$

According to question $\frac{3}{2}PV = N \times 2 \times 10^{-9} \text{ J}$

$$\frac{3}{2} \times 10^5 \times 1000 \times 10^{-6} = N \times 2 \times 10^{-9}$$

$$N = \frac{3}{4} \times \frac{10^5 \times 10^3 \times 10^{-6}}{10^{-9}} = \frac{3}{4} \times 10^{11} = 0.75 \times 10^{11}$$

Activate Windows
Go to Settings to activate Windows.

$|\overline{AD}|$ = Diameter of circle

$$|\overline{AD}| = 2|\overline{AO}| = 2|\overline{BO}|$$

7. A monoatomic gas is kept in a 1 litre container at pressure 1 atm. If average energy per molecule is 2×10^{-9} J, find no. of molecules in the container :
- (1) 0.75×10^{-6} (2) 0.75×10^9 (3) 0.5×10^{11} (4) 0.75×10^{11}

Ans. (4)

Sol. Total energy = $\frac{f}{2}nRT = \frac{3}{2}PV$

According to question $\frac{3}{2}PV = N \times 2 \times 10^{-9}$ J

$$\frac{3}{2} \times 10^5 \times 1000 \times 10^{-6} = N \times 2 \times 10^{-9}$$

$$N = \frac{3}{4} \times \frac{10^5 \times 10^3 \times 10^{-6}}{10^{-9}} = \frac{3}{4} \times 10^{11} = 0.75 \times 10^{11}$$

8. The relative permittivity of distilled water is 81. The velocity of light in it will be ($\mu_r = 1$)
- (1) 3.3×10^7 (2) 5.3×10^7 (3) 4.3×10^7 (4) None

Ans. (1)

Sol. $v = \frac{c}{\sqrt{\mu_r \epsilon_r}}$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222


To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in


This solution was download from Resonance JEE (MAIN) 2021 Solution portal | PAGE # 4

Activate Windows
Go to Settings to activate Windows.


9. Match the following column if all the rods have same density and same radius.




(a) $\frac{M\ell^2}{12}$



(b) $\frac{M\ell^2}{3}$



(c) $\frac{2M\ell^2}{3}$



(d) $\frac{8M\ell^2}{3}$

- (1) (i) b (ii) d (iii) a (iv) c
 (3) (i) a (ii) c (iii) b (iv) d

- (2) (i) c (ii) d (iii) a (iv) d
 (3) (i) d (ii) a (iii) c (iv) b

Ans. (1)

Sol. (1) $\frac{M\ell^2}{3}$ (a)

(2) $\frac{2M(2\ell)^2}{3} \Rightarrow \frac{8M\ell^2}{3}$ (b)

(3) $\frac{M\ell^2}{12}$ (c)

(4) $\frac{2M4\ell^2}{12} \Rightarrow \frac{2M\ell^2}{3}$ (d)

10. A particle executing SHM having maximum kinetic energy 'E' and amplitude 'A'. Find displacement from mean position when its kinetic energy is $\frac{3E}{4}$:

- ... A ... A ... $\sqrt{3}$...

Activate Windows
Go to Settings to activate Windows.

(2) $\frac{2M(2\ell)^2}{3} \Rightarrow \frac{8M\ell^2}{3}$ (b)

(3) $\frac{M\ell^2}{12}$ (c)

(4) $\frac{2M4\ell^2}{12} \Rightarrow \frac{2M\ell^2}{3}$ (d)

10. A particle executing SHM having maximum kinetic energy 'E' and amplitude 'A'. Find displacement from mean position when its kinetic energy is $\frac{3E}{4}$:

- (1) $\frac{A}{2}$ (2) $\frac{A}{\sqrt{2}}$ (3) $\frac{\sqrt{3}}{2} A$ (4) A

Ans. (1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal | PAGE # 5

Resonance Educating for better tomorrow | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

Sol. $V = A\omega$, $KE_{max} = \frac{1}{2} m\omega^2 A^2$

$V = \omega\sqrt{A^2 - x^2}$ $KE_x = \frac{1}{2} m\omega^2 (A^2 - x^2)$

Activate Windows
Go to Settings to activate Windows.

Sol. $V = A\omega$, $KE_{\max} = \frac{1}{2}m\omega^2A^2$

$V = \omega\sqrt{A^2 - x^2}$, $KE_x = \frac{1}{2}m\omega^2(A^2 - x^2)$

$\frac{3}{4}KE_{\max} = KE_x$


$\frac{3}{4} \cdot \frac{1}{2}m\omega^2A^2 = \frac{1}{2}m\omega^2(A^2 - x^2)$

$\frac{3}{4}A^2 = A^2 - x^2$

$x^2 = \frac{A^2}{4}$

$x = \pm \frac{A}{2}$

11. For given PV diagram AB is isothermal at T_1 temperature, CD is also isothermal at T_2 temperature. given that $T_1 > T_2$. Then which one of the following is correct.




- (1) $W_{AB} > W_{BC}$ (2) $W_{BC} = W_{AD}$ (3) $W_{AB} < W_{BC}$ (4) $W_{AB} = W_{BC}$

Ans. (1)

12. A container of area 'A' filled with liquid, a small hole of area 'a' is made at the bottom of curved surface


Activate Windows
Go to Settings to activate Windows.



- (1) $W_{AB} > W_{BC}$ (2) $W_{BC} = W_{AD}$ (3) $W_{AB} < W_{BC}$ (4) $W_{AB} = W_{BC}$

Ans. (1)

12. A container of area 'A' filled with liquid, a small hole of area 'a' is made at the bottom of curved surface as shown. Find the coefficient of friction between container and ground in order to prevent it from sliding.



- (1) $\frac{a}{A}$ (2) $\frac{2a}{A}$ (3) $\frac{a}{2A}$ (4) $\frac{3a}{2A}$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE # 6

Activate Windows
Go to Settings to activate Windows.

Ans. (2)

Sol. Force exerted by leaving water on the container,

$$F = \left(\frac{dm}{dt}\right)v = (\rho av)v$$

$$= \rho av^2$$

By conservation of energy $v = \sqrt{2gh}$

$$\mu mg = \rho a(2gh)$$

$$\mu \rho (Ah)g = \rho a2gh$$


$$\mu = \frac{2a}{A}$$

13. Two solid spheres of radius $4R$ and R and having same mass density ρ kept at a distance of $8R$. Find Moment of Inertia of the system about the axis passes through centre of line joining their centres.

- (1) $\frac{9660}{5} \pi \rho R^5$ (2) $\frac{5800}{3} \pi \rho R^5$ (3) $\frac{9664}{4} \pi \rho R^5$ (4) $\frac{9664}{3} \pi \rho R^5$

Ans. (2)

Sol.



Let mass density = ρ

$$m_2 = \frac{4}{3} \pi R^3 \rho$$

$$m_1 = \frac{4}{3} \pi (4R)^3 \rho$$

$$m_1 = 64 m_2$$

Activate Windows
Go to Settings to activate Windows.

$$m_2 = \frac{4}{3} \pi R^3 \rho$$

$$m_1 = \frac{4}{3} \pi (4R)^3 \rho$$

$$m_1 = 64 m_2$$

$$I_{\text{final}} = [I_2 + m_2 (4R)^2] + [I_1 + m_1 (4R)^2]$$

$$= \frac{2}{5} m_2 R^2 + 16 m_2 R^2 + \frac{2}{5} m_1 (4R)^2 + m_1 (4R)^2$$

$$= \frac{2}{5} m_2 R^2 + 16 m_2 R^2 + \frac{2}{5} 64 m_2 (4R)^2 + 64 m_2 (4R)^2 = \frac{5800}{3} \pi \rho R^5$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222


To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE # 7

Resonance | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS

14. For the given circuit, switch is closed at t = 0 find time after which voltage across capacitor becomes 50V



- (1) 69.3 μsec (2) 60 μsec (3) 50 μsec (4) 40 μsec

Ans. (1)


Activate Windows
Go to Settings to activate Windows.

$$50 = 100e^{-RC}$$

$$t = RC \ln 2$$

$$= 100 \times 10^{-6} \ln 2 = 10^{-4} \times 0.693 \text{ sec} = 69.3 \mu\text{sec}$$


15. Figure shows two capacitors in steady state. Now the cell is removed and a dielectric of dielectric constant k is inserted between the plates of the capacitor of capacitance C . Find new potential difference across any of the capacitors :




- (1) $\frac{V}{2+k}$ (2) $\frac{2V}{2+k}$ (3) $\frac{3V}{2+k}$ (4) $\frac{V}{1+2k}$

Ans. (3)
Sol.

Just after removing cell



After inserting dielectric



Conservation of charge

$$2CV + CV = 2C \cdot V_{ab} + Ck V_{ab}$$

$$\therefore V_{ab} = \frac{3CV}{2C + Ck}$$

$$V_{ab} = \frac{3V}{2 + K}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 8

Activate Windows
Go to Settings to activate Windows.

16. A particle of mass 9.1×10^{-31} kg is moving with velocity 6 m/s. Momentum of photon is 2×10^{-27} kg m/s. If de-Broglie wavelength of the particle is K times of the wavelength of photon. Find value of K :
 (1) 366 (2) 566 (3) 322 (4) 166

Ans. (1)

Sol. De-Broglie wavelength of particle

$$\lambda = \frac{h}{p} = \frac{h}{9.1 \times 10^{-31} \times 6}$$

Wavelength of photon

$$\lambda_{\text{photon}} = \frac{h}{p} = \frac{h}{2 \times 10^{-27}}$$


From question

$$K\lambda_{\text{photon}} = \lambda_{\text{particle}}$$

$$\Rightarrow K = \frac{2 \times 10^{-27}}{9.1 \times 10^{-31} \times 6} = 366$$

17. Two prism of same angle of refraction are arranged as shown. If the light incident, on the system goes undeviated. Then find wave length of incident light

$$\left(\begin{array}{l} \text{given } n_1 = 1.2 + \frac{10.8 \times 10^{-14}}{\lambda^2} \\ n_2 = 1.45 + \frac{1.8 \times 10^{-14}}{\lambda^2} \end{array} \right)$$




- (1) 600 nm (2) 675 nm (3) 575 nm (4) 475 nm

Ans. (1)

Activate Windows
Go to Settings to activate Windows.

18. For the given circuit find current 'I' and the phase difference between V and I :



- (1) 8.8A, $\tan^{-1} 1.83$ (2) 6.8A, $\tan^{-1} 2.83$ (3) 5.8A, $\tan^{-1} 0.83$ (4) 7.8A, $\tan^{-1} 2.53$

Ans. (1)

Sol. $\omega = 2\pi f = 100\pi$

Thus $x_L = \omega L = 100 \times \frac{22}{7} \times 0.07 = 22\Omega$


Thus, $z = \sqrt{R^2 + x_L^2}$

$z = \sqrt{(12)^2 + (22)^2}$

$z \approx 25.00$

$I = \frac{220}{25} = 8.8$

$\tan \phi = 22/12 = 1.83$



Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029


Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resowatch blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PAGE # 10

Activate Windows
Go to Settings to activate Windows.

19. A particle of mass 'm' is projected from the surface of planet 'A' of mass M and radius R. What should be the velocity of projection such that particle reaches to another planet 'B', which is at a distance of 8R from planet 'A'. Planet 'B' has mass 9M and radius 2R.



- (1) $\sqrt{\frac{1}{5} \frac{GM}{R}}$ (2) $\sqrt{\frac{4}{7} \frac{GM}{R}}$ (3) $\sqrt{\frac{2}{5} \frac{GM}{R}}$ (4) $\sqrt{\frac{7}{3} \frac{GM}{R}}$

Ans. (2)

Sol. Let at a distance 'X' from planet 'A', the net gravitational field becomes zero

$$\frac{GM}{X^2} = \frac{G \times 9M}{(8R - X)^2}$$

$$(3X)^2 = (8R - X)^2$$

$$X = 2R$$

Now, Particle should be projected such that it covers a minimum distance of '2R'.

Thus

$$\frac{1}{2}mv^2 - \frac{GMm}{R} - \frac{G(9M)m}{7R} = -\frac{GMm}{2R} - \frac{G(9M)m}{6R}$$

$$\frac{1}{2}v^2 = \frac{2}{7} \frac{GM}{R} \Rightarrow v = \sqrt{\frac{4}{7} \frac{GM}{R}}$$

20. Circular scale divisions of a screw gauge is 50. Five full rotations advances circular scale by 5mm.

Statement-1 : Least count of screw gauge is 0.001 cm

Statement-2 : Least count = $\frac{\text{pitch}}{\text{Total no. of divisions}}$

- (1) Statement-1 & 2 both are true
 (2) Statement-1 & 2 both are true statement-2 is correct explant of statement-1
 (3) Statement-1 is false Statement-2 is true

Activate Windows
Go to Settings to activate Windows.

20. Circular scale divisions of a screw gauge is 50. Five full rotations advances circular scale by 5mm.

Statement-1 : Least count of screw gauge is 0.001 cm

Statement-2 : Least count = $\frac{\text{pitch}}{\text{Total no. of divisions}}$

- (1) Statement-1 & 2 both are true
- (2) Statement-1 & 2 both are true statement-2 is correct explant of statement-1
- (3) Statement-1 is false Statement-2 is true
- (4) Statement-2 is true Statement-1 is false.

Ans. (3)

Sol. $LC = 1\text{mm}/50 = 0.02\text{ mm}$

Statement -1 is wrong. Statement-2 is right

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005


Ph. No.: +91-744-2777777, 2777700 | **FAX No.:** +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029


Toll Free : 1800 258 5555  **7340010333**  [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu)  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal


PAGE # 11

 **JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-1) | PAPER-1 | MEMORY BASED | PHYSICS**

21. Figure shows a conductor of tapered cone shape. As one goes from left to right on conductor, choose correct option



Activate Windows
Go to Settings to activate Windows.



- (1) current decreases
- (2) drift velocity of electron increases
- (3) electric field inside conductor decreases
- (4) All of the above

Ans. (2)

Sol. $i = nAev_d$
 $\therefore v_d$ increases as area A decrease

22. A body cools down from 61°C to 49°C in 4 min. and the surrounding temperature is 30°C. Find the time taken by this body to cool down from 49°C to 37°C.

- (1) 1.62 min
- (2) 1.25 min
- (3) 7.69 min
- (4) 0.92 min.

Ans. (3)

Sol. $\frac{61 - 49}{4} = K \left[\frac{61 + 49}{2} - 30 \right] \dots(1)$

$\frac{49 - 37}{t} = k \left[\frac{49 + 37}{2} - 30 \right] \dots(2)$

Dividing (1) and (2)

$\frac{t}{4} = \frac{25}{13}$

$t = \frac{4 \times 25}{13} = 7.69$

23. A bar-magnet of magnetic moment 9.85 A-m² and moment of inertia I = 10⁻⁶ kg-m² makes 10 oscillations in 5 sec. in uniform magnetic field. Find intensity of magnetic field.

(Take $\pi^2 = 9.85$)

- (1) 20 μ T
- (2) 25 μ T
- (3) 16 μ T
- (4) 10 μ T

Ans. (3)

Activate Windows
Go to Settings to activate Windows.

$= 16 \mu l$

24. If two discs are of same mass density but having different radii R and r.
 I_1 = Moment of inertia of disc of radius R, about an axis perpendicular to the plane of disc and passing through its centre.
 I_2 = Moment of inertia of disc of radius r, about one of its diameter. Then choose correct option

- (1) $\frac{I_1}{I_2} = \frac{2r^4}{R^2}$ (2) $\frac{I_1}{I_2} = \frac{2R^4}{r^4}$ (3) $\frac{I_1}{I_2} = \frac{2r^2}{R^2}$ (4) $\frac{I_1}{I_2} = \frac{2r}{R}$

Ans. (2)

Sol. Let mass density be σ


$M_1 = \pi R^2 \sigma, M_2 = \pi r^2 \sigma$

$I_1 = \frac{1}{2} M_1 R^2 = \frac{\pi \sigma R^4}{2}$

$I_2 = \frac{1}{4} M_2 r^2 = \frac{\pi \sigma r^4}{4}$

$\frac{I_1}{I_2} = \frac{2R^4}{r^4}$

25. Find the equivalent capacitance of shown arrangement if all the plates are identical having surface area A.



- (1) $\frac{24}{15} \frac{A \epsilon_0 d}{K}$ (2) $\frac{17}{15} \frac{A \epsilon_0 d}{K}$ (3) $\frac{15}{34} \frac{A \epsilon_0 d}{K}$ (4) $\frac{34}{15} \frac{A \epsilon_0 d}{4K}$

Ans. (3)

Activate Windows
Go to Settings to activate Windows.

$$C_{eq} = \frac{15 A \epsilon_0 d}{34 K}$$

26. An electron is revolving in a circular orbit of radius 0.5 m, with a speed of 2.2×10^{-6} m/s. Find equivalent current ?

- (1) 1.12×10^{-25} A (2) 2.15×10^{-25} A (3) 1.12×10^{-15} A (4) 2.15×10^{-13} A

Ans. (1)

Sol. $i = \frac{q}{t} = \frac{q}{2\pi r / v}$
 $= \frac{1.6 \times 10^{-19} \times 2.2 \times 10^{-6}}{2\pi \times 0.5} = 1.12 \times 10^{-25}$ A

27. A capacitor ($C = 100 \mu\text{F}$) discharging against a resistor R, at same time a radioactive substance decays with mean life 30 ms, if the ratio of charge on capacitor and activity of substance remain same for all the time then, find the resistance :

- (1) 300Ω (2) 432Ω (3) 450Ω (4) 250Ω

Ans. (1)

Sol. $q = q_0 e^{-\frac{t}{RC}} \dots(1)$

$A = A_0 e^{-\lambda t} \dots(2)$

Divide (1) & (2)

$$\Rightarrow \frac{q}{A} = \frac{q_0}{A_0} \frac{e^{-\frac{t}{RC}}}{e^{-\lambda t}} \Rightarrow -\lambda t = -\frac{t}{RC}$$

$RC = 1/\lambda = 30 \times 10^{-3}$ sec

$R = 300$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

Activate Windows
Go to Settings to activate Windows.