

### PART : PHYSICS

1. Two masses each of mass 1 kg are separated by a distance 2R, rotating under their mutual gravitation force. Find their angular velocity :

- (1)  $\frac{1}{2}\sqrt{\frac{G}{R^3}}$       (2)  $\sqrt{\frac{G}{R^3}}$       (3)  $\sqrt{\frac{2G}{R^3}}$       (4)  $\sqrt{\frac{G}{2R^3}}$

Ans. (1)

Sol.  $\frac{Gm^2}{4R^2} = m\omega^2 R$

$$\omega = \sqrt{\frac{Gm}{4R^3}}$$

$$\omega = \frac{1}{2}\sqrt{\frac{G}{R^3}}$$

2. A conducting wire has resistance 16Ω at 15°C and 20Ω at 100°C. Find temperature coefficient of resistance

- (1)  $\frac{1}{340} \text{ } ^\circ\text{C}^{-1}$       (2)  $\frac{1}{200} \text{ } ^\circ\text{C}^{-1}$       (3)  $\frac{1}{470} \text{ } ^\circ\text{C}^{-1}$       (4)  $\frac{1}{300} \text{ } ^\circ\text{C}^{-1}$


Ans. (1)

Sol.  $R' = R(1 + \alpha\Delta t)$

$$20 = 16(1 + \alpha \cdot 85)$$

$$\alpha = \frac{\frac{20}{16} - 1}{85} = \frac{1}{4 \times 85} = \frac{1}{340} \text{ } ^\circ\text{C}^{-1}$$

3. In the figure each side has length ℓ. Find electric field at centre :



Activate Windows  
Go to Settings to activate Windows.

$(1) \frac{kq}{2l^2}(\sqrt{2}-1)$       $(2) \frac{kq\sqrt{2}}{2l^2}$       $(3) \frac{kq}{l^2} \sqrt{\left(\frac{5}{4} + \frac{1}{\sqrt{2}}\right)}$       $(4) \frac{kq}{l^2}(\sqrt{2}+1)$

Ans. (3)


## Resonance Eduventures Ltd.

**Reg. Office & Corp. Office :** CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005  
**Ph. No.:** +91-744-2777777, 2777700 | **FAX No.:** +91-022-39167222  
**To Know more :** sms RESO at 56677 | **Website :** www.resonance.ac.in | **E-mail :** [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | **CIN :** U80302RJ2007PLC024029  
**Toll Free : 1800 258 5555**    [7340010333](https://www.facebook.com/ResonanceEdu)    [facebook.com/ResonanceEdu](https://twitter.com/ResonanceEdu)    [twitter.com/ResonanceEdu](https://www.youtube.com/resowatch)    [www.youtube.com/resowatch](https://www.youtube.com/resowatch)    [blog.resonance.ac.in](https://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal     PAGE # 1

**Resonance®** Educating for better tomorrow | **JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS**

Sol.



Activate Windows  
Go to Settings to activate Windows.



$$\frac{(\sqrt{2}l)^2}{l^2}$$

$$E = \frac{kq}{l^2} \left( \sqrt{1 + \frac{1}{4} + 2 \times \frac{1}{2} \times \frac{1}{\sqrt{2}}} \right)$$

$$= \frac{kq}{l^2} \sqrt{\frac{5}{4} + \frac{1}{\sqrt{2}}}$$

4. A force  $F = F_0 \left[ 1 - \left( \frac{T-t}{T} \right)^2 \right]$  start to act on a ball of mass  $m$  at  $t = 0$ . Initially ball was at rest. Find velocity of the ball at  $t = 2T$  :

- (1)  $\frac{F_0}{m} \left[ \frac{4T}{3} \right]$       (2)  $\frac{F_0}{m} \left[ \frac{3T}{2} \right]$       (3)  $\frac{F_0}{m} \left[ \frac{T}{3} \right]$       (4)  $\frac{F_0}{m} \left[ \frac{T}{2} \right]$

Ans. (1)

Sol. acceleration of ball

$$a = \frac{F}{m}$$

$$a = \frac{F_0}{m} \left[ 1 - \left( \frac{T-t}{T} \right)^2 \right]$$

$$\frac{dv}{dt} = \frac{F_0}{m} \left[ 1 - \left( \frac{T-t}{T} \right)^2 \right]$$

$$\int_0^v dv = \frac{F_0}{m} \int_0^{2T} \left[ 1 - \left( \frac{T-t}{T} \right)^2 \right] dt$$

$$V = \frac{F_0}{m} \left[ t + \frac{1}{3T^2} (T-t)^3 \right]_0^{2T}$$

$$V = \frac{F_0}{m} \left\{ \left[ 2T + \frac{1}{3T^2} (T-2T)^3 \right] - \left[ 0 + \frac{T^3}{3T^2} \right] \right\}$$


$$V = \frac{F_0}{m} \left[ \frac{4T}{3} \right]$$

### Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005  
 Ph. No. : +91 744 2777777, 2777700 | FAX No. : +91 022 29167222


Activate Windows  
Go to Settings to activate Windows.

5. A block of mass  $m$  as shown in figure is released from rest from the top of a fixed smooth hemisphere. Find the angle made by this particle with vertical at the instant when it loses contact with the hemisphere:



- (1)  $\cos^{-1}(2/3)$       (2)  $\cos^{-1}(1/3)$       (3)  $\cos^{-1}(1/2)$       (4)  $\cos^{-1}(1/4)$


Ans. (1)  
Sol.



From work energy theorem  $W = \Delta K$   
 $Mg(R - R \cos \theta) = \frac{1}{2} mv^2$   
 $v = \sqrt{2gR(1 - \cos \theta)}$

To loose contact  $\frac{mv^2}{R} = mg \cos \theta$   
 $M2g(1 - \cos \theta) = mg \cos \theta$   
 $2 - 2 \cos \theta = \cos \theta$   
 $\cos \theta = 2/3 ; \theta = \cos^{-1}(2/3)$

6. Figure shows variation of potential energy ( $U$ ) verses displacement ( $x$ ) graph :




Find the correct statement :

- (1)  $v$  &  $KE$  is least and body has constant speed

Activate Windows  
Go to Settings to activate Windows.





Find the correct statement :

- (1)  $x < x_1$ , KE is least and body has constant speed.
- (2)  $x = x_2$ , K.E is minimum
- (3)  $x > x_2$  K.E. is maximum and velocity is maximum
- (4)  $x > x_2$  K.E. is minimum so velocity is minimum

Ans. (1)

Sol.  $K + U = E$  mechanical energy = constant

### Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No.: +91-022-39167222

To Know more : sms RESO at 56677 | Website : [www.resonance.ac.in](http://www.resonance.ac.in) | E-mail : [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](http://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE # 3


**Resonance** Educating for better tomorrow | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

7. Magnetic flux in a circular loop having resistance of  $2\Omega$  is varying with time as  $\phi = 10t^2 + 20t$ . What will be the current in circuit at  $t = 5$ sec. (in Ampere)

- (1) 20
- (2) 40
- (3) 60
- (4) 80

Ans. (3)

Sol.



Activate Windows  
Go to Settings to activate Windows.

thus,  $i = \frac{\varepsilon}{R} = \frac{120}{2} = 60 \text{ Amp.}$

8. Match list-I with list-II and select the correct option from below the list

**List-I**

1. Electric field intensity (E)
2. Magnetic permeability ( $\mu_0$ )
3. Electrical permittivity ( $\epsilon_0$ )
4. Capacitance (C)

- (1) 1 → (ii), 2 → (iii), 3 → (iv), 4 → (i)  
(3) 1 → (iii), 2 → (iv), 3 → (ii), 4 → (i)

**List-II (Dimension)**

- (i)  $M^{-1}L^{-2}I^2T^{+4}$
  - (ii)  $M^{-1}L^{-3}T^4I^2$
  - (iii)  $MLI^{-1}T^{-3}$
  - (iv)  $ML^2T^{-4}I^{-2}$
- (2) 1 → (iii), 2 → (i), 3 → (iii), 4 → (ii)  
(4) 1 → (ii), 2 → (iv), 3 → (i), 4 → (iii)

Ans. (3)

9. A block of mass 1kg connected to a massless spring fixed from one end executing SHM. Initially at mean position. Its amplitude is 5 cm and time period is 0.2 sec. Find potential energy after 0.05 sec.

- (1)  $\frac{1}{2} \text{ J}$                       (2)  $\frac{3}{4} \text{ J}$                       (3)  $\frac{5}{6} \text{ J}$                       (4)  $\frac{5}{4} \text{ J}$

Ans. (4)

Sol. At  $t = \frac{T}{4}$  particle is at extreme.  $v = K.E_{\text{max}} = \frac{1}{2} m\omega^2 A^2$ ,  $T = 2\pi\sqrt{\frac{m}{K}}$

$$= \frac{1}{2} m \left( \frac{2\pi}{0.2} \right)^2 A^2$$

$$= \frac{1}{2} \times 100 \pi^2 \times \frac{25}{10000} = \frac{5}{4} \text{ J}$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : [www.resonance.ac.in](http://www.resonance.ac.in) | E-mail : [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](http://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal


PAGE # 4

Activate Windows  
Go to Settings to activate Windows.




**Resonance** Educating for better tomorrow | **JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS**

10. Which of the following is correct graph between deviation ( $\delta$ ) and angle of incident 'i', if a ray of light passes through a prism :




Ans. (2)

Sol.  $\therefore \delta = i + e - A$ .




11. Pendulum bob of mass  $m$  and charge  $q$  is hinge between the plates of a parallel plate capacitor. The first half of space between the plates is filled with dielectric of dielectric constant  $K$  and another half is vacuum, where bob is hinged. What will be the angular deflection in string of pendulum with vertical if bob is in equilibrium. The voltage across capacitor is  $V$ .



Activate Windows  
Go to Settings to activate Windows.

11. Pendulum bob of mass  $m$  and charge  $q$  is hinge between the plates of a parallel plate capacitor. The first half of space between the plates is filled with dielectric of dielectric constant  $K$  and another half is vacuum, where bob is hinged. What will be the angular deflection in string of pendulum with vertical if bob is in equilibrium. The voltage across capacitor is  $V$ .



- (1)  $\tan^{-1} \frac{qvk}{(k+1)mgd}$     (2)  $\tan^{-1} \frac{2qvk}{(k+1)mgd}$     (3)  $\tan^{-1} \frac{3qvk}{(k+1)mgd}$     (4)  $\tan^{-1} \frac{4qv(k+1)}{(k)mgd}$

Ans. (2)

### Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](http://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal | PAGE # 5

Resonance® Educating for better tomorrow | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

Sol.  $(C_{eq})$  Equivalent capacitance =  $\frac{\epsilon_0 A}{\frac{d}{2k} + \frac{d}{2}} = \frac{2\epsilon_0 Ak}{d(k+1)}$

potential difference =  $V$

charge on capacitor =  $v \times C_{eq} = \frac{2\epsilon_0 AVk}{d(k+1)}$


Activate Windows  
Go to Settings to activate Windows.



Sol.  $(C_{eq})$  Equivalent capacitance =  $\frac{\epsilon_0 A}{\frac{d}{2k} + \frac{d}{2}} = \frac{2\epsilon_0 Ak}{d(k+1)}$


potential difference = V

charge on capacitor =  $v \times C_{eq} = \frac{2\epsilon_0 AVk}{d(k+1)}$



$\Rightarrow$  Electric field in vacuum =  $\frac{Q}{A\epsilon_0} \Rightarrow \frac{2Vk\epsilon_0 A}{d(k+1)A\epsilon_0}$

$E = \frac{2Vk}{d(k+1)}$



As bob is in equilibrium

$T \cos \theta = mg$

$T \sin \theta = qE$


$\tan \theta = \frac{qE}{mg}$

$\tan \theta = \frac{q2Vk}{mgd(k+1)}$

$\theta = \tan^{-1} \left( \frac{2qVk}{mgd(k+1)} \right)$

Activate Windows  
Go to Settings to activate Windows.

12. For the given gate circuit, choose the correct truth table ?



A	B	C
0	0	0
0	1	0
1	0	1
1	1	1

A	B	C
0	0	1
0	1	1
1	0	1
1	1	1

A	B	C
0	0	1
0	1	0
1	0	1
1	1	1

A	B	C
0	0	1
0	1	0
1	0	0
1	1	0



Ans. (3)

13. Write the approx. Value of plank's constant & permittivity constant :

- (1)  $6.6 \times 10^{-34}$  J-s,  $8.85 \times 10^{-12}$  F/m
- (2)  $6.6 \times 10^{-19}$  J-s,  $8.85 \times 10^{-12}$  F/m
- (3)  $6.6 \times 10^{-34}$  J-s,  $9 \times 10^9$  m/F
- (4) None

Ans. (1)

14. Two long wires having same current having radius a & b ( $a < b$ ) what will be the correct representation of magnetic field intensity v/s r :



Activate Windows  
Go to Settings to activate Windows.



1 | 1 | 1 | 1 | 1 | 0


**Ans. (3)**

13. Write the approx. Value of plank's constant & permittivity constant :

(1)  $6.6 \times 10^{-34}$  J-s,  $8.85 \times 10^{-12}$  F/m      (2)  $6.6 \times 10^{-19}$  J-s,  $8.85 \times 10^{-12}$  F/m  
 (3)  $6.6 \times 10^{-34}$  J-s,  $9 \times 10^9$  m/F      (4) None

**Ans. (1)**

14. Two long wires having same current having radius a & b ( $a < b$ ) what will be the correct representation of magnetic field intensity v/s r :



**Ans. (2)**

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : [www.resonance.ac.in](http://www.resonance.ac.in) | E-mail : [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](http://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal      PAGE # 7

Activate Windows  
Go to Settings to activate Windows.


Sol. B due to wire :

$$r < R \quad B = \frac{\mu_0 J r}{2}$$

where  $J = \frac{i}{\pi R^2}$

for wire of radius a  $J_1 = \frac{i}{\pi a^2}$

for wire of radius b  $J_2 = \frac{i}{\pi b^2}$



slope of OA = J

$$B_{\max} = \frac{\mu_0 J R}{2} = \frac{\mu_0 i R}{2\pi R^2} = \frac{\mu_0 i}{2\pi R}$$

we can see slope of wire of radius a ( $J_1$ ) > slope of wire of radius b ( $J_2$ )  
as  $b > a$  then  $B_{\max}$  for a >  $B_{\max}$  for b.

So Ans (B)


15. If Thomson model is considered and  $\alpha$  rays are bombard on this model then,  $\alpha$  rays will :

- (1) deflected at wide angle
- (2) reflected all at
- (3) will pass undeviated
- (4) all deflected at same angle

Ans. (3)

Sol. Theory based


16. A man crosses a river flowing with speed same as speed of man with respect to river. If man cross the river along path AB making an angle of  $30^\circ$  with the direction of river flow. Man starts swimming along line Ac, making an angle  $\theta$  with path AB. then value of  $\theta$  was ?





(3) will pass undeviated (4) all deflected at same angle  
**Ans. (3)**  
**Sol.** Theory based

16. A man crosses a river flowing with speed same as speed of man with respect to river. If man cross the river along path AB making an angle of  $30^\circ$  with the direction of river flow. Man starts swimming along line Ac, making an angle  $\theta$  with path AB. then value of  $\theta$  was ?



**Ans. (2)** (1)  $60^\circ$  (2)  $30^\circ$  (3)  $45^\circ$  (4)  $75^\circ$

### Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005  
Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222  
To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | CIN : U80302RJ2007PLC024029  
Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](http://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE # 8

Resonance® Educating for better tomorrow | JEE MAIN-2021 | DATE : 27-07-2021 (SHIFT-2) | PAPER-1 | MEMORY BASED | PHYSICS

**Sol.**  $\vec{V}_M = \vec{V}_{M,R} + \vec{V}_R$   
 $V_M$  should be along line AB  
 $\Rightarrow |\vec{V}_{M,R}| = |\vec{V}_R|$   
As  $|\vec{V}_{M,R}| = |\vec{V}_R|$


Activate Windows  
Go to Settings to activate Windows.

Sol.  $\vec{V}_M = \vec{V}_{M,R} + \vec{V}_R$

$V_M$  should be along line AB

$\Rightarrow |\vec{V}_{M,R}| = |\vec{V}_R|$

As  $|\vec{V}_{M,R}| = |\vec{V}_R|$



$\therefore V_M$  should be along angle bisector of angle between  $\vec{V}_{M,R}$  and  $\vec{V}_R$


$\therefore \theta = 30^\circ$

17. Two Carnot engines A and B are operated in series. The first one, A receives heat at  $T_1 (=600 \text{ K})$  and rejects to a reservoir at temperature  $T_2$ . The second engine B receives heat rejected by the first engine and, in turn, rejects to a heat reservoir at  $T_3 (=400\text{K})$ . Calculate the temperature  $T_2$  if the work outputs of the two engines are equal :

- (1) 500 k
- (2) 300 K
- (3) 600 K
- (4) 400 K

Ans. (1)

Sol.



$W = Q_1 - Q_2$

$W = Q_2 - Q_3$

$Q_1 - Q_2 = Q_2 - Q_3$

$Q_1 + Q_3 = 2Q_2$


$\frac{Q_1}{Q_2} + \frac{Q_3}{Q_2} = 2$

$T_1, T_3 = 2 \dots T = \frac{T_1 + T_3}{2} = 500 \text{ K}$

Activate Windows  
Go to Settings to activate Windows.



(1) 500 K (2) 300 K (3) 600 K (4) 400 K  
**Ans. (1)**  
**Sol.**



$$W = Q_1 - Q_2$$

$$W = Q_2 - Q_3$$

$$Q_1 - Q_2 = Q_2 - Q_3$$

$$Q_1 + Q_3 = 2Q_2$$

$$\frac{Q_1}{Q_2} + \frac{Q_3}{Q_2} = 2$$

$$\frac{T_1}{T_2} + \frac{T_3}{T_2} = 2 \Rightarrow T_2 = \frac{T_1 + T_3}{2} = 500 \text{ K.}$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : [www.resonance.ac.in](http://www.resonance.ac.in) | E-mail : [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | CIN : U80302RJ2007PLC024029

**Toll Free : 1800 258 5555** **7340010333** [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) [www.youtube.com/resowatch](https://www.youtube.com/resowatch) [blog.resonance.ac.in](http://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE # 9

Activate Windows  
Go to Settings to activate Windows.

**18.** An electron and a proton combined to form a H-atom in which electron is in 2<sup>nd</sup> excited state. From this excited state it releases a photon that strike a metal and emits an electron. The threshold wavelength for metal is 4000 Å , then find maximum possible kinetic energy for emitted electron.  
 (1) 9.0 eV                      (2) 3.1 eV                      (3) 7.1 eV                      (4) 13.6 eV

**Ans. (1)**

**Sol.**  $E = 13.6 \left[ \frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \text{ eV}$   
 $E = 13.6 \left[ \frac{1}{1} - \frac{1}{(3)^2} \right]$   
 $E = 13.6 \left[ \frac{8}{9} \right] = 12.1 \text{ eV}$   
 $KE_{\text{max}} = E - \frac{hc}{\lambda_0} = 12.1 - \frac{12400}{4000} \text{ eV}$   
 $= 12.1 - 3.1 \text{ eV} = 9.0 \text{ eV}$

**19.** Rain drops are falling vertically on earth with speed of 20m/s. Now wind start blowing horizontally with speed of 5m/s and a cyclist is moving with speed of 35 m/s opposite to the wind. Then find the velocity of rain with which rain hitting the cyclist.  
 (1)  $10\sqrt{5}$                       (2)  $22\sqrt{5}$                       (3)  $20\sqrt{5} \text{ m/s}$                       (4)  $\frac{22}{\sqrt{5}}$

**Ans. (3)**


**Sol.**  $V_{rg} = V_{rw} + V_{wg}$   
 $= -20\hat{j} + 5\hat{i}$   
 $V_{\text{rain,cy}} = V_{\text{rain,g}} - V_{\text{cy,g}}$   
 $= -20\hat{j} + 5\hat{i} - 35(-\hat{i}) = -20\hat{j} + 40(\hat{i})$   
 $V_{\text{rain,cy}} = \sqrt{20^2 + 40^2} = 20\sqrt{5} \text{ m/s}$

**20.** A beaker filled to the height of 12 cm was given, find the location where a hole should be made for max range.

Activate Windows  
Go to Settings to activate Windows.



20. A beaker filled to the height of 12 cm was given, find the location where a hole should be made for max range.



Ans. 6

### Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005


Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](http://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2021 Solution portal PAGE # 10

Sol.  $R = 2\sqrt{hh'}$



R is maximum when  $h' = h$

Activate Windows  
Go to Settings to activate Windows.

$\pi$  is maximum when  $\pi = \pi$   
 $\therefore h' = h = H/2$   
 $= 12/2 = 6$

21. A particle performing SHM is given by  $x = A \sin(\omega T + \phi)$ . At  $t = 0$ , particle is at  $x = 2$  and its velocity is  $2\omega$  then find amplitude :  
(1)  $2\sqrt{2}$                       (2)  $5\sqrt{2}$                       (3)  $4\sqrt{2}$                       (6)  $6\sqrt{2}$

Ans. (1)

Sol.  $2 = A \sin(0 + \phi)$

$\Rightarrow \sin \phi = \frac{2}{A}$

$\Rightarrow \cos \phi = \sqrt{1 - \frac{4}{A^2}} = \sqrt{\frac{A^2 - 4}{A^2}}$

$V = \frac{dx}{dt}$

$V = A\omega \cos(\omega t + \phi)$

$\Rightarrow 2\omega = A\omega \cos(0 + \phi)$

$\Rightarrow 2 = A \cos \phi$

$\Rightarrow A = \frac{2}{\cos \phi}$  ..... (i)

$\Rightarrow A = \frac{2}{\sqrt{\frac{A^2 - 4}{A^2}}}$

$\Rightarrow A = \frac{2A}{\sqrt{A^2 - 4}}$

$\Rightarrow \sqrt{A^2 - 4} = 2$

$\Rightarrow A = \pm \sqrt{8}$

### Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Activate Windows  
Go to Settings to activate Windows.



22. A body of mass  $m$  at rest starts moving along straight line by a machine delivering a constant power. Distance travelled by body in time  $t$  is :

- (1)  $\frac{4\sqrt{2p}}{3} \times t^{3/2}$       (2)  $\frac{2\sqrt{2p}}{3} \times t^{3/2}$       (3)  $\frac{\sqrt{2p}}{3} \times t^{3/2}$       (4)  $2\sqrt{3\frac{2p}{m}} \times t^{3/2}$

Ans. (2)

Sol. Energy supply =  $Pt$

in  $t$  sec

$$Pt = \frac{1}{2}mv^2$$

$$v = \sqrt{\frac{2pt}{m}}$$

$$\frac{dS}{dt} = \sqrt{\frac{2p}{m}}\sqrt{t}$$

$$\int_0^S dS = \sqrt{\frac{2p}{m}} \int_0^t t^{1/2} dt$$

$$S = \frac{2\sqrt{\frac{2p}{m}} t^{3/2}}{3}$$

$$t^{3/2} = \frac{3S}{2\sqrt{\frac{2p}{m}}}$$

$$S = \frac{2\sqrt{\frac{2p}{m}}}{3} \times t^{3/2}$$

23. In a communication, a message signal of amplitude 4 V is modulated with carrier signal of amplitude 12V, then find modulation index.

- (1)  $\frac{1}{2}$       (2)  $\frac{1}{3}$       (3)  $\frac{1}{4}$       (4)  $\frac{1}{6}$

Ans. (2)

Sol.  $A_{max} = 12 + 4 = 16$  V

Activate Windows  
Go to Settings to activate Windows.

24. 1 moles of an ideal gas undergoes adiabatic process, which increases the temperature from 27°C to 37°C. Gas is polyatomic has 4 vibrational modes of freedom. Find net work :

- (1) Work done by the gas 528 J                      (2) Work done on the gas 582 J  
 (3) Work done on the gas 382 J                      (4) Work done by the gas 382 J

Ans. (2)

Sol.  $f = 3 + 3 + (4 \times 2)$

$$f = 14$$

$$W = \frac{P_1 V_1 - P_2 V_2}{\gamma - 1} = \frac{nR(T_1 - T_2)}{\gamma - 1}$$

$$= \frac{1 \times 8.314 \times (-10)}{\left(\frac{8}{7} - 1\right)} \quad \left(\because \gamma = 1 + \frac{2}{f} = \frac{8}{7}\right)$$

$$= -582 \text{ J}$$