



Resonance[®]
Educating for better tomorrow

JEE

(Main)

PAPER-1 (B.E./B. TECH.)

2021

COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

Date: 25 July, 2021 (SHIFT-2) | TIME : (3.00 p.m. to 6.00 p.m)

Duration: 3 Hours | Max. Marks: 300






SUBJECT: PHYSICS

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2021 Solution portal

PART : PHYSICS

1 Value of force $F = A \sin(Bt) + C \cos(Dx)$ find dimension of $\frac{AB}{D}$

- (1) ML^3T^{-1} (2) ML^2T^{-3} (3) ML^1T^{-3} (4) ML^2T^3

Ans. (2)

Sol. Dimension of $A = MLT^{-2}$, $B = T^{-1}$, $D = L^{-1}$

$$\text{Dim} = \frac{AB}{D} = \frac{MLT^{-2}T^{-1}}{L^{-1}} = ML^2T^{-3}$$

2 Force is given by $F = (5y + 20) \hat{j}$ Find work done for moving particle from $y = 0$ to $y = 5$:

- (1) 162.5 J (2) 165 J (3) 132.5 J (4) 140.5 J

Ans. (1)

Sol. $w = \int F \cdot dy$

$$w = \int_0^5 (5y + 20) dy$$

$$= \left[\frac{5y^2}{2} + 20y \right]_0^5 \Rightarrow \frac{5 \times 25}{2} + 100 = 162.5 \text{ J}$$

3 A hot air balloon is ascending with constant velocity of 10 m/s. when balloon reaches a height of 75 m, a stone is dropped from balloon. what will be the height of balloon, when stone reaches earth?

- (1) 125 m. (2) 135 m. (3) 140 m. (4) 145 m.

Ans. (1)

Sol. For stone

$$75 = -10t + \frac{1}{2}gt^2$$

$$75 = -10t + 5t^2$$


$$t^2 - 2t - 15 = 0$$

$$t = 5 \text{ sec.}$$

Height of balloon

$$H = vt + 75$$

$$H = 10 \times 5 + 75 = 125 \text{ m.}$$



Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

4. Relation between position and time of a particle moving along straight line is given by $t = x + 3x^2$. Find acceleration of particle at $t = 10$ s

(1) $\frac{-5}{1331}$ (2) $\frac{6}{1331}$ (3) $\frac{-6}{1331}$ (4) $\frac{5}{1331}$

Ans. (3)

Sol. $t = x + 3x^2$ (1)

$$1 = \frac{dx}{dt} + 6x \frac{dx}{dt} \Rightarrow v = \frac{1}{(1+6x)}$$

$$0 = \frac{d^2x}{dt^2} + 6 \left(x \frac{d^2x}{dt^2} + \left(\frac{dx}{dt} \right)^2 \right)$$

$$0 = a + 6xa + 6v^2$$

$$a = \frac{-6v^2}{(1+6x)} \dots(2)$$

$$a = -\frac{6}{(1+6x)^3}$$

From equation(1)

$$10 = x + 3x^2$$

$$3x^2 + x - 10 = 0$$

$$3x^2 + 6x - 5x - 10 = 0$$

$$3x(x + 2) - 5(x + 2)$$

$$(3x - 5)(x + 2) \Rightarrow x = \frac{5}{3}$$

From equation (2)

$$a = \frac{-6}{\left(1 + 6 \times \frac{5}{3}\right)^3} = \frac{-6}{1331}$$

5. Two particle of same mass & charges Q_1 and Q_2 are moving perpendicular to an uniform magnetic field where

the ratio of charges is $\frac{Q_1}{Q_2} = \frac{1}{2}$ and ratio of velocities is $\frac{V_1}{V_2} = \frac{3}{2}$ then find the ratio of the radius $\frac{R_1}{R_2}$:

(1) 2 : 1 (2) 3 : 1 (3) 4 : 1 (4) 1 : 1

Ans. (2)

Sol. Given

$$\frac{Q_1}{Q_2} = \frac{1}{2} \quad \& \quad \frac{V_1}{V_2} = \frac{3}{2}$$

$$R = \frac{mv}{qB}$$

$$\frac{R_1}{R_2} = \frac{V_1}{V_2} \times \frac{Q_2}{Q_1} = \frac{3}{2} \times \frac{2}{1} = \frac{3}{1}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

6. A particle performing SHM with amplitude A. Find the ratio of kinetic energy and total energy when particle is at A/2
- (1) $\frac{3}{4}$ (2) $\frac{2}{3}$ (3) $\frac{4}{3}$ (4) $\frac{1}{2}$

Ans. (1)

Sol. $V_{A/2} = \omega\sqrt{A^2 - x^2}$

$$= \omega\sqrt{A^2 - \left(\frac{A}{2}\right)^2} = \omega\left(\frac{\sqrt{3}}{2}A\right)$$

$$= \frac{\sqrt{3}}{2}V_{\max}$$

$$KE = \frac{1}{2}m\left(\frac{\sqrt{3}}{2}V_{\max}\right)^2$$

$$TE = \frac{1}{2}m(V_{\max})^2$$

$$\frac{KE}{TE} = \frac{3}{4}$$

Ans.

7. In photoelectric effect stopping potential is $3V_0$ for incident wave length λ_0 and stopping potential V_0 for incident wavelength $2\lambda_0$. Find threshold wavelength.
- (1) $3\lambda_0$ (2) $2\lambda_0$ (3) $4\lambda_0$ (4) $8\lambda_0$

Ans. (3)

Sol. $KE = h\nu - W$

$$eV = \frac{hc}{\lambda} - W$$

For first case

$$e(3V_0) = \frac{hc}{\lambda_0} - W \quad \dots(i)$$

For second case

$$eV_0 = \frac{hc}{2\lambda_0} - W \quad \dots(ii)$$

From equation (i) and (ii)

$$W = \frac{hc}{4\lambda_0}$$

For λ_{th}

$$W = \frac{hc}{\lambda_{th}}$$

$$\Rightarrow \frac{hc}{4\lambda_0} = \frac{hc}{\lambda_{th}} \Rightarrow \lambda_{th} = 4\lambda_0$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

8. Efficiency of heat engine is $\eta = 1/6$. If temperature of sink is decreased by 62K, then efficiency becomes 1/3. Find temperature of source :

(1) 372K (2) 272K (3) 350K (4) 450K

Ans. (1)

Sol. $\eta = \left(1 - \frac{T_2}{T_1}\right)$

$$\frac{T_2}{T_1} = 1 - \eta = 1 - \frac{1}{6} \quad \dots(1)$$

$$\frac{T_2 - 62}{T_1} = 1 - \frac{1}{3} \quad \dots(2)$$

Equation $\frac{(1)}{(2)}$:

$$\Rightarrow \frac{T_2}{T_2 - 62} = \frac{5}{6} \times \frac{3}{2} = \frac{5}{4}$$

$$\Rightarrow T_2 = 5 \times 62$$

From eq. (1)

$$T_1 = \frac{T_2}{1 - \eta} = \frac{5 \times 62}{1 - \frac{1}{6}} = 5 \times 62 \times \frac{6}{5} = 372\text{K}$$

9. Activity of an element x becomes 1/8 of initial in 30 years. Find half-life :

(1) 10 Year. (2) 12 Year (3) 15 Year (4) 17 Year

Ans. (1)

Sol. $A = A_0 e^{-\lambda t}$

For half life

$$A/2 = A e^{-\lambda t_{1/2}}$$

$$\frac{1}{2} = e^{-\lambda t_{1/2}} \quad \dots(1)$$

$$\text{Given } 1/8 = e^{-\lambda 30} \quad \dots(2)$$

Solving (1) and (2)

$$e^{-3\lambda t_{1/2}} = e^{-\lambda 30}$$

$$T_{1/2} = 10 \text{ Yrs.}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

10. If De-Broglie wavelengths of photon and electron are equal, what will be the ratio of kinetic energy of electron and energy of photon? Given that velocity of electron is v and velocity of light is c :

(1) $\frac{2v}{c}$ (2) $\frac{v}{2c}$ (3) $\frac{3v}{c}$ (4) $\frac{c}{3v}$

Ans. (2)

Sol. De-Broglie wavelength is given by $\lambda = \frac{h}{p}$

$$KE_{pn} = mc^2 = pc \quad \dots(1)$$

$$KE_e = \frac{1}{2} mv^2 = \frac{pv}{2} \quad \dots(2)$$


$$\frac{KE_e}{KE_{pn}} = \frac{pv/2}{pc} = \frac{v}{2c}$$

11. A square loop of total resistance 16Ω . If a battery of $2V$ and 1Ω internal resistance is connected across one of its side then find potential difference across its diagonal :

(1) $1V$ (2) $2V$ (3) $3V$ (4) $4V$


Ans. (1)

Sol.



$$V_{AB} = ??$$

$$R_{eq} = \frac{12 \times 4}{12 + 4} = 3$$



$$i = \frac{2}{3+1} = \frac{1}{2} \text{ A}$$

$$i_2 = \frac{r_2}{r_2 + r_1} i = \frac{1}{3+1} \times \frac{1}{2} = \frac{1}{8}$$

$$V_{AB} = \frac{1}{8} \times 8 = 1V$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

12. \vec{A} and \vec{B} are two vectors such that $|\vec{A}| = 2$ and $|\vec{B}| = 5$. If $|\vec{A} \times \vec{B}| = 8$, then $|\vec{A} \cdot \vec{B}| = ?$

- (1) 2 (2) 6 (3) 7 (4) 9

Ans. (2)

Sol. $|\vec{A} \times \vec{B}| = |\vec{A}||\vec{B}|\sin\theta$

$$\Rightarrow 10\sin\theta = 8$$

$$\sin\theta = \frac{4}{5}$$

$$\text{Now } |\vec{A} \cdot \vec{B}| = |\vec{A}||\vec{B}|\cos\theta = 10 \times \frac{3}{5} = 6$$

13. Find significant figure for the value 0.00346.

- (1) 5 (2) 4 (3) 3 (4) 2

Ans. (3)

Sol. There are 3 non zero digit after the decimal point so significant number is 3.

0.00346

14. For a prism, if angle of minimum deviation is equal to angle of prism. If refractive index of prism material is μ . Then angle of prism should be?

- (1) $2\sin^{-1}\left(\frac{\mu}{2}\right)$ (2) $2\cos^{-1}\left(\frac{\mu}{2}\right)$ (3) $3\cos^{-1}\left(\frac{\mu}{2}\right)$ (4) $3\sin^{-1}\left(\frac{\mu}{2}\right)$

Ans. (2)

Sol. $\mu = \frac{\sin\left(\frac{A + \delta_m}{2}\right)}{\sin\left(\frac{A}{2}\right)}$

$$\mu = \frac{\sin A}{\sin A/2}$$

$$\mu = 2\cos\frac{A}{2}$$






$$A = 2\cos^{-1}\left(\frac{\mu}{2}\right)$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in

15. A photon of wavelength 500 nm falls on a metal surface of work function 1.3eV. An electron releases from metal moved in a perpendicular magnetic field. In a circular path of radius 30 cm. Then the magnitude of magnetic field will be ?

- (1) 12.2 μT (2) 10.2 μT (3) 8.2 μT (4) 6.2 μT

Ans. (1)

Sol. $\frac{hc}{\lambda} = \phi + KE_{\max}$

$$\frac{1240}{500} = 1.3 + KE_{\max}$$

$$KE_{\max} = 1.18 \text{ eV}$$

$$\text{Now } R = \frac{mv}{qB} = \frac{\sqrt{2mKE}}{qB}$$

$$B = \frac{\sqrt{2mKE}}{qR}$$


$$B = \frac{\sqrt{2 \times 9.1 \times 10^{-31} \times 1.18 \times 1.6 \times 10^{-19}}}{1.6 \times 10^{-19} \times 30 \times 10^{-2}}$$

$$B = 0.122 \times 10^{-4}$$

$$B = 12.2 \times 10^{-6}$$

$$\text{i.e., } B = 12.2 \mu\text{T}$$

16. Two electric dipole \vec{P}_1 and \vec{P}_2 are kept as shown in figure. Net electric field at point S is E makes an angle 37° with \vec{P}_1 then find the ratio of $|\vec{P}_1|$ and $|\vec{P}_2|$.



- (1) $\frac{3}{2}$ (2) $\frac{1}{2}$ (3) $\frac{2}{3}$ (4) $\frac{3}{4}$

Ans. (3)

Resonance Eduventures Ltd.


Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [7340010333](https://www.facebook.com/ResonanceEdu) | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

Sol.



Electric field due to \vec{P}_1 at axis point S

$$E_{\text{axis}} = \frac{2KP_1}{r^3}$$

$$\Rightarrow E \cos 37^\circ = \frac{2KP_1}{r^3} \quad \dots(1)$$

Electric field due to \vec{P}_2 at perpendicular bisector at point S.


$$E_{\perp} = \frac{KP_2}{r^3}$$

$$\Rightarrow E \sin 37^\circ = \frac{KP_2}{r^3} \quad \dots(2)$$

$$\therefore \frac{\frac{2KP_1}{r^3}}{\frac{KP_2}{r^3}} = \frac{E \cos 37^\circ}{E \sin 37^\circ}$$

$$\Rightarrow \frac{2P_1}{P_2} = \frac{4}{3} \Rightarrow \frac{P_1}{P_2} = \frac{2}{3}$$

17. Power in both the given circuit are same then find angular frequency of AC source.



(1) 200

(2) 300

(3) 400

(4) 500

Ans. (4)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

Sol. $P_1 = P_2$

$$\left(\frac{V^2}{R}\right)_1 = \left(\frac{V^2}{Z}\right)_2 \Rightarrow R = Z$$

$$R = \sqrt{\left(\omega L - \frac{1}{\omega C}\right)^2 + R^2}$$

$$10 = \sqrt{\left(\omega L - \frac{1}{\omega C}\right)^2 + R^2}$$


$$100 = \left[\omega(0.1) - \frac{1}{\omega(40 \times 10^{-6})}\right]^2 + 100$$

$$\omega^2(0.1) = \frac{1}{40 \times 10^{-6}}$$

$$\omega^2 = \frac{1}{4} \times 10^6$$

$$\omega = 500$$

18. For the given circuit, find the potential drop across 2Ω resistance ?



The wire AB is of length 10 cm, and its resistance is $1\Omega/\text{cm}$. Point D is mid-point of wire AB.

(1) 2.44 V


(2) 4.44 V

(3) 3.44 V

(4) 10.44 V

Ans. (2)

Sol.



$$V_{2\Omega} = \frac{20}{\frac{10}{7} + 5} \times \frac{10}{7}$$

$$V_{2\Omega} = 4.44 \text{ V}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

19. Mass of a planet is double the mass of earth. Both the planet have same mass density. A body has weight W on surface of earth, then weight of the same body on surface of planet ?

(1) $2^{2/3} W$ (2) $2^{1/3} W$ (3) W (4) $3^{1/2} W$

Ans. (2)

Sol. $2M_E = M_P$

$$2\rho \times \frac{4}{3} R_E^3 = \rho \times \frac{4}{3} \pi R_P^3 \text{ (same density)}$$

$$R_P = 2^{1/3} R_E$$

$$g_P = \frac{GM_P}{R_P^2} \text{ (acceleration due to gravity)}$$

$$g_P = \frac{G2M_E}{(2^{1/3} R_E)^2} = \frac{G2M_E}{2^{2/3} R_E^2}$$

$$g_P = 2^{1/3} g_e$$

weight on planet = $2^{1/3}$ weight on earth

$$W_P = 2^{1/3} W$$

20. A force $\vec{F} = 40\hat{i} + 10\hat{j}$ is applied on a stationary object of mass 5kg. What will be the position of object after 10s, if initially object was at origin ?

(1) (200, 100) (2) (400, 400) (3) (400, 100) (4) (100, 100)

Ans. (3)


Sol. $\vec{a} = 8\hat{i} + 2\hat{j}$

$$\vec{s} = \vec{u}t + \frac{1}{2} \vec{a}t^2$$

$$\vec{s} = \frac{1}{2} (8\hat{i} + 2\hat{j}) \times 100$$

$$\vec{s} = 400\hat{i} + 100\hat{j}$$

21. An AC Source with $V_{\max} = 200 \text{ V}$ and $f = 50 \text{ Hz}$ connected across 10Ω resistance. Find the time in which source voltage changes from maximum to rms value.



(1) $\frac{1}{200} \text{ s}$ (2) $\frac{1}{400} \text{ s}$ (3) $\frac{1}{300} \text{ s}$ (4) $\frac{1}{500} \text{ s}$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resowatch blog.resonance.ac.in

23. For two vector \vec{X} and \vec{Y} , $|\vec{X}| = |\vec{Y}|$ and $|\vec{X} - \vec{Y}| = n |\vec{X} + \vec{Y}|$. Then find angle between \vec{X} and \vec{Y} ?

(1) $\cos^{-1} \frac{1-n^2}{1+n^2}$ (2) $\cos^{-1} \frac{1+n^2}{1-n^2}$ (3) $\cos^{-1} \frac{2-n^2}{2+n^2}$ (4) $\cos^{-1} \frac{2+n^2}{2-n^2}$

Ans. (1)

Sol. $|\vec{X} - \vec{Y}| = n |\vec{X} + \vec{Y}|$

$$|\vec{X}|^2 + |\vec{Y}|^2 - 2|\vec{X}||\vec{Y}|\cos\theta = n^2 [|\vec{X}|^2 + |\vec{Y}|^2 + 2|\vec{X}||\vec{Y}|\cos\theta]$$

As $|\vec{X}| = |\vec{Y}|$

$$2|\vec{X}|^2 - 2|\vec{X}|^2 \cos\theta = 2n^2|\vec{X}|^2 + 2n^2|\vec{X}|^2 \cos\theta$$

$$1 - \cos\theta = n^2 + n^2 \cos\theta$$

$$\cos\theta = \frac{1-n^2}{1+n^2}$$

$$\theta = \cos^{-1} \frac{1-n^2}{1+n^2}$$

24 Find energy required to break an Aluminium nucleus into its constituent nucleons.

($m_n = 1.00867$ u, $m_p = 1.00783$ u, $m_{Al} = 26.98154$ u)

(1) 225 MeV (2) 230 MeV (3) 235 MeV (4) 245 MeV

Ans. (1)

Sol. Binding Energy = $\Delta m C^2$

$$\Delta m = [13 \times 1.00783 + 14 \times 1.00867 - 26.98154]$$

$$= [13.10179 + 14.12138 - 26.98154] = 0.24163$$

$$\therefore \text{B.E} = 0.24163 C^2 \times 931 \text{ MeV}/C^2$$

$$= 224.95 \text{ MeV} \approx 225 \text{ MeV.}$$

25. A Cell of Voltage ' V_0 ' is connected across a capacitor of capacitance 'C'. Now the space between the plates is filled with a material of dielectric constant K. Find the ratio of charge appear on the plates of capacitor before and after filling.

(1) 1 : K (2) K : 1 (3) 2 K : 1 (4) K : 2

Ans. (1)

Resonance Eduventures Ltd.


Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

Sol.



$$Q_1 = CV$$

$$Q_2 = KCV$$

$$\frac{Q_1}{Q_2} = \frac{1}{K}$$

26. Pure Si at room temperature has equal electron (n_e) and hole (n_h) concentration of $1.5 \times 10^{16} \text{ m}^{-3}$. Doping by indium increases n_h to $3 \times 10^{22} \text{ m}^{-3}$. Calculate n_e in the doped Si.

- (1) $7.5 \times 10^9 \text{ m}^{-3}$ (2) $6.5 \times 10^9 \text{ m}^{-3}$ (3) $7.5 \times 10^8 \text{ m}^{-3}$ (4) $7.5 \times 10^7 \text{ m}^{-3}$

Ans. (1)

Sol. For a doped semi-conductor in thermal equilibrium

$$n_e n_h = n_i^2$$

$$\Rightarrow n_e = \frac{n_i^2}{n_h} = \frac{(1.5 \times 10^{16})^2}{3 \times 10^{22}} = 7.5 \times 10^9 \text{ m}^{-3}$$

27. A particle starts from rest and moves with a variable acceleration $a = \alpha t + \beta t^2$, where α and β are positive constants. Find the distance covered by particle in $t = 1 \text{ sec}$ to $t = 2 \text{ sec}$?

- (1) $\frac{11}{6}\alpha + \frac{15}{12}\beta$ (2) $\frac{7}{6}\alpha + \frac{17}{12}\beta$ (3) $\frac{7}{6}\alpha + \frac{15}{12}\beta$ (4) $\frac{1}{3}\alpha + \frac{15}{12}\beta$

Ans. (3)

Sol. $\int_0^v dv = \int_0^t a dt$

$$v = \frac{\alpha t^2}{2} + \frac{\beta t^3}{3}$$

Now

$$\int_0^s ds = \int_1^2 v dt$$

$$s = \left[\frac{\alpha t^3}{6} + \frac{\beta t^4}{12} \right]_1^2 \Rightarrow s = \frac{7}{6}\alpha + \frac{15}{12}\beta$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

28. A carrier frequency of 1 MHz and peak value of 10 v is amplitude modulated with a signal frequency of 10 KHz with peak value of 0.5 v. Find modulation index.

- (1) 0.02 (2) 0.03 (3) 0.04 (4) 0.05

Ans. (4)

Sol. $A_{\max.} = 10 + 0.5 = 10.5$

$$A_{\min} = 10 - 0.5 = 9.5$$


$$m_a = \frac{A_{\max.} - A_{\min.}}{A_{\max.} + A_{\min.}} = \frac{10.5 - 9.5}{10.5 + 9.5} = 0.05$$

29. Two soap bubbles of radius r_1 and r_2 in vacuum are combined isothermally to form a new bubble. Find the radius of this new bubble ?

- (1) $\sqrt{r_1^2 + r_2^2}$ (2) $\sqrt{r_1^2 - r_2^2}$ (3) $\sqrt{\frac{r_1 r_2}{r_1 + r_2}}$ (4) $\sqrt{\frac{r_1 r_2}{r_1 - r_2}}$

Ans. (1)

Sol.



By surface energy conservation

$$\sigma A_1 + \sigma A_2 = \sigma A$$

$$\sigma [2 \times 4\pi r_1^2] + \sigma [2 \times 4\pi r_2^2] = \sigma [2 \times 4\pi r^2]$$

$$r_1^2 + r_2^2 = r^2$$

$$r = \sqrt{r_1^2 + r_2^2}$$

Resonance Eduventures Ltd.


Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | facebook.com/ResonanceEdu | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

30. A ray is incident on a slab of refractive index $\frac{5}{4}$ at an angle θ as shown in figure. Find maximum angle θ . So that TIR occur at surface AD.



(1) $\sin^{-1} \frac{\sqrt{5}}{3}$

(2) $\sin^{-1} \frac{\sqrt{3}}{2}$

(3) $\sin^{-1} \frac{3}{4}$

(4) $\sin^{-1} \frac{\sqrt{5}}{4}$

Ans. (3)

Sol. $1 \times \sin\theta = \frac{5}{4} \sin(90 - C)$

$$\sin\theta = \frac{5}{4} \cos C$$

but $\sin C = \frac{1}{\mu} = \frac{4}{5}$

$$\cos C = \frac{3}{5}$$

$$\sin\theta = \frac{5}{4} \times \frac{3}{5} = \frac{3}{4}$$

For T.I.R. $\sin\theta < \frac{3}{4}$

$$\theta = \sin^{-1} \frac{3}{4}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in



**RESULT: JEE (Advanced),
JEE (Main), NEET**

HIGHEST No. of Classroom Selections
in JEE (Advanced) 2020 from any Institute of Kota

5 AIRs in TOP-50 in JEE (Adv.) 2020 from Classroom

**AIR-2
(GEN-EWS)**
AIR-15
**DHANANJAY
KEJRIWAL**
With us Since Class 9th

**Zonal Topper
IIT-Kharagpur**
AIR-25
**SAMARTH
AGARWAL**
With us Since Class 11th

**2nd Rank in
IIT-Kharagpur Zone**
AIR-29
**SANKALP
PARASHAR**
With us Since Class 11th

AIR-30
**AARYAN K.
GUPTA**
With us Since Class 9th

AIR-41
**UTKARSH P.
SINGH**
With us Since Class 10th

Total Selections in JEE (Advanced) 2020

4505

Classroom: 3441 | Distance: 1064

Eligible for JEE (Advanced) Through JEE (Main) 2020

14755

Classroom: 11047 | Distance: 3708

NEET 2020

2646

Classroom: 1833 | Distance: 813

ADMISSION OPEN for Session 2021-22

ONLINE + OFFLINE PROGRAMS

CLASS 11, 12 & 12+

Target: JEE (Main+Adv.) | JEE (Main) | NEET

Scholarship Upto 90%*

Toll Free: 1800 258 5555 | Visit us: www.resonance.ac.in

