

FINAL JEE-MAIN EXAMINATION – JULY, 2021
(Held On Tuesday 27th July, 2021)
TIME : 9 : 00 AM to 12 : 00 NOON
MATHEMATICS
TEST PAPER WITH ANSWER
SECTION-A

1. If the mean and variance of the following data:
 6, 10, 7, 13, a, 12, b, 12
 are 9 and $\frac{37}{4}$ respectively, then $(a - b)^2$ is equal to:

(1) 24 (2) 12 (3) 32 (4) 16

Official Ans. by NTA (4)

2. The value of $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \frac{(2j-1) + 8n}{(2j-1) + 4n}$ is equal to :

 (1) $5 + \log_e \left(\frac{3}{2}\right)$ (2) $2 - \log_e \left(\frac{2}{3}\right)$

 (3) $3 + 2 \log_e \left(\frac{2}{3}\right)$ (4) $1 + 2 \log_e \left(\frac{3}{2}\right)$
Official Ans. by NTA (4)

3. Let $\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = -\hat{i} + 2\hat{j} + 3\hat{k}$. Then the vector product $(\vec{a} + \vec{b}) \times \left((\vec{a} \times ((\vec{a} - \vec{b}) \times \vec{b})) \times \vec{b} \right)$ is equal to :

 (1) $5(34\hat{i} - 5\hat{j} + 3\hat{k})$ (2) $7(34\hat{i} - 5\hat{j} + 3\hat{k})$

 (3) $7(30\hat{i} - 5\hat{j} + 7\hat{k})$ (4) $5(30\hat{i} - 5\hat{j} + 7\hat{k})$
Official Ans. by NTA (2)

4. The value of the definite integral

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{dx}{(1 + e^{\cos x})(\sin^4 x + \cos^4 x)}$$

is equal to :

 (1) $-\frac{\pi}{2}$ (2) $\frac{\pi}{2\sqrt{2}}$ (3) $-\frac{\pi}{4}$ (4) $\frac{\pi}{\sqrt{2}}$
Official Ans. by NTA (2)

5. Let C be the set of all complex numbers. Let

$$S_1 = \{z \in C \mid |z - 3 - 2i|^2 = 8\},$$

$$S_2 = \{z \in C \mid \operatorname{Re}(z) \geq 5\} \text{ and}$$

$$S_3 = \{z \in C \mid |z - \bar{z}| \geq 8\}.$$

 Then the number of elements in $S_1 \cap S_2 \cap S_3$ is equal to

(1) 1 (2) 0 (3) 2 (4) Infinite

Official Ans. by NTA (1)

6. If the area of the bounded region
 $R = \left\{ (x, y) : \max\{0, \log_e x\} \leq y \leq 2^x, \frac{1}{2} \leq x \leq 2 \right\}$

 is, $\alpha(\log_e 2)^{-1} + \beta(\log_e 2) + \gamma$, then the value of $(\alpha + \beta - 2\gamma)^2$ is equal to :

(1) 8 (2) 2 (3) 4 (4) 1

Official Ans. by NTA (2)

7. A ray of light through (2,1) is reflected at a point P on the y-axis and then passes through the point (5, 3). If this reflected ray is the directrix of an ellipse with eccentricity $\frac{1}{3}$ and the distance of the

 nearer focus from this directrix is $\frac{8}{\sqrt{53}}$, then the

equation of the other directrix can be:

 (1) $11x + 7y + 8 = 0$ or $11x + 7y - 15 = 0$

 (2) $11x - 7y - 8 = 0$ or $11x + 7y + 15 = 0$

 (3) $2x - 7y + 29 = 0$ or $2x - 7y - 7 = 0$

 (4) $2x - 7y - 39 = 0$ or $2x - 7y - 7 = 0$
Official Ans. by NTA (3)

8. If the coefficients of x^7 in $\left(x^2 + \frac{1}{bx}\right)^{11}$ and x^{-7} in

 $\left(x - \frac{1}{bx^2}\right)^{11}$, $b \neq 0$, are equal, then the value of b

is equal to:

(1) 2 (2) -1 (3) 1 (4) -2

Official Ans. by NTA (3)

9. The compound statement $(P \vee Q) \wedge (\sim P) \Rightarrow Q$ is equivalent to:

 (1) $P \vee Q$ (2) $P \wedge \sim Q$

 (3) $\sim(P \Rightarrow Q)$ (4) $\sim(P \Rightarrow Q) \Leftrightarrow P \wedge \sim Q$
Official Ans. by NTA (4)

10. If $\sin \theta + \cos \theta = \frac{1}{2}$, then

 $16(\sin(2\theta) + \cos(4\theta) + \sin(6\theta))$ is equal to:

(1) 23 (2) -27 (3) -23 (4) 27

Official Ans. by NTA (3)

11. Let $A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix}$. If $A^{-1} = \alpha I + \beta A$, $\alpha, \beta \in \mathbf{R}$, I is a 2×2 identity matrix, then $4(\alpha - \beta)$ is equal to :
- (1) 5 (2) $\frac{8}{3}$ (3) 2 (4) 4

Official Ans. by NTA (4)

12. Let $f : \left(-\frac{\pi}{4}, \frac{\pi}{4}\right) \rightarrow \mathbf{R}$ be defined as

$$f(x) = \begin{cases} (1 + |\sin x|)^{\frac{3a}{|\sin x|}}, & -\frac{\pi}{4} < x < 0 \\ b, & x = 0 \\ e^{\cot 4x / \cot 2x}, & 0 < x < \frac{\pi}{4} \end{cases}$$

If f is continuous at $x = 0$, then the value of $6a + b^2$ is equal to:

- (1) $1 - e$ (2) $e - 1$ (3) $1 + e$ (4) e

Official Ans. by NTA (3)

13. Let $y = y(x)$ be solution of the differential equation

$$\log_e \left(\frac{dy}{dx} \right) = 3x + 4y, \text{ with } y(0) = 0.$$

If $y \left(-\frac{2}{3} \log_e 2 \right) = \alpha \log_e 2$, then the value of α is equal to:

- (1) $-\frac{1}{4}$ (2) $\frac{1}{4}$ (3) 2 (4) $-\frac{1}{2}$

Official Ans. by NTA (1)

14. Let the plane passing through the point $(-1, 0, -2)$ and perpendicular to each of the planes $2x + y - z = 2$ and $x - y - z = 3$ be $ax + by + cz + 8 = 0$. Then the value of $a + b + c$ is equal to:

- (1) 3 (2) 8 (3) 5 (4) 4

Official Ans. by NTA (4)

15. Two tangents are drawn from the point $P(-1, 1)$ to the circle $x^2 + y^2 - 2x - 6y + 6 = 0$. If these tangents touch the circle at points A and B , and if D is a point on the circle such that length of the segments AB and AD are equal, then the area of the triangle ABD is equal to:

- (1) 2 (2) $(3\sqrt{2} + 2)$
 (3) 4 (4) $3(\sqrt{2} - 1)$

Official Ans. by NTA (3)

16. Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be a function such that $f(2) = 4$ and $f'(2) = 1$. Then, the value of $\lim_{x \rightarrow 2} \frac{x^2 f(2) - 4f(x)}{x - 2}$ is equal to :

- (1) 4 (2) 8 (3) 16 (4) 12

Official Ans. by NTA (4)

17. Let P and Q be two distinct points on a circle which has center at $C(2, 3)$ and which passes through origin O . If OC is perpendicular to both the line segments CP and CQ , then the set $\{P, Q\}$ is equal to

- (1) $\{(4, 0), (0, 6)\}$
 (2) $\{(2 + 2\sqrt{2}, 3 - \sqrt{5}), (2 - 2\sqrt{2}, 3 + \sqrt{5})\}$
 (3) $\{(2 + 2\sqrt{2}, 3 + \sqrt{5}), (2 - 2\sqrt{2}, 3 - \sqrt{5})\}$
 (4) $\{(-1, 5), (5, 1)\}$

Official Ans. by NTA (4)

18. Let α, β be two roots of the equation

$$x^2 + (20)^{1/4} x + (5)^{1/2} = 0. \text{ Then } \alpha^8 + \beta^8 \text{ is equal to}$$

- (1) 10 (2) 100 (3) 50 (4) 160

Official Ans. by NTA (3)

19. The probability that a randomly selected 2-digit number belongs to the set $\{n \in \mathbf{N} : (2^n - 2) \text{ is a multiple of } 3\}$ is equal to

- (1) $\frac{1}{6}$ (2) $\frac{2}{3}$ (3) $\frac{1}{2}$ (4) $\frac{1}{3}$

Official Ans. by NTA (3)

20. Let

$$A = \{(x, y) \in \mathbf{R} \times \mathbf{R} \mid 2x^2 + 2y^2 - 2x - 2y = 1\},$$

$$B = \{(x, y) \in \mathbf{R} \times \mathbf{R} \mid 4x^2 + 4y^2 - 16y + 7 = 0\} \text{ and}$$

$$C = \{(x, y) \in \mathbf{R} \times \mathbf{R} \mid x^2 + y^2 - 4x - 2y + 5 \leq r^2\}.$$

Then the minimum value of $|r|$ such that $A \cup B \subseteq C$ is equal to

- (1) $\frac{3 + \sqrt{10}}{2}$ (2) $\frac{2 + \sqrt{10}}{2}$
 (3) $\frac{3 + 2\sqrt{5}}{2}$ (4) $1 + \sqrt{5}$

Official Ans. by NTA (3)

SECTION-B

1. For real numbers α and β , consider the following system of linear equations :

$$x + y - z = 2, x + 2y + \alpha z = 1, 2x - y + z = \beta.$$

If the system has infinite solutions, then $\alpha + \beta$ is equal to _____.

Official Ans. by NTA (5)

2. Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, \vec{b} and $\vec{c} = \hat{j} - \hat{k}$ be three vectors such that $\vec{a} \times \vec{b} = \vec{c}$ and $\vec{a} \cdot \vec{b} = 1$. If the length of projection vector of the vector \vec{b} on the vector $\vec{a} \times \vec{c}$ is l , then the value of $3l^2$ is equal to _____.

Official Ans. by NTA (2)

3. If $\log_3 2, \log_3(2^x - 5), \log_3\left(2^x - \frac{7}{2}\right)$ are in an arithmetic progression, then the value of x is equal to _____.

Official Ans. by NTA (3)

4. Let the domain of the function $f(x) = \log_4\left(\log_5\left(\log_3(18x - x^2 - 77)\right)\right)$ be (a, b) .

Then the value of the integral

$$\int_a^b \frac{\sin^3 x}{(\sin^3 x + \sin^3(a + b - x))} dx \text{ is equal to } \underline{\hspace{2cm}}.$$

Official Ans. by NTA (1)

5. Let
- $$f(x) = \begin{vmatrix} \sin^2 x & -2 + \cos^2 x & \cos 2x \\ 2 + \sin^2 x & \cos^2 x & \cos 2x \\ \sin^2 x & \cos^2 x & 1 + \cos 2x \end{vmatrix}, x \in [0, \pi]$$

Then the maximum value of $f(x)$ is equal to _____.

Official Ans. by NTA (6)

6. Let $F : [3, 5] \rightarrow \mathbf{R}$ be a twice differentiable function on $(3, 5)$ such that

$$F(x) = e^{-x} \int_3^x (3t^2 + 2t + 4F'(t)) dt.$$

If $F'(4) = \frac{\alpha e^\beta - 224}{(e^\beta - 4)^2}$, then $\alpha + \beta$ is equal to _____.

Official Ans. by NTA (16)

7. Let a plane P pass through the point $(3, 7, -7)$ and contain the line, $\frac{x-2}{-3} = \frac{y-3}{2} = \frac{z+2}{1}$. If distance of the plane P from the origin is d , then d^2 is equal to _____.

Official Ans. by NTA (3)

8. Let $S = \{1, 2, 3, 4, 5, 6, 7\}$. Then the number of possible functions $f : S \rightarrow S$ such that $f(m \cdot n) = f(m) \cdot f(n)$ for every $m, n \in S$ and $m \cdot n \in S$ is equal to _____.

Official Ans. by NTA (490)

9. If $y = y(x)$, $y \in \left[0, \frac{\pi}{2}\right)$ is the solution of the differential equation

$$\sec y \frac{dy}{dx} - \sin(x+y) - \sin(x-y) = 0, \text{ with } y(0) = 0,$$

then $5y'\left(\frac{\pi}{2}\right)$ is equal to _____.

Official Ans. by NTA (2)

10. Let $f : [0, 3] \rightarrow \mathbf{R}$ be defined by $f(x) = \min \{x - [x], 1 + [x] - x\}$ where $[x]$ is the greatest integer less than or equal to x . Let P denote the set containing all $x \in [0, 3]$ where f is discontinuous, and Q denote the set containing all $x \in (0, 3)$ where f is not differentiable. Then the sum of number of elements in P and Q is equal to _____.

Official Ans. by NTA (5)