

2.

FINAL JEE-MAIN EXAMINATION - JULY, 2021

(Held On Tuesday 27th July, 2021)

TEST PAPER WITH ANSWER

TIME: 3:00 PM to 6:00 PM

- (1) (a) (v), (b) (i), (c) (ii), (d) (iv)
- (2) (a) (v), (b) (ii), (c) (iv), (d) (i)
- (3) (a) (iv), (b) (iii), (c) (i), (d) (ii)
- (4) (a) (v), (b) (iii), (c) (ii), (d) (i)

Official Ans. by NTA (4)

- Given below are two statement : one is labelled as Assertion A and the other is labelled as Reason R.
 Assertion A : SO₂(g) is adsorbed to a large extent than H₂(g) on activated charcoal.
 - **Reason** R : $SO_2(g)$ has a higher critical temperature than $H_2(g)$.

In the light of the above statements, choose the most appropriate answer from the options given below.

- (1)Both **A** and **R** are correct but **R** is not the correct explanation fo **A**
- (2) Both **A** and **R** are correct and **R** is the correct explanation of **A**.
- (3) A is not correct but **R** is correct.
- (4) **A** is correct but **R** is not correct.

Official Ans. by NTA (2)

- **6.** The **CORRECT** order of first ionisation enthalpy is:
 - (1) Mg < S < Al < P
- (2) Mg < Al < S < P
- (3) Al < Mg < S < P
- (4) Mg < Al < P < S

Official Ans. by NTA (3)

7. Given below are two statements:

Statement I : Hyperconjugation is a permanent effect.

Statement II : Hyperconjugation in ethyl cation $\left(CH_3 - \overset{+}{C}H_2\right)$ involves the overlapping of $C_{sp^2} - H_{ls}$ bond with empty 2p orbital of other carbon.

Choose the **correct** option:

- (1) Both **statement I** and **statement II** are false
- (2) **Statement I** is incorrect but **statement II** is true
- (3) Statement I is correct but statement II is false
- (4) Both **Statement I** and **statement II** are true.
- Official Ans. by NTA (3)

CHEMISTRY

SECTION-A

- 1. Which one of the following set of elements can be detected using sodium fusion extract?
 - (1) Sulfur, Nitrogen, Phosphorous, Halogens
 - (2) Phosphorous, Oxygen, Nitrogen, Halogens
 - (3) Nitrogen, Phosphorous, Carbon, Sulfur
 - (4) Halogens, Nitrogen, Oxygen, Sulfur

Official Ans. by NTA (1)

Consider the above reaction, the major product "P" formed is:-

(3)
$$CH_3$$
 $C-OCH_3$ CH_3 CH_3 $C-Br$

Official Ans. by NTA (2)

- 3. The number of neutrons and electrons, respectively, present in the radioactive isotope of hydrogen is:-
 - (1) 1 and 1
- (2) 3 and 1
- (3) 2 and 1
- (4) 2 and 2

Official Ans. by NTA (3)

4. Match List - I with List II:

List - I		List - II					
(a)	Li	(i)	photoelectric cell				
(b)	Na	(ii)	absorbent of CO ₂				
(c)	K	(iii)	coolant in fast breeder nuclear reactor				
(d)	Cs	(iv)	treatment of cancer				
		(v)	bearings for motor engines				

Choose the **correct** answer from the options given below:



8. Given below are two **statements**:

Statement I: $[Mn(CN)_6]^{3-}$, $[Fe(CN)_6]^{3-}$ and $[Co(C_2O_4)_3]^{3-}$ are d^2sp^3 hybridised.

Statement II: $[MnCl_6]^{3-}$ and $[FeF_6]^{3-}$ are paramagnetic and have 4 and 5 unpaired electrons, respectively.

In the light of the above statements, choose the **correct** answer from the options given below:

- (1) Statement I is correct but statement II is false
- (2) Both **statement I** and **statement II** are false
- (3) **Statement I** is incorrect but **statement II** is true
- (4) Both **statement I** and **statement II** are are true **Official Ans. by NTA (4)**
- 9. To an aqueous solution containing ions such as Al^{3+} , Zn^{2+} , Ca^{2+} , Fe^{3+} , Ni^{2+} , Ba^{2+} and Cu^{2+} was added conc. HCl, followed by H_2S .

The total number of cations precipitated during this reaction is/are:

(1) 1

(2) 3

(3)4

(4) 2

Official Ans. by NTA (1)

10. Given below are two **statements**:

Statement I : Penicillin is a bacteriostatic type antibiotic.

Statement II : The general structure of Penicillin is:

Choose the correct option:

- (1) Both **statement I** and **statement II** are false
- (2) Statement I is incorrect but statement II is true
- (3) Both **statement I** and **statement II** are true
- (4) Statement I is correct but statement II is false Official Ans. by NTA (2)
- **11.** Compound **A** gives D-Galactose and D-Glucose on hydrolysis. The compound **A** is :

(1) Amylose

(2) Sucrose

(3) Maltose

(4) Lactose

Official Ans. by NTA (4)

12.
$$R - CN \xrightarrow{(i) DIBAL-H \atop (ii) H_2O} R - Y$$

Consider the above reaction and identify "Y"

(1) – CH_2NH_2

(2) –CONH₂

(3) –CHO

(4) -COOH

Official Ans. by NTA (3)

OH Conc.
$$H_2SO_4$$
 A A B

consider the above reaction, and choose the correct statement:

- (1) The reaction is not possible in acidic medium
- (2) Both compounds **A** and **B** are formed equally
- (3) Compound A will be the major product
- (4) Compound **B** will be the major product

Official Ans. by NTA (3)

14. Match List - I with List - II:

	List - I	List - II							
	(compound)	(effect/affected species)							
(a)	Carbon monoxide	(i)	Carcinogenic						
(b)	Sulphur dioxide	(ii)	Metabolized	by					
			pyrus plants						
(c)	Polychlorinated	(iii)	Haemoglobin						
	biphenyls								
(d)	Oxides of Nitrogen	(iv)	Stiffness	of					
			flower buds						

Choose the **correct** answer from the options given below:

$$(1)$$
 (a) - (iii) , (b) - (iv) , (c) - (i) , (d) - (ii)

$$(3)$$
 (a) - (i), (b) - (ii), (c) - (iii), (d) - (iv)

Official Ans. by NTA (1)

- 15. If the Thompson model of the atom was correct, then the result of Rutherford's gold foil experiment would have been:
 - (1) All of the α -particles pass through the gold foil without decrease in speed.
 - (2) α -Particles are deflected over a wide range of angles.
 - (3) All α -particles get bounced back by 180°
 - (4) α -Particles pass through the gold foil deflected by small angles and with reduced speed.

Official Ans. by NTA (4)

Final JEE-Main Exam July, 2021/27-07-2021/ Evening Session

- Number of Cl = O bonds in chlorous acid, chloric 16. acid and perchloric acid respectively are:
 - (1) 3, 1 and 1
- (2) 4, 1 and 0
- (3) 1, 1 and 3
- (4) 1, 2 and 3

Official Ans. by NTA (3)

- Select the correct statements. 17.
 - (A) Crystalline solids have long range order.
 - (B) Crystalline solids are isotropic.
 - (C) Amorphous solid are sometimes called pseudo solids.
 - (D) Amorphous solids soften over a range of temperatures.
 - (E) Amorphous solids have a definite heat of fusion. Choose the most appropriate answer from the options given below.
 - (1)(A),(B),(E) only
 - (2) (B), (D) only
 - (3) (C), (D) only
 - (4) (A), (C), (D) only

Official Ans. by NTA (4)

What is A in the following reaction? 18.

$$CH_2Br \xrightarrow{(i) \qquad \qquad N^{\odot}K^{\oplus}} A$$

$$(ii) \xrightarrow{O} A$$

$$(iii) \xrightarrow{O}OH/H_2O \qquad (Major Product)$$

$$(1) \bigcirc NH-CH_2 - \bigcirc$$

Official Ans. by NTA (4)

19. The correct sequence of correct reagents for the following transformation is:-

$$\stackrel{\text{NO}_2}{\bigodot} \longrightarrow \stackrel{\text{OH}}{\bigodot}_{\text{Cl}}$$

- (1) (i) Fe, HCl
- (ii) Cl₂, HCl,
- (iii) NaNO₂, HCl, 0°C (iv) H₂O/H⁺ (2) (i) Fe, HCl
 - (ii) NaNO2, HCl, 0°C
 - (iii) H₂O/H⁺
- (iv) Cl₂, FeCl₃
- (3) (i) Cl₂, FeCl₃
- (ii) Fe, HCl
- (iii) NaNO₂, HCl, 0°C (iv) H₂O/H⁺
- (4) (i) Cl₂, FeCl₃
- (ii) NaNO2, HCl, 0°C
- (iii) Fe, HCl
- (iv) H_2O/H^+

Official Ans. by NTA (3)

- The addition of silica during the extraction of 20. copper from its sulphide ore :-
 - (1) converts copper sulphide into copper silicate
 - (2) converts iron oxide into iron silicate
 - (3) reduces copper sulphide into metallic copper
 - (4) reduces the melting point of the reaction mixture

Official Ans. by NTA (2)

SECTION-B

1. The equilibrium constant for the reaction

$$A(s) \longrightarrow M(s) + \frac{1}{2}O_2(g)$$

is $K_p = 4$. At equilibrium, the partial pressure of O_2 is atm. (Round off to the nearest integer)

Official Ans. by NTA (16)

When 400 mL of 0.2M H₂SO₄ solution is mixed with 600 mL of 0.1 M NaOH solution, the increase in temperature of the final solution is \times 10⁻² K. (Round off to the nearest integer).

[Use :
$$H^+$$
 (aq) + OH^- (aq) \rightarrow H_2O :

$$\Delta_{v}H = -57.1 \text{ kJ mol}^{-1}$$

Specific heat of $H_2O = 4.18 \text{ J K}^{-1} \text{ g}^{-1}$

density of $H_2O = 1.0 \text{ g cm}^{-3}$

Assume no change in volume of solution on mixing.

Official Ans. by NTA (2)

ALLEN Ans. (82)

3. $2SO_2(g) + O_2(g) \rightarrow 2SO_3(g)$

The above reaction is carried out in a vessel starting with partial pressure $P_{SO_2}=250\,\text{m}\,\text{bar}$, $P_{O_2}=750\,\text{m}$ bar and $P_{SO_3}=0$ bar. When the reaction is complete, the total pressure in the reaction vessel is _____ m bar. (Round off of the nearest integer).

Official Ans. by NTA (875)

4. 10.0 mL of 0.05 M KMnO_4 solution was consumed in a titration with 10.0 mL of given oxalic acid dihydrate solution. The strength of given oxalic acid solution is $\times 10^{-2} \text{ g/L}$.

(Round off to the nearest integer)

Official Ans. by NTA (1575)

5. The total number of electrons in all bonding molecular orbitals of O_2^{2-} is

(Round off to the nearest integer)

Official Ans. by NTA (10)

6. 3 moles of metal complex with formula Co(en)₂Cl₃ gives 3 moles of silver chloride on treatment with excess of silver nitrate. The secondary valency of Co in the complex is _____.

(Round off to the nearest integer)

Official Ans. by NTA (6)

7. In a solvent 50% of an acid HA dimerizes and the rest dissociates. The van't Hoff factor of the acid is $\times 10^{-2}$.

(Round off to the nearest integer)

Official Ans. by NTA (125)

8. The dihedral angle in staggered form of Newman projection of 1, 1, 1-Trichloro ethane is degree. (Round off to the nearest integer)

(Round off to the nearest integer)

Official Ans. by NTA (60)

9. For the first order reaction A → 2B, 1 mole of reactant A gives 0.2 moles of B after 100 minutes. The half life of the reaction is min. (Round off to the nearest integer).

[Use: $\ln 2 = 0.69$, $\ln 10 = 2.3$]

Properties of logarithms : $\ln x^y = y \ln x$;

$$\ln\left(\frac{x}{y}\right) = \ln x - \ln y$$

(Round off to the nearest integer)

Official Ans. by NTA (300) ALLEN Ans. (600 to 700)

10. For the cell

$$Cu(s) \mid Cu^{2^{+}}(aq) \ (0.1M) \parallel Ag^{+} \ (aq) \ (0.01M) \mid Ag(s)$$

the cell potential $E_1 = 0.3095 \text{ V}$

For the cell

$$Cu(s) | Cu^{2+}(aq) (0.01 \text{ M}) || Ag^{+}(aq) (0.001 \text{ M}) | Ag(s)$$

the cell potential = $___ \times 10^{-2}$ V. (Round off the Nearest Integer).

[Use :
$$\frac{2.303 \text{ RT}}{\text{F}} = 0.059 \text{]}$$

Official Ans. by NTA (28)