

परिक्षार्थी प्रवेश-पत्र के अनुसार भरें

पत्र Subject: **MATHEMATICS STANDARD**

पत्र कोड Subject Code: **041**

परीक्षा का दिन एवं तिथि
Day & Date of the Examination: **THURSDAY 12.3.2020**

उत्तर देने का माध्यम
Medium of answering the paper: **ENGLISH**

प्रश्न पत्र के ऊपर लिखें
कोड को दर्शाएँ
Write code No. as written on
the top of the question paper.

Code Number
30/5/2

Set Number
① ● ③ ④

अतिरिक्त उत्तर-पुस्तिका (ओं) की संख्या
No. of supplementary answer-book(s) used

—

बेधमाक विकलांग व्यक्ति
Person with Benchmark Disabilities

हाँ / नहीं
Yes / No

NO

विकलांगता का कोड
(प्रवेश पत्र के अनुसार)
Code of Disabilities
(as given on Admit Card)

—

क्या लेखन - लिपिक उपलब्ध कराया गया :
Whether writer provided :

हाँ / नहीं
Yes / No

NO

यदि दृष्टिहीन हैं तो उपयोग में लाए गए
सॉफ्टवेयर का नाम :
If Visually challenged, name of software used :

—

* एक खाने में एक अक्षर लिखें। नाम के प्रत्येक भाग के बीच एक खाना रिक्त छोड़ दें। यदि परीक्षार्थी का नाम 24 अक्षरों से अधिक है, तो केवल नाम के प्रथम 24 अक्षर ही लिखें।
Each letter be written in one box and one box be left blank between each part of the name. In case Candidate's Name exceeds 24 letters, write first 24 letters.

कार्यालय उपयोग के लिए
Space for office use

SECTION - D

choice - ①

$$P(x) = 2x^4 - x^3 - 11x^2 + 5x + 5$$

Given zeroes: $\sqrt{5}, -\sqrt{5}$

sum of zeroes = $\sqrt{5} - \sqrt{5} = 0$

Product of zeroes = $\sqrt{5} \times -\sqrt{5} = -5$

$$f(x) = x^2 + 0x - 5$$

$$\begin{array}{r} 2x^2 - 12x - 1 \\ \hline x^2 + 0x - 5 \end{array} \quad \begin{array}{r} 2x^4 - x^3 - 11x^2 + 5x + 5 \\ -2x^4 + 0x^3 + 10x^2 \\ \hline 0x^3 - x^3 - 11x^2 + 5x + 5 \\ -x^3 + 0x^2 + 5x + 5 \\ \hline 0x^3 - 1x^2 + 5x + 5 \\ +x^2 + 0x + 5 \\ \hline 0x^3 - 1x^2 + 5x + 5 \\ +1x^2 + 0x + 5 \\ \hline 0x^3 + 0x^2 + 5x + 10 \\ 0x^3 + 0x^2 + 5x + 10 \end{array}$$

$$\begin{array}{r} 0x^3 - 1x^2 + 5x + 5 \\ +x^2 + 0x + 5 \\ \hline 0x^3 - 1x^2 + 5x + 5 \\ +1x^2 + 0x + 5 \\ \hline 0x^3 + 0x^2 + 5x + 10 \\ 0x^3 + 0x^2 + 5x + 10 \end{array}$$

[Faint handwritten notes and calculations, including a table with columns for powers of x and coefficients.]

$$g(x) = 2x^2 - 1x - 1$$

$$g(x) = 0$$

$$2x^2 - 1x - 1 = 0$$

$$2x^2 - 2x + 1x - 1 = 0$$

$$2x(x-1) + 1(x-1) = 0$$

$$x = 1$$

$$x = -\frac{1}{2}$$

$$\frac{1}{-8} + \frac{1}{8} = \frac{-11}{4}$$

$$\frac{-11}{4} \times \frac{2}{2} = \frac{-11}{2}$$

$$2 - 1 - 11 + 5 + 5 = -2x + 1$$

$$\frac{1 + 1 - 22 - 20 + 40}{8} = \frac{-11}{4}$$

$$\frac{-11}{4} \times \frac{2}{2} = \frac{-11}{2}$$

$$\frac{4 - 1 - 22}{8} = \frac{-19}{8}$$

Ans: The other zeroes of $P(x)$ are: $1, -\frac{1}{2}$

Zeroes of $P(x)$: $\sqrt{5}, -\sqrt{5}, 1, -\frac{1}{2}$

(39)




Given: Right $\triangle ABC$, $\angle B = 90^\circ$

To prove: $AC^2 = AB^2 + BC^2$

Construction:

Draw $BD \perp AC$.



Proof:

In $\triangle ABC$ and $\triangle ADB$,

$$\angle ABC = \angle ADB = 90^\circ$$

$\angle 1 = \angle 1$ (common)

$\therefore \triangle ABC \sim \triangle ADB$ (AA)

by CPST, $\frac{AB}{AD} = \frac{BC}{DB} = \frac{AC}{AB}$

$$\Rightarrow \frac{AB}{AD} = \frac{AC}{AB}$$

$$AB^2 = AC \cdot AD \rightarrow ①$$

In $\triangle ABC$ and $\triangle BDC$

$$\angle ABC = \angle BDC = 90^\circ$$

$\angle 2 = \angle 2$ (common)

$\therefore \triangle ABC \sim \triangle BDC$ (AA)

by CPST, $\frac{AB}{BD} = \frac{BC}{DC} = \frac{AC}{BC}$

$$\frac{BC}{DC} = \frac{AC}{BC} \Rightarrow BC^2 = AC \cdot DC \rightarrow ②$$


Adding ① and ②,

$$AB^2 + BC^2 = AC \cdot AD + AC \cdot DC$$

$$AB^2 + BC^2 = AC(AD + DC)$$

$$AB^2 + BC^2 = AC^2 //$$

Hence proved.



Let AB \rightarrow Transmission tower

BC \rightarrow Building - 20m

P \rightarrow Point on the ground.

In ΔPBC , $\angle C = 90^\circ$


$$\tan 45^\circ = \frac{20}{x} \Rightarrow x = \frac{20}{1} = 20$$

In ΔPAC , $\angle C = 90^\circ$

$$\tan 60^\circ = \frac{h+20}{x}$$

$$\frac{20+h}{20} = \sqrt{3}$$

$$h = 20\sqrt{3} - 20$$



$$\frac{h+20}{20} = \sqrt{3}$$

$$h+20 = 20\sqrt{3}$$

$$h = 20(\sqrt{3} - 1)$$

$$h = 20 \times 1.73 - 20$$

$$h = 34.6 - 20$$

$$h = 14.6$$

$h = 20(\sqrt{3}-1)$
 $h = 14.64m$

Ans:- Height of transmission tower = 14.64m

choice-1

(37)

Avg (in yrs) No. of persons less than c.f


(C.o.I)	(f)	(cf)
D-10	5	5
10-20	15	20
20-30	20	40
30-40	25	65
40-50	15	80
50-60	11	91
60-70	9	100

Median Age:

$L + \left(\frac{\frac{n}{2} - cf}{f} \right) \times h \Rightarrow 30 + \frac{50-40}{25} \times 10$
 $\Rightarrow 30 + \frac{100}{25} \Rightarrow 34 \text{ years}$

Median of the distribution: 34 years (by graph and calculation)

(36)



For Hemisphere:

$$r = 7 \text{ cm}$$

For cone:

$$h = 10 \text{ cm}$$

$$r = 7 \text{ cm}$$

$$d = \sqrt{100 + 49} = 12.2 \text{ cm}$$

Volume of the toy: Volume of cone + Volume of hemisphere.

$$\frac{1}{3} \pi r^2 h + \frac{2}{3} \pi r^3$$

$$\frac{1}{3} \pi r^2 (h + 2r)$$

$$\frac{1}{3} \times \frac{22}{7} \times 7 \times 7 (10 + 14)$$

$$\frac{1}{3} \times \frac{22}{7} \times 7 \times 7 \times 24$$

$$\Rightarrow 1232 \text{ cm}^3$$

Area of coloured sheet required : CSA of cone + CSA of hemisphere

$$\Rightarrow \pi r l + 2\pi r^2$$

$$\Rightarrow \pi r (l + 2r)$$

$$\Rightarrow \frac{22}{7} \times 7 \times (12 + 2 \times 7)$$

$$\Rightarrow \frac{22}{7} \times 7 \times 26 = 26 \times 22$$

$$\Rightarrow 576.4 \text{ cm}^2$$

Ans: Volume of the toy = 1232 cm^3

Area of coloured sheet required = 576.4 cm^2 .

For motorboat:

Speed in still water = 18 km/hr

Let speed of stream = $x \text{ km/hr}$

upstream speed = $18 - x \text{ km/hr}$

downstream speed = $18 + x \text{ km/hr}$

35

$$t = \frac{d}{s}$$

$$\frac{24}{18-x} - \frac{24}{18+x} = 1$$

$$24 \left(\frac{1}{18-x} - \frac{1}{18+x} \right) = 1$$

$$18+x > 18+x$$

$$(18-x)(18+x) = 24$$

$$\frac{2x}{324-x^2} = \frac{1}{24}$$

$$48x = 324 - x^2$$

$$x^2 + 48x - 324 = 0$$

$$(x+54)(x-6) = 0$$

$$x = -54 \text{ (invalid - speed cannot be negative)}$$

$$x = 6$$

Ans:

Speed of the stream = 6 km/hr.

$$\begin{array}{r} 6 \\ 18 \overline{) 108} \\ \underline{18} \\ 144 \\ \underline{144} \\ 0 \end{array}$$

$$24$$

$$-12$$

$$211$$

$$414$$

$$-207$$

$$6$$

$$118$$

$$1$$

$$24$$

$$48x = 324 - x^2$$

$$x^2 + 48x - 324 = 0$$

$$(x+54)(x-6) = 0$$

$$x = -54 \text{ (invalid - speed cannot be negative)}$$

$$x = 6$$

Ans:

Speed of the stream = 6 km/hr.

$$\frac{0}{65} \frac{0}{50}$$

SECTION - C

(34) (i) Total number of 'Numbers' on spinner = 6
 'Even numbers' = 3

$P(\text{Shweta being allowed to pick a marble}) = \frac{\text{No. of favourable outcomes}}{\text{Total number of outcomes}}$

Ans: Probability of being allowed to pick a marble $\Rightarrow \frac{3}{6}$

(ii) Total number of marbles = 20
 black marbles = 6.

$$P(\text{Shweta winning a prize}) = \frac{6}{20} \Rightarrow \frac{3}{10} \Rightarrow 0.3$$

$\left. \begin{array}{l} \text{Number of favourable} \\ \text{outcomes} \end{array} \right\}$
 $\left. \begin{array}{l} \text{Total number of} \\ \text{outcomes} \end{array} \right\}$

Ans: Probability of getting a prize = $\frac{3}{10}$

(33)

$$a = 54$$

$$d = -3$$

$$a_n = 0$$

$$a_n = 0$$

$$a + (n-1)d = 0$$

$$54 - 3(n-1) = 0$$

$$78(n-1) = 18$$

$$n = 19$$

Ans: $n = 19$

$$a_{19} = 0$$

$$S_n = \frac{n}{2} (2a + (n-1)d)$$

$$S_n = \frac{n}{2} [108 - 3(n-1)]$$

$$S_n = \frac{n}{2} [108 - 3n + 3] \Rightarrow \frac{n}{2} (111 - 3n)$$

$$\frac{54 - 3(18)}{54 - 54} = \frac{0}{0} = \frac{19}{3}$$

$$\frac{54}{108} = \frac{1}{2}$$

$$\frac{54}{108} = \frac{1}{2}$$

$$a = 54$$

$$d = -3$$

$$54 - 3(n-1) = 0$$

$$78(n-1) = 18$$

$$n = 19$$

$$a_{19} = 0$$

$$S_n = \frac{n}{2} (2a + (n-1)d)$$

$$S_n = \frac{n}{2} [108 - 3(n-1)]$$

$$S_n = \frac{n}{2} [108 - 3n + 3] \Rightarrow \frac{n}{2} (111 - 3n)$$

Here, $n = 19$

$$S_{19} = \frac{19}{2} [111 - 17]$$

$$S_{19} = \frac{19 \times 54}{2}$$

Ans: $S_{19} = 513$

Ans: $n = 19$

$$S_n = S_{19} = 513$$

choice - ①


R: 1 B(-2, -7)



Let the y axis meet the line segment joining points A(6, -4) and B(-2, -7)

Let P divide AB in the ratio R:1

Let P be (0, y)



$$\frac{y - (-4)}{-4 - (-7)} = \frac{0 - 6}{-7 - 6}$$

$$y + 4 = \frac{6}{13} \times 13$$

$$y + 4 = 6$$

$$y = 6 - 4$$

$$y = 2$$

$$P = (0, 2)$$

14 A P(0,y)

coordinates of P: $P\left(\frac{-2R+6}{R+1}, \frac{-7R-4}{R+1}\right)$

$$\frac{-2R+6}{R+1} = 0$$

$$-2R+6=0$$

$$2R=6$$

$$R=3$$

Ans: Ratio in which y axis divides AB = 3:1

$$y = \frac{-7(3)-4}{(3)+1}$$

$$y = \frac{-21-4}{4}$$

$$y = \frac{-25}{4}$$

Ans: Point of intersection = $P\left(0, \frac{-25}{4}\right)$ of y axis and line segment.

(31)

$$870 - 3 = 867$$

$$258 - 3 = 255$$

HCF(867, 255) by Euclid's Division Algorithm:

$$867 = 255 \times 3 + 102$$

$$255 = 102 \times 2 + 51$$

$$102 = 51 \times 2 + 0$$

$$\text{HCF}(867, 255) = 51$$

Ans: The largest number which divides 870 and 258 leaving remainder 3 in each case is 51.

$$\begin{array}{r} 255 \\ 3 \overline{) 867} \\ \underline{765} \\ 1020 \\ \underline{765} \\ 255 \end{array}$$

$$\begin{array}{r} 255 \\ 2 \overline{) 510} \\ \underline{510} \\ 0 \end{array}$$

$$\begin{array}{r} 255 \\ 3 \overline{) 765} \\ \underline{765} \\ 0 \end{array}$$

$$\begin{array}{r} 255 \\ 3 \overline{) 765} \\ \underline{765} \\ 0 \end{array}$$

$$\begin{array}{r} 255 \\ 3 \overline{) 765} \\ \underline{765} \\ 0 \end{array}$$

$$\begin{array}{r} 17 \\ 51 \overline{) 867} \\ \underline{867} \\ 0 \end{array}$$

$$\begin{array}{r} 51 \\ 2 \overline{) 102} \\ \underline{102} \\ 0 \end{array}$$

choice - ②

Let the present age of

Father = x yearsson = y years

$$x = 3y + 3$$

$$x - 3y - 3 = 0 \rightarrow \textcircled{1}$$

$$x + 3 = 2y + 10$$

$$x + 3 = 2(y + 3) + 10$$

$$x + 3 = 2y + 6 + 10$$

$$x - 2y = 16 + 3 - 3$$

$$x - 2y - 13 = 0 \rightarrow \textcircled{2}$$

Solving ① and ②,

$$x - 2y - 3 = 0$$

$$-x + 2y + 13 = 0$$

$$\rightarrow -y + 10 = 0$$

$$y = 10$$

$$x = 33$$

Ans: Present age of father = 33 years
 son = 10 years

$$\frac{2 \cos^3 \theta - \cos \theta}{\sin \theta - 2 \sin^3 \theta} = \cot \theta$$

LHS:

$$\frac{\cos \theta (2 \cos^2 \theta - 1)}{\sin \theta (1 - 2 \sin^2 \theta)}$$

$$\frac{\cos \theta [2(1 - \sin^2 \theta) - 1]}{\sin \theta (1 - 2 \sin^2 \theta)}$$


$$\frac{\cos \theta [2 - 2 \sin^2 \theta - 1]}{\sin \theta (1 - 2 \sin^2 \theta)} \Rightarrow \frac{\cos \theta}{\sin \theta} \times \frac{(1 - 2 \sin^2 \theta)}{(1 - 2 \sin^2 \theta)} \Rightarrow \cot \theta$$

LHS = RHS = $\cot \theta$ //

Q2) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (2) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (3) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (4) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (5) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (6) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (7) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (8) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (9) With M as centre circle is drawn. B and P to meet the circumference of A and B.
 (10) With M as centre circle is drawn. B and P to meet the circumference of A and B.

Choice - (2)

(28)



STEPS OF CONSTRUCTION:

- ① Draw a circle with centre O and radius 3.5 cm
- ② Take a point P outside the circle so that $OP = 7\text{ cm}$.
- ③ Join OP .
- ④ Construct perpendicular bisector for OP and let it meet OP at M .
- ⑤ With M as centre and OM as radius draw a circle, passing through O and P to meet the previous circle at A and B .
- ⑥ Join AP, BP . AP and BP are the required tangents.

SECTION - B

26 Since A, B, C are interior angles of $\triangle ABC$,

$$\angle A + \angle B + \angle C = 180^\circ \text{ (ASP)}$$

$$\angle A + \angle B + \angle C = 90^\circ$$

$$\frac{\angle B + \angle C}{2} = \frac{90^\circ - \angle A}{2} \rightarrow \textcircled{1}$$

$$\text{LHS: } \cot\left(\frac{B+C}{2}\right) = \tan\left(\frac{A}{2}\right) \rightarrow \text{To Prove:}$$

$$\cot\left(\frac{B+C}{2}\right)$$

sub(1),

$$\cot\left(90^\circ - \frac{\angle A}{2}\right)$$

$$\left. \begin{aligned} \cot\left(90^\circ - \theta\right) &= \tan \theta \end{aligned} \right\}$$

$$\cot\left(\frac{\angle A}{2}\right) = \text{RHS} //$$

Proved.

(25)

Let us assume to the contrary that $\sqrt{5} + 2\sqrt{7}$ is rational.

Then $\sqrt{5} + 2\sqrt{7}$ is of the form $\frac{p}{q}$ where p and q are co-primes and $q \neq 0$.

$$\frac{p}{q} = \sqrt{5} + 2\sqrt{7}$$

$$\frac{p - \sqrt{5}q}{q} = 2\sqrt{7}$$

$$\frac{p - \sqrt{5}q}{2q} = \sqrt{7}$$


$\frac{p - \sqrt{5}q}{2q}$ is rational as p and q are integers

This contradicts the given fact that $\sqrt{7}$ is irrational.
 \therefore Our assumption is wrong.

$\sqrt{5} + 2\sqrt{7}$ is irrational //

Proved.

Given: $\triangle ABC$



$DE \parallel BC$
 $DF \parallel AE$

To Prove: $\frac{BF}{FE} = \frac{BE}{EC}$

Proof: In $\triangle ABE$, $DF \parallel AE$

by BPT, $\frac{BF}{FE} = \frac{BD}{DA}$ \rightarrow (1)

In $\triangle ABC$, $DE \parallel AC$

by BPT, $\frac{BE}{EC} = \frac{BD}{DA}$ \rightarrow (2)


From (1) and (2),

$\frac{BF}{FE} = \frac{BE}{EC}$ ✓

Proved

37

Age in years



Scale:
 x axis: 1cm = 5 years
 y axis: 1cm = 5 units

23

For Small cube: $a = 2\text{cm}$

Large cube: $A = 10\text{cm}$

Let number of cubes = n

$$A^3 = n \times a^3$$

$$n = \frac{A^3}{a^3}$$


$$n = \frac{10^3}{2^3} = \frac{1000}{8} = 125$$

Ans:

Number of cubes that can be made = 125 cubes

24

choice - 1



Lengths of

- ① $AP = AR$
- ② $DP = DS$
- ③ $BR = BQ$
- ④ $CR = CQ$

Tangents from external points A, B, C, D to the circle are equal

Let the circle and square meet at P, Q, R, S.

Adding ①, ②, ③, ④,

$$AP + DP + BR + CR = AS + BS + CS + DS + CS$$

$$AD + BC = AB + DC \quad // \quad \text{Proved.}$$

Marks No. of students (f)

0-10 4

10-20 6

20-30 7

30-40 12 modal class

40-50 5

50-60 6

Mode: $L + \frac{f_1 - f_0}{f_1 - f_0 + f_1 - f_2} \times h$

$$= 30 + \frac{12-7}{12-7+12-5} \times 10$$

$$= 30 + \frac{5}{10} \times 10 = 35$$

~~35 marks~~

$$30 + \frac{12-7}{24-7-10} \times 10$$

$$30 + \frac{12-7}{24-12} \times 10$$

$$30 + \frac{12-7}{12-6} \times 10$$

$$30 + \frac{5}{12-6} \times 10 = 30 + \frac{50}{6}$$


$h=10$

$$\text{Mode} = 30 + \frac{12-7}{24-7-5} \times 10$$

$$30 + \frac{5}{17} \times 10 = 34.17 \text{ marks (approx)}$$

Ans: Medal marks = 34.17 marks (approx)

SECTION - A



Let hgt of tower = AB = h m

In right ΔABC ,

$$\tan 30^\circ = \frac{h}{30}$$

$$\frac{1}{\sqrt{3}} = \frac{h}{30} \Rightarrow h = \frac{30\sqrt{3}}{3}$$

Ans: Height of tower = 17.32 m (approx)

4	16
6	25
	24
10	
6	
40	
36	
40	

4.17

(20)

(19) $2 \sec 30^\circ \times \tan 60^\circ$

$2 \times \frac{2}{\sqrt{3}} \times \sqrt{3} = 4$

(18) $n = 100$

Sum of first 100 natural numbers = $\frac{n(n+1)}{2}$

Ans: Sum of 1st 100 natural nos. $\Rightarrow \frac{100 \times 101}{2} = 5050$

(17) Sum of zeroes = -3

Product of zeroes = 2.

Ans: The required Polynomial:

$P(x) = k(x^2 + 3x + 2)$

Product of 2 nos. = $L.C.M \times H.C.F$

Let other number be x

$x \times 26 = 182 \Rightarrow x = 7$

$x = 7$

Ans: other number is 7

✓ $(1 - \frac{1}{2})^2$ (2)

(15) $(1 - \cos^2 A) (1 + \cot^2 A)$ $(1 + \frac{\cos^2 A}{\sin^2 A})$

$(1 - \cos^2 A) (1 + \frac{\cos^2 A}{\sin^2 A})$

$(1 - \cos^2 A) (\frac{\sin^2 A + \cos^2 A}{\sin^2 A})$


$(1 - \cos^2 A) \Rightarrow \frac{\sin^2 A}{\sin^2 A} = 1$

ans 21

(14) $P(\text{sure event}) = 1$

(13) Similar

(12) $u_i = \frac{x_i - a}{h}$

(11)  length of diagonal AB = $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
 $= \sqrt{(0 - 4)^2 + (-3 - 0)^2} = \sqrt{16 + 9} = 5$ units

Ans: length of diagonal of rectangle ABC is 5 units.

$$\text{Volume} = 12\pi$$

$$\frac{4}{3}\pi r^3 = 12\pi$$

$$\frac{4}{3}\pi r^3 = 12\pi \quad (\sqrt[3]{12\pi}) \quad (\sqrt[3]{12\pi})$$

$$r^3 = 9$$

$$r^3 = 9$$

$$r = \sqrt[3]{9} \text{ cm.}$$

Ans: (c) $\sqrt[3]{9}$ cm ✓

(A) 50° ✓

$$\frac{3}{2}x + \frac{5}{3}y = 7$$

$$9x + 10y = 14$$

$$\frac{a_1}{a_2} = \frac{3}{2} \times \frac{1}{9} = \frac{1}{6}$$

$$\frac{b_1}{b_2} = \frac{5}{3} \times \frac{1}{10} = \frac{1}{6}$$

$$\frac{c_1}{c_2} = \frac{+7}{+14} = \frac{1}{2}$$

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$$

Ans (B) Inconsistent

7 $A(-6, 3)$

$B(6, 4)$

$$O = \left(\frac{-6+6}{2}, \frac{3+4}{2} \right)$$

$$O = \left(0, \frac{7}{2} \right)$$

Ans - (c) $\left[0, \frac{7}{2} \right]$

8 $AB^2 = AC^2 + BC^2$

$$AB^2 = 2AC^2$$

Ans - (A) $AB^2 = 2AC^2$

(5) (A) 2 ✓

(4) A (m, -n)
B (-m, n)

$$AB = \sqrt{(m+m)^2 + (-n-n)^2}$$
$$AB = \sqrt{4m^2 + 4n^2}$$

$$AB = 2\sqrt{m^2 + n^2}$$

ans = (c) $2\sqrt{m^2 + n^2}$

(3) (B) 4cm ✓

(2) (c) $\frac{4}{3}, \frac{7}{3}, \frac{9}{3}, \frac{12}{3}$

$\frac{1}{2} + \frac{1}{2} = 1$
 $\frac{1}{4} + \frac{1}{4} = \frac{1}{2}$
 $\frac{1}{8} + \frac{1}{8} = \frac{1}{4}$
 $\frac{1}{16} + \frac{1}{16} = \frac{1}{8}$

ans (B) 10 ✓

(F) $\sqrt{(-1)^2}$

(B) $\sqrt{(-1)^2}$

$$\frac{1+3}{2} = 2$$

$$\frac{1}{2} + \frac{1}{2} = 1$$

$$\frac{0.8}{1.2} = \frac{2.8}{3.6}$$
$$\frac{0.8}{2.0} = \frac{0.8}{2.0}$$

$$3 + \sqrt{2} - 3 = \sqrt{2}$$

$$\frac{3}{3} + \frac{2}{3} = \frac{5}{3}$$
$$\frac{1}{3} - \frac{4}{3} = -\frac{3}{3} = -1$$

$$-\frac{2}{5} + 1 = \frac{3}{5}$$
$$-\frac{3}{5} + \frac{12}{5} = \frac{9}{5}$$

$$2x^2 + kx + 2 = 0$$

For equal roots;

$$b^2 - 4ac = 0$$

$$k^2 - 16 = 0$$

$$k^2 = 16$$

$$k = \pm 4$$

Ans (B) ± 4