

**FINAL JEE–MAIN EXAMINATION – JUNE, 2022**

 (Held On Saturday 25<sup>th</sup> June, 2022)

TIME : 9 : 00 AM to 12 : 00 PM

**CHEMISTRY**
**SECTION-A**

1. Bonding in which of the following diatomic molecule(s) become(s) stronger, on the basis of MO Theory, by removal of an electron ?

(A) NO (B) N<sub>2</sub>  
 (C) O<sub>2</sub> (D) C<sub>2</sub>  
 (E) B<sub>2</sub>

Choose the most appropriate answer from the options given below :-

(A) (A), (B), (C) only (B) (B), (C), (E) only  
 (C) (A), (C) only (D) (D) only

**Official Ans. by NTA (C)**

**Allen Ans. (C)**

- Sol.** Bond strength  $\propto$  Bond order  
 removal of electron from antibonding MO increases B.O.

NO & O<sub>2</sub> has valence e<sup>-</sup> in  $\pi^*$ orbital.

2. Incorrect statement for Tyndall effect is :-  
 (A) The refractive indices of the dispersed phase and the dispersion medium differ greatly in magnitude.  
 (B) The diameter of the dispersed particles is much smaller than the wavelength of the light used.  
 (C) During projection of movies in the cinemas hall, Tyndall effect is noticed.  
 (D) It is used to distinguish a true solution from a colloidal solution.

**Official Ans. by NTA (B)**

**Allen Ans. (B)**

- Sol.** The diameter of dispersed particle should be somewhat below or near the wavelength of light.

3. The pair, in which ions are isoelectronic with Al<sup>3+</sup> is :-

(A) Br<sup>-</sup> and Be<sup>2+</sup> (B) Cl<sup>-</sup> and Li<sup>+</sup>  
 (C) S<sup>2-</sup> and K<sup>+</sup> (D) O<sup>2-</sup> and Mg<sup>2+</sup>

**Official Ans. by NTA (D)**

**Allen Ans. (D)**

- Sol.** Isoelectronic species have same no. of electrons  
 Al<sup>3+</sup>, O<sup>2-</sup>, Mg<sup>2+</sup> all have 10 electrons.

**TEST PAPER WITH SOLUTION**

4. Leaching of gold with dilute aqueous solution of NaCN in presence of oxygen gives complex [A], which on reaction with zinc forms the elemental gold and another complex [B]. [A] and [B], respectively are :-

(A) [Au(CN)<sub>4</sub>]<sup>-</sup> and [Zn(CN)<sub>2</sub>(OH)<sub>2</sub>]<sup>2-</sup>  
 (B) [Au(CN)<sub>2</sub>]<sup>-</sup> and [Zn(OH)<sub>4</sub>]<sup>2-</sup>  
 (C) [Au(CN)<sub>2</sub>]<sup>-</sup> and [Zn(CN)<sub>4</sub>]<sup>2-</sup>  
 (D) [Au(CN)<sub>4</sub>]<sup>2-</sup> and [Zn(CN)<sub>6</sub>]<sup>4-</sup>

**Official Ans. by NTA (C)**

**Allen Ans. (C)**

- Sol.** Au + NaCN  $\rightarrow$  Na[Au(CN)<sub>2</sub>]  
 Zn + Na[Au(CN)<sub>2</sub>]  $\rightarrow$  Na<sub>2</sub>[Zn(CN)<sub>4</sub>] + Au

5. Number of electron deficient molecules among the following

PH<sub>3</sub>, B<sub>2</sub>H<sub>6</sub>, CCl<sub>4</sub>, NH<sub>3</sub>, LiH and BCl<sub>3</sub> is

(A) 0 (B) 1  
 (C) 2 (D) 3

**Official Ans. by NTA (C)**

**Allen Ans. (C)**

- Sol.** Electron deficient species have less than 8 electrons (or two electrons for H) in their valence (incomplete octet)

B<sub>2</sub>H<sub>6</sub>, BCl<sub>3</sub> have incomplete octet.

6. Which one of the following alkaline earth metal ions has the highest ionic mobility in its aqueous solution?

(A) Be<sup>2+</sup> (B) Mg<sup>2+</sup>  
 (C) Ca<sup>2+</sup> (D) Sr<sup>2+</sup>

**Official Ans. by NTA (D)**

**Allen Ans. (D)**

- Sol.** Highest ionic mobility corresponds to lowest extent of hydration and highest size of gaseous ion.

Hence Sr<sup>2+</sup> has the highest ionic mobility in its aqueous solution

7. White precipitate of AgCl dissolves in aqueous ammonia solution due to formation of :

- (A)  $[\text{Ag}(\text{NH}_3)_4]\text{Cl}_2$       (B)  $[\text{Ag}(\text{Cl})_2(\text{NH}_3)_2]$   
(C)  $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$       (D)  $[\text{Ag}(\text{NH}_3)\text{Cl}]\text{Cl}$

**Official Ans. by NTA (C)**

**Allen Ans. (C)**

**Sol.**  $\text{AgCl} + 2\text{NH}_3 \rightarrow [\text{Ag}(\text{NH}_3)_2]^+\text{Cl}^-$   
soluble


8. Cerium (IV) has a noble gas configuration. Which of the following is correct statement about it?

- (A) It will not prefer to undergo redox reactions.  
(B) It will prefer to gain electron and act as an oxidizing agent  
(C) It will prefer to give away an electron and behave as reducing agent  
(D) It acts as both, oxidizing and reducing agent.

**Official Ans. by NTA (B)**

**Allen Ans. (B)**

**Sol.** Cerium exists in two different oxidation state +3, +4



It shows  $\text{Ce}^{+4}$  acts as a strong oxidising agent & accepts electron.


9. Among the following, which is the strongest oxidizing agent ?

- (A)  $\text{Mn}^{3+}$       (B)  $\text{Fe}^{3+}$   
(C)  $\text{Ti}^{3+}$       (D)  $\text{Cr}^{3+}$

**Official Ans. by NTA (A)**

**Allen Ans. (A)**

**Sol.** Strongest oxidising agent have highest reduction potential value



10. The eutrophication of water body results in :

- (A) loss of Biodiversity  
(B) breakdown of organic matter  
(C) increase in biodiversity  
(D) decrease in BOD.

**Official Ans. by NTA (A)**

**Allen Ans. (A)**

**Sol.** Eutrophication of water body results in loss of Biodiversity.


11. Phenol on reaction with dilute nitric acid, gives two products. Which method will be most effective for large scale separation ?

- (A) Chromatographic separation  
(B) Fractional Crystallisation  
(C) Steam distillation  
(D) Sublimation

**Official Ans. by NTA (C)**

**Allen Ans. (C)**


**Sol.**



Para product has higher boiling point than ortho as intermolecular H-bond is possible in former, whereas intramolecular H-bond is possible in ortho product.

Steam distillation can separate them as ortho product is steam volatile.

12. In the following structures, which one is having staggered conformation with maximum dihedral angle?








**Official Ans. by NTA (C)**

**Allen Ans. (C)**

**Sol.** Dihedral angle : It's the angle b/w 2 specified groups ( $-\text{CH}_3$  here)


Staggered form is Given in option (C) & the angle is  $180^\circ$

**13.** The products formed in the following reaction.



**Official Ans. by NTA (B)**

**Allen Ans. (B)**




**14.** The IUPAC name of ethylidene chloride is :-

- (A) 1-Chloroethene
- (B) 1-Chloroethyne
- (C) 1,2-Dichloroethane
- (D) 1,1-Dichloroethane


**Official Ans. by NTA (D)**

**Allen Ans. (D)**



“1, 1-Dichloroethane is Ethylidene chloride”

**15.** The major product in the reaction




- (A) t-Butyl ethyl ether
- (B) 2,2-Dimethyl butane
- (C) 2-Methyl pent-1-ene
- (D) 2-Methyl prop-1-ene






**Official Ans. by NTA (D)**

**Allen Ans. (D)**

**Sol.** We have been given a bulky base, hence elimination will take place & not substitution.




**16.** The intermediate X, in the reaction



**Official Ans. by NTA (C)**


**Allen Ans. (C)**

Sol. It's a classic Reimer-Tiemann reaction.







Will be the intermediate formed.

17. In the following reaction :




The compounds A and B respectively are :-

- (A) ,  $\text{CH}_3\text{COOH}$
- (B) ,  $\text{CH}_3\text{COOH}$
- (C) ,  $\text{CH}_3\text{COCH}_3$
- (D) ,  $\text{CH}_3\text{COCH}_3$

Official Ans. by NTA (C)

Allen Ans. (C)

Sol. Given reaction is cumene-Peroxide method for the preparation of phenol. In this reaction




18. The reaction of  $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}_2$  with bromine and KOH gives  $\text{RNH}_2$  as the end product. Which one of the following is the intermediate product formed in this reaction ?

- (A)  $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}-\text{Br}$
- (B)  $\text{R}-\text{NH}-\text{Br}$
- (C)  $\text{R}-\text{N}=\text{C}=\text{O}$
- (D)  $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NBr}_2$

Official Ans. by NTA (C)

Allen Ans. (A & C)

Sol. The given reaction is Hoffmann-Bromide degradation method.



19. Using very little soap while washing clothes, does not serve the purpose of cleaning of clothes because
- (A) soap particles remain floating in water as ions  
 (B) the hydrophobic part of soap is not able to take away grease  
 (C) the micelles are not formed due to concentration of soap, below its CMC value  
 (D) colloidal structure of soap in water is completely disturbed.

**Official Ans. by NTA (C)**

**Allen Ans. (C)**

- Sol.** Micelle formation only takes place above CMC.
20. Which one of the following is an example of artificial sweetner ?
- (A) Bithional (B) Alitame  
 (C) Salvarsan (D) Lactose

**Official Ans. by NTA (B)**

**Allen Ans. (B)**

- Sol.** Alitame is a second generation dipeptide sweetner that is 200 times sweeter than sucrose.

### SECTION-B

1. The number of N atoms is 681 g of  $C_7H_5N_3O_6$  is  $x \times 10^{21}$ . The value of x is \_\_\_\_\_ ( $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$ ) (Nearest Integer)

**Official Ans. by NTA (5418)**

**Allen Ans. (5418)**

- Sol.** M.M. of  $C_7H_5N_3O_6$  is  $84 + 5 + 42 + 96 = 227$

$$n_{C_7H_5N_3O_6} = \frac{681}{227} = 3$$

$$n_N = \frac{681}{227} \times 3 = 9 \text{ mol}$$

$$\text{no. of N atoms} = 9 \times 6.02 \times 10^{23}$$

$$= 5418 \times 10^{21}$$

$\therefore$  The answer is 5418.

2. The distance between  $Na^+$  and  $Cl^-$  ions in solid NaCl of density  $43.1 \text{ g cm}^{-3}$  is \_\_\_\_\_  $\times 10^{-10} \text{ m}$ . (Nearest Integer)  
 (Given :  $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$ )

**Official Ans. by NTA (1)**

**Allen Ans. (1)**

- Sol.** Unit cell formula –  $Na_4Cl_4$

$$\text{Mass per unit cell} = \frac{Z \times \text{M.M.}}{N_A} \text{ g}$$

$$= \frac{4 \times 58.5}{N_A} \text{ g}$$

$$d_{\text{unit cell}} = \frac{m}{V} = \frac{m}{a^3}$$

$$\Rightarrow \frac{4 \times 58.5}{N_A \cdot a^3} = 43.1$$

$$\Rightarrow a^3 = 9.02 \times 10^{-24} \text{ cm}^3$$

$$\Rightarrow a = 2.08 \times 10^{-8} \text{ cm}$$

$$\Rightarrow a = 2.08 \times 10^{-10} \text{ m}$$

$$\text{Also } a = 2(r_{Na^+} + r_{Cl^-})$$

$$\Rightarrow r_{Na^+} + r_{Cl^-} = 1.04 \times 10^{-10} \text{ m}$$

$\therefore$  The answer is 1

3. The longest wavelength of light that can be used for the ionisation of lithium atom (Li) in its ground state is  $x \times 10^{-8} \text{ m}$ . The value of x is \_\_\_\_\_. (Nearest Integer)

(Given : Energy of the electron in the first shell of the hydrogen atom is  $-2.2 \times 10^{-18} \text{ J}$ ;  $h = 6.63 \times 10^{-34} \text{ Js}$  and  $c = 3 \times 10^8 \text{ ms}^{-1}$ )

**Official Ans. by NTA (4)**

**Allen Ans. (Bonus)**

- Sol.** We can not calculate I.E. of lithium atom.

4. The standard entropy change for the reaction  $4Fe(s) + 3O_2(g) \rightarrow 2Fe_2O_3(s)$  is  $-550 \text{ JK}^{-1}$  at  $298 \text{ K}$ .

[Given : The standard enthalpy change for the reaction is  $-165 \text{ kJ mol}^{-1}$ ]. The temperature in K at which the reaction attains equilibrium is \_\_\_\_\_. (Nearest Integer)

**Official Ans. by NTA (300)**

**Allen Ans. (300)**

**Sol.**  $\Delta G = \Delta H - T\Delta S = 0$  at equilibrium

$$\Rightarrow -165 \times 10^3 - T \times (-505) = 0$$

$$\Rightarrow T = 300\text{K}$$

The answer is 300

**5.** 1 L aqueous solution of  $\text{H}_2\text{SO}_4$  contains 0.02 m mol  $\text{H}_2\text{SO}_4$ . 50% of this solution is diluted with deionized water to give 1 L solution (A). In solution (A), 0.01 m mol of  $\text{H}_2\text{SO}_4$  are added. Total m mols of  $\text{H}_2\text{SO}_4$  in the final solution is \_\_\_\_\_  $\times 10^3$  m mols.

**Official Ans. by NTA (0)**

**Allen Ans. (0)**

**Sol.**  $n_{\text{H}_2\text{SO}_4}$  in Sol<sup>n</sup> A = 50% of original solution

$$= 0.01 \text{ m mol.}$$

$$n_{\text{H}_2\text{SO}_4} \text{ in Final solution} = 0.01 + 0.01$$

$$= 0.02 \text{ mmol}$$

$$= 0.00002 \times 10^3 \text{ mmol}$$

The answer 0

**6.** The standard free energy change ( $\Delta G^\circ$ ) for 50% dissociation of  $\text{N}_2\text{O}_4$  into  $\text{NO}_2$  at  $27^\circ\text{C}$  and 1 atm pressure is  $-x \text{ J mol}^{-1}$ . The value of x is \_\_\_\_\_. (Nearest Integer)

[Given :  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ ,  $\log 1.33 = 0.1239$   
 $\ln 10 = 2.3$ ]

**Official Ans. by NTA (710)**

**Allen Ans. (710)**

**Sol.**  $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$

$$t = 0 \quad 1 \text{ mol}$$

$$t = t \quad (1-0.5) \text{ mol} \quad 0.5 \times 2 \text{ mol}$$

$$= 0.5 \text{ mol} \quad 1 \text{ mol}$$

$$k_p = \frac{\left(\frac{1}{1.5} \times 1\right)^2}{\left(\frac{0.5}{1.5} \times 1\right)} = \frac{1}{0.75} = \frac{100}{75}$$

$$= 1.33$$

$$\Delta G^\circ = -RT \ln k_p$$

$$= -8.31 \times 300 \times \ln(1.33) = -710.45 \text{ J/mol}$$

$$= -710 \text{ J/mol.}$$

**7.** In a cell, the following reactions take place



The standard electrode potential for the spontaneous reaction in the cell is  $x \times 10^{-2} \text{ V}$  298 K. The value of x is \_\_\_\_\_ (Nearest Integer)

**Official Ans. by NTA (23)**

**Allen Ans. (23)**

**Sol.**  $\text{Fe}^{+3} + \text{I}^- \xrightarrow{\text{Cathode anode}} \text{I}_2 + \text{Fe}^{+2}$


$$E_{\text{Cell}}^\circ = E_{\text{cathode}}^\circ - E_{\text{anode}}^\circ$$

$$= 0.77 - 0.54$$

$$= 0.23$$

$$= 23 \times 10^{-2} \text{ V}$$

**8.** For a given chemical reaction



Concentration of C changes from 10 mmol  $\text{dm}^{-3}$  to 20 mmol  $\text{dm}^{-3}$  in 10 seconds. Rate of appearance of D is 1.5 times the rate of disappearance of B which is twice the rate of disappearance A. The rate of appearance of D has been experimentally determined to be 9 mmol  $\text{dm}^{-3} \text{ s}^{-1}$ . Therefore the rate of reaction is \_\_\_\_\_ mmol  $\text{dm}^{-3} \text{ s}^{-1}$ . (Nearest Integer)

**Official Ans. by NTA (1)**

**Allen Ans. (1)**

**Sol.**  $\gamma_1\text{A} + \gamma_2\text{B} \rightarrow \gamma_3\text{C} + \gamma_4\text{D}$

$$\text{Given: } +\frac{d[\text{D}]}{dt} = \frac{-3}{2} \frac{d[\text{B}]}{dt}$$



$$\Rightarrow \frac{-1}{2} \frac{d[\text{B}]}{dt} = \frac{+1}{3} \frac{d[\text{D}]}{dt}$$

$$-\frac{d[B]}{dt} = -2\frac{d[A]}{dt} \Rightarrow -\frac{1}{2}\frac{d[B]}{dt} = \frac{-d(A)}{dt}$$

$$+\frac{d[B]}{dt} = 9 \text{ mmol dm}^{-3}\text{s}^{-1}$$

$$\frac{+d[C]}{dt} = \frac{20-10}{10} = 1 \text{ mmol dm}^{-3}\text{s}^{-1}$$

$$\frac{+d[C]}{dt} = \frac{1}{9} \times \frac{+d[D]}{dt}$$



$$\text{Rate of reaction} = \frac{+d[C]}{dt} = 1 \text{ mmol dm}^{-3} \text{ s}^{-1}$$

9. If  $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$  absorbs a light of wavelength 600 nm for d-d transition, then the value of octahedral crystal field splitting energy for  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  will be \_\_\_\_\_  $\times 10^{-21}$  J. (Nearest Integer)

(Given :  $h = 6.63 \times 10^{-34}$  Js

and  $c = 3.08 \times 10^8 \text{ ms}^{-1}$ )

**Official Ans. by NTA (746)**

**Allen Ans. (766)**

**Sol.** 
$$\Delta_t = \frac{hc}{\lambda} = \frac{6.63 \times 10^{-34} \times 3.08 \times 10^8}{600 \times 10^{-9}}$$

$$= \frac{6.63 \times 3.08 \times 10^{-17}}{600}$$

$$= 0.034034 \times 10^{-17}$$

$$= 340.34 \times 10^{-21} \text{ J}$$

$$\Delta_0 = \frac{9}{4} \Delta_t$$

$$= \frac{9}{4} \times 340.34 \times 10^{-21}$$

$$= 765.765 \times 10^{-21} \text{ J}$$


$$\approx 766 \times 10^{-21} \text{ J}$$

Answer = 766

10. Number of grams of bromine that will completely react with 5.0g of pent-1-ene is \_\_\_\_\_  $\times 10^{-2}$ g. (Atomic mass of Br = 80 g/mol) [Nearest Integer]

**Official Ans. by NTA (1143)**

**Allen Ans. (1143)**



moles of  $\text{Br}_2$  = moles of  $\text{C}_5\text{H}_{10}$

$$\Rightarrow \frac{w}{160} = \frac{5}{70}$$

$$\Rightarrow w = \frac{5 \times 160}{70} \text{ g}$$

$$= 11.428 \text{ g}$$

$$= 1142.8 \times 10^{-2} \text{ g} \approx 1143 \times 10^{-2} \text{ g}$$