

## 0.1: Physical Constants

Speed of light	$c$	$3 \times 10^8$ m/s
Planck constant	$h$	$6.63 \times 10^{-34}$ J s
	$hc$	1242 eV-nm
Gravitation constant	$G$	$6.67 \times 10^{-11}$ m <sup>3</sup> kg <sup>-1</sup> s <sup>-2</sup>
Boltzmann constant	$k$	$1.38 \times 10^{-23}$ J/K
Molar gas constant	$R$	8.314 J/(mol K)
Avogadro's number	$N_A$	$6.023 \times 10^{23}$ mol <sup>-1</sup>
Charge of electron	$e$	$1.602 \times 10^{-19}$ C
Permeability of vacuum	$\mu_0$	$4\pi \times 10^{-7}$ N/A <sup>2</sup>
Permittivity of vacuum	$\epsilon_0$	$8.85 \times 10^{-12}$ F/m
Coulomb constant	$\frac{1}{4\pi\epsilon_0}$	$9 \times 10^9$ N m <sup>2</sup> /C <sup>2</sup>
Faraday constant	$F$	96485 C/mol
Mass of electron	$m_e$	$9.1 \times 10^{-31}$ kg
Mass of proton	$m_p$	$1.6726 \times 10^{-27}$ kg
Mass of neutron	$m_n$	$1.6749 \times 10^{-27}$ kg
Atomic mass unit	$u$	$1.66 \times 10^{-27}$ kg
Atomic mass unit	$u$	931.49 MeV/c <sup>2</sup>
Stefan-Boltzmann constant	$\sigma$	$5.67 \times 10^{-8}$ W/(m <sup>2</sup> K <sup>4</sup> )
Rydberg constant	$R_\infty$	$1.097 \times 10^7$ m <sup>-1</sup>
Bohr magneton	$\mu_B$	$9.27 \times 10^{-24}$ J/T
Bohr radius	$a_0$	$0.529 \times 10^{-10}$ m
Standard atmosphere	atm	$1.01325 \times 10^5$ Pa
Wien displacement constant	$b$	$2.9 \times 10^{-3}$ m K

# 1 MECHANICS


## 1.1: Vectors

**Notation:**  $\vec{a} = a_x \hat{i} + a_y \hat{j} + a_z \hat{k}$

**Magnitude:**  $a = |\vec{a}| = \sqrt{a_x^2 + a_y^2 + a_z^2}$

**Dot product:**  $\vec{a} \cdot \vec{b} = a_x b_x + a_y b_y + a_z b_z = ab \cos \theta$

**Cross product:**



$$\vec{a} \times \vec{b} = (a_y b_z - a_z b_y) \hat{i} + (a_z b_x - a_x b_z) \hat{j} + (a_x b_y - a_y b_x) \hat{k}$$

$$|\vec{a} \times \vec{b}| = ab \sin \theta$$

## 1.2: Kinematics

**Average and Instantaneous Vel. and Accel.:**

$$\vec{v}_{av} = \Delta \vec{r} / \Delta t,$$

$$\vec{v}_{inst} = d\vec{r}/dt$$

$$\vec{a}_{av} = \Delta \vec{v} / \Delta t$$


$$\vec{a}_{inst} = d\vec{v}/dt$$

**Motion in a straight line with constant  $a$ :**

$$v = u + at, \quad s = ut + \frac{1}{2}at^2, \quad v^2 - u^2 = 2as$$

**Relative Velocity:**  $\vec{v}_{A/B} = \vec{v}_A - \vec{v}_B$

**Projectile Motion:**



$$x = ut \cos \theta, \quad y = ut \sin \theta - \frac{1}{2}gt^2$$

$$y = x \tan \theta - \frac{g}{2u^2 \cos^2 \theta} x^2$$

$$T = \frac{2u \sin \theta}{g}, \quad R = \frac{u^2 \sin 2\theta}{g}, \quad H = \frac{u^2 \sin^2 \theta}{2g}$$

## 1.3: Newton's Laws and Friction

**Linear momentum:**  $\vec{p} = m\vec{v}$

**Newton's first law:** inertial frame.

**Newton's second law:**  $\vec{F} = \frac{d\vec{p}}{dt}, \quad \vec{F} = m\vec{a}$

**Newton's third law:**  $\vec{F}_{AB} = -\vec{F}_{BA}$

**Frictional force:**  $f_{static, max} = \mu_s N, \quad f_{kinetic} = \mu_k N$

**Banking angle:**  $\frac{v^2}{rg} = \tan \theta, \quad \frac{v^2}{rg} = \frac{\mu + \tan \theta}{1 - \mu \tan \theta}$


**Centripetal force:**  $F_c = \frac{mv^2}{r}, \quad a_c = \frac{v^2}{r}$

**Pseudo force:**  $\vec{F}_{pseudo} = -m\vec{a}_0, \quad F_{centrifugal} = -\frac{mv^2}{r}$

**Minimum speed to complete vertical circle:**

$$v_{min, bottom} = \sqrt{5gl}, \quad v_{min, top} = \sqrt{gl}$$

**Conical pendulum:**  $T = 2\pi \sqrt{\frac{l \cos \theta}{g}}$



## 1.4: Work, Power and Energy

**Work:**  $W = \vec{F} \cdot \vec{S} = FS \cos \theta, \quad W = \int \vec{F} \cdot d\vec{S}$

**Kinetic energy:**  $K = \frac{1}{2}mv^2 = \frac{p^2}{2m}$

**Potential energy:**  $F = -\partial U / \partial x$  for conservative forces.

$$U_{gravitational} = mgh, \quad U_{spring} = \frac{1}{2}kx^2$$

**Work done by conservative forces** is path independent and depends only on initial and final points:  
 $\oint \vec{F}_{conservative} \cdot d\vec{r} = 0.$

**Work-energy theorem:**  $W = \Delta K$


**Mechanical energy:**  $E = U + K$ . Conserved if forces are conservative in nature.

**Power**  $P_{av} = \frac{\Delta W}{\Delta t}$ ,  $P_{inst} = \vec{F} \cdot \vec{v}$

**Rotation about an axis with constant  $\alpha$ :**

$$\omega = \omega_0 + \alpha t, \quad \theta = \omega t + \frac{1}{2}\alpha t^2, \quad \omega^2 - \omega_0^2 = 2\alpha\theta$$

**Moment of Inertia:**  $I = \sum_i m_i r_i^2$ ,  $I = \int r^2 dm$



### 1.5: Centre of Mass and Collision

**Centre of mass:**  $x_{cm} = \frac{\sum x_i m_i}{\sum m_i}$ ,  $x_{cm} = \frac{\int x dm}{\int dm}$

**CM of few useful configurations:**


- $m_1, m_2$  separated by  $r$ :
- Triangle (CM  $\equiv$  Centroid)  $y_c = \frac{h}{3}$
- Semicircular ring:  $y_c = \frac{2r}{\pi}$
- Semicircular disc:  $y_c = \frac{4r}{3\pi}$
- Hemispherical shell:  $y_c = \frac{r}{2}$
- Solid Hemisphere:  $y_c = \frac{3r}{8}$
- Cone: the height of CM from the base is  $h/4$  for the solid cone and  $h/3$  for the hollow cone.

**Motion of the CM:**  $M = \sum m_i$

$$\vec{v}_{cm} = \frac{\sum m_i \vec{v}_i}{M}, \quad \vec{p}_{cm} = M \vec{v}_{cm}, \quad \vec{a}_{cm} = \frac{\vec{F}_{ext}}{M}$$

**Impulse:**  $\vec{J} = \int \vec{F} dt = \Delta \vec{p}$

**Collision:**



Momentum conservation:  $m_1 v_1 + m_2 v_2 = m_1 v'_1 + m_2 v'_2$

Elastic Collision:  $\frac{1}{2} m_1 v_1^2 + \frac{1}{2} m_2 v_2^2 = \frac{1}{2} m_1 v'^2_1 + \frac{1}{2} m_2 v'^2_2$

Coefficient of restitution:


$$e = \frac{-(v'_1 - v'_2)}{v_1 - v_2} = \begin{cases} 1, & \text{completely elastic} \\ 0, & \text{completely in-elastic} \end{cases}$$

If  $v_2 = 0$  and  $m_1 \ll m_2$  then  $v'_1 = -v_1$ .


If  $v_2 = 0$  and  $m_1 \gg m_2$  then  $v'_2 = 2v_1$ .

Elastic collision with  $m_1 = m_2$ :  $v'_1 = v_2$  and  $v'_2 = v_1$ .

**Theorem of Parallel Axes:**  $I_{||} = I_{cm} + md^2$




**Theorem of Perp. Axes:**  $I_z = I_x + I_y$



**Radius of Gyration:**  $k = \sqrt{I/m}$

**Angular Momentum:**  $\vec{L} = \vec{r} \times \vec{p}$ ,  $\vec{L} = I\vec{\omega}$

**Torque:**  $\vec{\tau} = \vec{r} \times \vec{F}$ ,  $\vec{\tau} = \frac{d\vec{L}}{dt}$ ,  $\tau = I\alpha$



**Conservation of  $\vec{L}$ :**  $\vec{\tau}_{ext} = 0 \implies \vec{L} = \text{const.}$

**Equilibrium condition:**  $\sum \vec{F} = \vec{0}$ ,  $\sum \vec{\tau} = \vec{0}$

**Kinetic Energy:**  $K_{rot} = \frac{1}{2} I \omega^2$


**Dynamics:**

$$\vec{\tau}_{cm} = I_{cm} \vec{\alpha}, \quad \vec{F}_{ext} = m \vec{a}_{cm}, \quad \vec{p}_{cm} = m \vec{v}_{cm}$$

$$K = \frac{1}{2} m v_{cm}^2 + \frac{1}{2} I_{cm} \omega^2, \quad \vec{L} = I_{cm} \vec{\omega} + \vec{r}_{cm} \times m \vec{v}_{cm}$$

### 1.7: Gravitation

**Gravitational force:**  $F = G \frac{m_1 m_2}{r^2}$



**Potential energy:**  $U = -\frac{GMm}{r}$

**Gravitational acceleration:**  $g = \frac{GM}{R^2}$

**Variation of g with depth:**  $g_{inside} \approx g \left(1 - \frac{h}{R}\right)$

**Variation of g with height:**  $g_{outside} \approx g \left(1 - \frac{2h}{R}\right)$

**Effect of non-spherical earth shape on g:**


$$g_{at \text{ pole}} > g_{at \text{ equator}} \quad (\because R_e - R_p \approx 21 \text{ km})$$

**Effect of earth rotation on apparent weight:**

### 1.6: Rigid Body Dynamics

**Angular velocity:**  $\omega_{av} = \frac{\Delta \theta}{\Delta t}$ ,  $\omega = \frac{d\theta}{dt}$ ,  $\vec{v} = \vec{\omega} \times \vec{r}$


**Angular Accel.:**  $\alpha_{av} = \frac{\Delta \omega}{\Delta t}$ ,  $\alpha = \frac{d\omega}{dt}$ ,  $\vec{a} = \vec{\alpha} \times \vec{r}$



**Orbital velocity of satellite:**  $v_o = \sqrt{\frac{GM}{R}}$

**Escape velocity:**  $v_e = \sqrt{\frac{2GM}{R}}$

**Kepler's laws:**




**First:** Elliptical orbit with sun at one of the focus.

**Second:** Areal velocity is constant. ( $\therefore d\vec{L}/dt = 0$ ).

**Third:**  $T^2 \propto a^3$ . In circular orbit  $T^2 = \frac{4\pi^2}{GM} a^3$ .

**Superposition of two SHM's:**



$x_1 = A_1 \sin \omega t, \quad x_2 = A_2 \sin(\omega t + \delta)$

$x = x_1 + x_2 = A \sin(\omega t + \epsilon)$

$A = \sqrt{A_1^2 + A_2^2 + 2A_1A_2 \cos \delta}$

$\tan \epsilon = \frac{A_2 \sin \delta}{A_1 + A_2 \cos \delta}$

**1.8: Simple Harmonic Motion**

**Hooke's law:**  $F = -kx$  (for small elongation  $x$ .)


**Acceleration:**  $a = \frac{d^2x}{dt^2} = -\frac{k}{m}x = -\omega^2x$

**Time period:**  $T = \frac{2\pi}{\omega} = 2\pi\sqrt{\frac{m}{k}}$


**Displacement:**  $x = A \sin(\omega t + \phi)$

**Velocity:**  $v = A\omega \cos(\omega t + \phi) = \pm\omega\sqrt{A^2 - x^2}$

**Potential energy:**  $U = \frac{1}{2}kx^2$




**Kinetic energy**  $K = \frac{1}{2}mv^2$



**Total energy:**  $E = U + K = \frac{1}{2}m\omega^2A^2$


**Simple pendulum:**  $T = 2\pi\sqrt{\frac{l}{g}}$




**Physical Pendulum:**  $T = 2\pi\sqrt{\frac{I}{mgl}}$




**Torsional Pendulum**  $T = 2\pi\sqrt{\frac{I}{k}}$



**Springs in series:**  $\frac{1}{k_{eq}} = \frac{1}{k_1} + \frac{1}{k_2}$



**Springs in parallel:**  $k_{eq} = k_1 + k_2$



**1.9: Properties of Matter**

**Modulus of rigidity:**  $Y = \frac{F/A}{\Delta l/l}, \quad B = -V \frac{\Delta P}{\Delta V}, \quad \eta = \frac{F}{A\theta}$

**Compressibility:**  $K = \frac{1}{B} = -\frac{1}{V} \frac{dV}{dP}$

**Poisson's ratio:**  $\sigma = \frac{\text{lateral strain}}{\text{longitudinal strain}} = \frac{\Delta D/D}{\Delta l/l}$

**Elastic energy:**  $U = \frac{1}{2} \text{stress} \times \text{strain} \times \text{volume}$

**Surface tension:**  $S = F/l$

**Surface energy:**  $U = SA$

**Excess pressure in bubble:**

$\Delta p_{\text{air}} = 2S/R, \quad \Delta p_{\text{soap}} = 4S/R$

**Capillary rise:**  $h = \frac{2S \cos \theta}{r\rho g}$

**Hydrostatic pressure:**  $p = \rho gh$

**Buoyant force:**  $F_B = \rho Vg = \text{Weight of displaced liquid}$

**Equation of continuity:**  $A_1v_1 = A_2v_2$




**Bernoulli's equation:**  $p + \frac{1}{2}\rho v^2 + \rho gh = \text{constant}$


**Torricelli's theorem:**  $v_{\text{efflux}} = \sqrt{2gh}$

**Viscous force:**  $F = -\eta A \frac{dv}{dx}$

**Stoke's law:**  $F = 6\pi\eta r v$



**Poiseuille's equation:**  $\frac{\text{Volume flow}}{\text{time}} = \frac{\pi p r^4}{8\eta l}$



**Terminal velocity:**  $v_t = \frac{2r^2(\rho - \sigma)g}{9\eta}$

## 2 Waves

### 2.1: Waves Motion

**General equation of wave:**  $\frac{\partial^2 y}{\partial x^2} = \frac{1}{v^2} \frac{\partial^2 y}{\partial t^2}$ .

**Notation:** Amplitude  $A$ , Frequency  $\nu$ , Wavelength  $\lambda$ , Period  $T$ , Angular Frequency  $\omega$ , Wave Number  $k$ ,


$$T = \frac{1}{\nu} = \frac{2\pi}{\omega}, \quad v = \nu\lambda, \quad k = \frac{2\pi}{\lambda}$$



**Progressive wave travelling with speed  $v$ :**

$$y = f(t - x/v), \rightsquigarrow +x; \quad y = f(t + x/v), \rightsquigarrow -x$$


**Progressive sine wave:**




$$y = A \sin(kx - \omega t) = A \sin(2\pi(x/\lambda - t/T))$$



4. 1<sup>st</sup> overtone/2<sup>nd</sup> harmonics:  $\nu_1 = \frac{2}{2L} \sqrt{\frac{T}{\mu}}$  
5. 2<sup>nd</sup> overtone/3<sup>rd</sup> harmonics:  $\nu_2 = \frac{3}{2L} \sqrt{\frac{T}{\mu}}$  
6. All harmonics are present.

**String fixed at one end:**



1. Boundary conditions:  $y = 0$  at  $x = 0$
2. Allowed Freq.:  $L = (2n + 1) \frac{\lambda}{4}$ ,  $\nu = \frac{2n+1}{4L} \sqrt{\frac{T}{\mu}}$ ,  $n = 0, 1, 2, \dots$
3. Fundamental/1<sup>st</sup> harmonics:  $\nu_0 = \frac{1}{4L} \sqrt{\frac{T}{\mu}}$  
4. 1<sup>st</sup> overtone/3<sup>rd</sup> harmonics:  $\nu_1 = \frac{3}{4L} \sqrt{\frac{T}{\mu}}$  
5. 2<sup>nd</sup> overtone/5<sup>th</sup> harmonics:  $\nu_2 = \frac{5}{4L} \sqrt{\frac{T}{\mu}}$  
6. Only odd harmonics are present.

### 2.2: Waves on a String

**Speed of waves** on a string with mass per unit length  $\mu$  and tension  $T$ :  $v = \sqrt{T/\mu}$

**Transmitted power:**  $P_{av} = 2\pi^2 \mu v A^2 \nu^2$

**Interference:**

$$y_1 = A_1 \sin(kx - \omega t), \quad y_2 = A_2 \sin(kx - \omega t + \delta)$$


$$y = y_1 + y_2 = A \sin(kx - \omega t + \epsilon)$$

$$A = \sqrt{A_1^2 + A_2^2 + 2A_1 A_2 \cos \delta}$$

$$\tan \epsilon = \frac{A_2 \sin \delta}{A_1 + A_2 \cos \delta}$$

$$\delta = \begin{cases} 2n\pi, & \text{constructive;} \\ (2n+1)\pi, & \text{destructive.} \end{cases}$$

**Standing Waves:**





$$y_1 = A_1 \sin(kx - \omega t), \quad y_2 = A_2 \sin(kx + \omega t)$$

$$y = y_1 + y_2 = (2A \cos kx) \sin \omega t$$

$$x = \begin{cases} (n + \frac{1}{2}) \frac{\lambda}{2}, & \text{nodes; } n = 0, 1, 2, \dots \\ n \frac{\lambda}{2}, & \text{antinodes. } n = 0, 1, 2, \dots \end{cases}$$

**String fixed at both ends:**



1. Boundary conditions:  $y = 0$  at  $x = 0$  and at  $x = L$
2. Allowed Freq.:  $L = n \frac{\lambda}{2}$ ,  $\nu = \frac{n}{2L} \sqrt{\frac{T}{\mu}}$ ,  $n = 1, 2, 3, \dots$
3. Fundamental/1<sup>st</sup> harmonics:  $\nu_0 = \frac{1}{2L} \sqrt{\frac{T}{\mu}}$  

### 2.3: Sound Waves

**Displacement wave:**  $s = s_0 \sin \omega(t - x/v)$

**Pressure wave:**  $p = p_0 \cos \omega(t - x/v)$ ,  $p_0 = (B\omega/v)s_0$

**Speed of sound waves:**

$$v_{\text{liquid}} = \sqrt{\frac{B}{\rho}}, \quad v_{\text{solid}} = \sqrt{\frac{Y}{\rho}}, \quad v_{\text{gas}} = \sqrt{\frac{\gamma P}{\rho}}$$


**Intensity:**  $I = \frac{2\pi^2 B}{v} s_0^2 \nu^2 = \frac{p_0^2 v}{2B} = \frac{p_0^2}{2\rho v}$



**Standing longitudinal waves:**

$$p_1 = p_0 \sin \omega(t - x/v), \quad p_2 = p_0 \sin \omega(t + x/v)$$

$$p = p_1 + p_2 = 2p_0 \cos kx \sin \omega t$$

**Closed organ pipe:**




1. Boundary condition:  $y = 0$  at  $x = 0$
2. Allowed freq.:  $L = (2n + 1) \frac{\lambda}{4}$ ,  $\nu = (2n + 1) \frac{v}{4L}$ ,  $n = 0, 1, 2, \dots$
3. Fundamental/1<sup>st</sup> harmonics:  $\nu_0 = \frac{v}{4L}$  
4. 1<sup>st</sup> overtone/3<sup>rd</sup> harmonics:  $\nu_1 = 3\nu_0 = \frac{3v}{4L}$  


5. 2<sup>nd</sup> overtone/5<sup>th</sup> harmonics:  $\nu_2 = 5\nu_0 = \frac{5v}{4L}$  


6. Only odd harmonics are present.


**Open organ pipe:**



1. Boundary condition:  $y = 0$  at  $x = 0$   
 Allowed freq.:  $L = n\frac{\lambda}{2}$ ,  $\nu = n\frac{v}{4L}$ ,  $n = 1, 2, \dots$


2. Fundamental/1<sup>st</sup> harmonics:  $\nu_0 = \frac{v}{2L}$  

3. 1<sup>st</sup> overtone/2<sup>nd</sup> harmonics:  $\nu_1 = 2\nu_0 = \frac{2v}{2L}$  

4. 2<sup>nd</sup> overtone/3<sup>rd</sup> harmonics:  $\nu_2 = 3\nu_0 = \frac{3v}{2L}$  

5. All harmonics are present.

**Resonance column:**



$$l_1 + d = \frac{\lambda}{2}, \quad l_2 + d = \frac{3\lambda}{4}, \quad v = 2(l_2 - l_1)\nu$$

**Beats:** two waves of almost equal frequencies  $\omega_1 \approx \omega_2$

$$p_1 = p_0 \sin \omega_1(t - x/v), \quad p_2 = p_0 \sin \omega_2(t - x/v)$$

$$p = p_1 + p_2 = 2p_0 \cos \Delta\omega(t - x/v) \sin \omega(t - x/v)$$


$$\omega = (\omega_1 + \omega_2)/2, \quad \Delta\omega = \omega_1 - \omega_2 \quad (\text{beats freq.})$$

**Doppler Effect:**

$$\nu = \frac{v + u_o}{v - u_s} \nu_0$$

where,  $v$  is the speed of sound in the medium,  $u_o$  is the speed of the observer w.r.t. the medium, considered positive when it moves towards the source and negative when it moves away from the source, and  $u_s$  is the speed of the source w.r.t. the medium, considered positive when it moves towards the observer and negative when it moves away from the observer.

**Path difference:**  $\Delta x = \frac{dy}{D}$



**Phase difference:**  $\delta = \frac{2\pi}{\lambda} \Delta x$

**Interference Conditions:** for integer  $n$ ,

$$\delta = \begin{cases} 2n\pi, & \text{constructive;} \\ (2n + 1)\pi, & \text{destructive,} \end{cases}$$

$$\Delta x = \begin{cases} n\lambda, & \text{constructive;} \\ (n + \frac{1}{2})\lambda, & \text{destructive} \end{cases}$$

**Intensity:**

$$I = I_1 + I_2 + 2\sqrt{I_1 I_2} \cos \delta,$$

$$I_{\max} = (\sqrt{I_1} + \sqrt{I_2})^2, \quad I_{\min} = (\sqrt{I_1} - \sqrt{I_2})^2$$

$$I_1 = I_2 : I = 4I_0 \cos^2 \frac{\delta}{2}, \quad I_{\max} = 4I_0, \quad I_{\min} = 0$$


**Fringe width:**  $w = \frac{\lambda D}{d}$

**Optical path:**  $\Delta x' = \mu \Delta x$

**Interference of waves transmitted through thin film:**

$$\Delta x = 2\mu d = \begin{cases} n\lambda, & \text{constructive;} \\ (n + \frac{1}{2})\lambda, & \text{destructive.} \end{cases}$$


**Diffraction from a single slit:**




For Minima:  $n\lambda = b \sin \theta \approx b(y/D)$


**Resolution:**  $\sin \theta = \frac{1.22\lambda}{b}$

**Law of Malus:**  $I = I_0 \cos^2 \theta$



**2.4: Light Waves**

**Plane Wave:**  $E = E_0 \sin \omega(t - \frac{x}{v})$ ,  $I = I_0$  


**Spherical Wave:**  $E = \frac{aE_0}{r} \sin \omega(t - \frac{r}{v})$ ,  $I = \frac{I_0}{r^2}$  

**Young's double slit experiment**

### 3 Optics


#### 3.1: Reflection of Light

**Laws of reflection:**




Incident ray, reflected ray, and normal lie in the same plane (ii)  $\angle i = \angle r$

**Plane mirror:**



(i) the image and the object are equidistant from mirror (ii) virtual image of real object

**Spherical Mirror:**




1. Focal length  $f = R/2$
2. Mirror equation:  $\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$
3. Magnification:  $m = -\frac{v}{u}$


#### 3.2: Refraction of Light

**Refractive index:**  $\mu = \frac{\text{speed of light in vacuum}}{\text{speed of light in medium}} = \frac{c}{v}$


**Snell's Law:**  $\frac{\sin i}{\sin r} = \frac{\mu_2}{\mu_1}$




**Apparent depth:**  $\mu = \frac{\text{real depth}}{\text{apparent depth}} = \frac{d}{d'}$



**Critical angle:**  $\theta_c = \sin^{-1} \frac{1}{\mu}$




**Deviation by a prism:**



$\delta = i + i' - A$ , general result  
 $\mu = \frac{\sin \frac{A + \delta_m}{2}}{\sin \frac{A}{2}}$ ,  $i = i'$  for minimum deviation

$\delta_m = (\mu - 1)A$ , for small A


**Refraction at spherical surface:**



$\frac{\mu_2}{v} - \frac{\mu_1}{u} = \frac{\mu_2 - \mu_1}{R}$ ,  $m = \frac{\mu_1 v}{\mu_2 u}$

**Lens maker's formula:**  $\frac{1}{f} = (\mu - 1) \left[ \frac{1}{R_1} - \frac{1}{R_2} \right]$

**Lens formula:**  $\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$ ,  $m = \frac{v}{u}$



**Power of the lens:**  $P = \frac{1}{f}$ , P in diopter if f in metre.


**Two thin lenses separated by distance d:**

$\frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2} - \frac{d}{f_1 f_2}$

#### 3.3: Optical Instruments


**Simple microscope:**  $m = D/f$  in normal adjustment.

**Compound microscope:**



1. Magnification in normal adjustment:  $m = \frac{v}{u} \frac{D}{f_e}$
2. Resolving power:  $R = \frac{1}{\Delta d} = \frac{2\mu \sin \theta}{\lambda}$

**Astronomical telescope:**



1. In normal adjustment:  $m = -\frac{f_o}{f_e}$ ,  $L = f_o + f_e$
2. Resolving power:  $R = \frac{1}{\Delta \theta} = \frac{1}{1.22 \lambda}$

#### 3.4: Dispersion

**Cauchy's equation:**  $\mu = \mu_0 + \frac{A}{\lambda^2}$ ,  $A > 0$


**Dispersion by prism with small A and i:**

1. Mean deviation:  $\delta_y = (\mu_y - 1)A$
2. Angular dispersion:  $\theta = (\mu_v - \mu_r)A$

**Dispersive power:**  $\omega = \frac{\mu_v - \mu_r}{\mu_y - 1} \approx \frac{\theta}{\delta_y}$  (if A and i small)

**Dispersion without deviation:**

$(\mu_y - 1)A + (\mu'_y - 1)A' = 0$



**Deviation without dispersion:**

$(\mu_v - \mu_r)A = (\mu'_v - \mu'_r)A'$

## 4 Heat and Thermodynamics

### 4.1: Heat and Temperature

Temp. scales:  $F = 32 + \frac{9}{5}C$ ,  $K = C + 273.16$

Ideal gas equation:  $pV = nRT$ ,  $n$ : number of moles

van der Waals equation:  $(p + \frac{a}{V^2})(V - b) = nRT$


Thermal expansion:  $L = L_0(1 + \alpha\Delta T)$ ,  
 $A = A_0(1 + \beta\Delta T)$ ,  $V = V_0(1 + \gamma\Delta T)$ ,  $\gamma = 2\beta = 3\alpha$

Thermal stress of a material:  $\frac{F}{A} = Y \frac{\Delta L}{L}$

### 4.2: Kinetic Theory of Gases

General:  $M = mN_A$ ,  $k = R/N_A$

Maxwell distribution of speed:



RMS speed:  $v_{rms} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}}$

Average speed:  $\bar{v} = \sqrt{\frac{8kT}{\pi m}} = \sqrt{\frac{8RT}{\pi M}}$

Most probable speed:  $v_p = \sqrt{\frac{2kT}{m}}$

Pressure:  $p = \frac{1}{3}\rho v_{rms}^2$

Equipartition of energy:  $K = \frac{1}{2}kT$  for each degree of freedom. Thus,  $K = \frac{f}{2}kT$  for molecule having  $f$  degrees of freedoms.

Internal energy of  $n$  moles of an ideal gas is  $U = \frac{f}{2}nRT$ .

### 4.3: Specific Heat

Specific heat:  $s = \frac{Q}{m\Delta T}$

Latent heat:  $L = Q/m$

Specific heat at constant volume:  $C_v = \left. \frac{\Delta Q}{n\Delta T} \right|_V$

Specific heat at constant pressure:  $C_p = \left. \frac{\Delta Q}{n\Delta T} \right|_p$

Relation between  $C_p$  and  $C_v$ :  $C_p - C_v = R$

Ratio of specific heats:  $\gamma = C_p/C_v$

Relation between  $U$  and  $C_v$ :  $\Delta U = nC_v\Delta T$

Specific heat of gas mixture:

$$C_v = \frac{n_1C_{v1} + n_2C_{v2}}{n_1 + n_2}, \quad \gamma = \frac{n_1C_{p1} + n_2C_{p2}}{n_1C_{v1} + n_2C_{v2}}$$

Molar internal energy of an ideal gas:  $U = \frac{f}{2}RT$ ,  
 $f = 3$  for monatomic and  $f = 5$  for diatomic gas.

### 4.4: Thermodynamic Processes

First law of thermodynamics:  $\Delta Q = \Delta U + \Delta W$

Work done by the gas:

$$\Delta W = p\Delta V, \quad W = \int_{V_1}^{V_2} pdV$$


$$W_{\text{isothermal}} = nRT \ln \left( \frac{V_2}{V_1} \right)$$

$$W_{\text{isobaric}} = p(V_2 - V_1)$$

$$W_{\text{adiabatic}} = \frac{p_1V_1 - p_2V_2}{\gamma - 1}$$

$$W_{\text{isochoric}} = 0$$


Efficiency of the heat engine:



$$\eta = \frac{\text{work done by the engine}}{\text{heat supplied to it}} = \frac{Q_1 - Q_2}{Q_1}$$

$$\eta_{\text{carnot}} = 1 - \frac{Q_2}{Q_1} = 1 - \frac{T_2}{T_1}$$

Coeff. of performance of refrigerator:



$$\text{COP} = \frac{Q_2}{W} = \frac{Q_2}{Q_1 - Q_2}$$

Entropy:  $\Delta S = \frac{\Delta Q}{T}$ ,  $S_f - S_i = \int_i^f \frac{\Delta Q}{T}$

$$\text{Const. } T : \Delta S = \frac{Q}{T}, \quad \text{Varying } T : \Delta S = ms \ln \frac{T_f}{T_i}$$


Adiabatic process:  $\Delta Q = 0$ ,  $pV^\gamma = \text{constant}$

### 4.5: Heat Transfer


Conduction:  $\frac{\Delta Q}{\Delta t} = -KA \frac{\Delta T}{x}$

Thermal resistance:  $R = \frac{x}{KA}$

$$R_{\text{series}} = R_1 + R_2 = \frac{1}{A} \left( \frac{x_1}{K_1} + \frac{x_2}{K_2} \right)$$




$$\frac{1}{R_{\text{parallel}}} = \frac{1}{R_1} + \frac{1}{R_2} = \frac{1}{x} (K_1A_1 + K_2A_2)$$



Kirchhoff's Law:  $\frac{\text{emissive power}}{\text{absorptive power}} = \frac{E_{\text{body}}}{a_{\text{body}}} = E_{\text{blackbody}}$

Wien's displacement law:  $\lambda_m T = b$




Stefan-Boltzmann law:  $\frac{\Delta Q}{\Delta t} = \sigma eAT^4$

Newton's law of cooling:  $\frac{dT}{dt} = -bA(T - T_0)$


# 5 Electricity and Magnetism

## 5.1: Electrostatics

Coulomb's law:  $\vec{F} = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2} \hat{r}$



Electric field:  $\vec{E}(\vec{r}) = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{r}$




Electrostatic energy:  $U = -\frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r}$


Electrostatic potential:  $V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$

$$dV = -\vec{E} \cdot d\vec{r}, \quad V(\vec{r}) = -\int_{\infty}^{\vec{r}} \vec{E} \cdot d\vec{r}$$

Electric dipole moment:  $\vec{p} = q\vec{d}$




Potential of a dipole:  $V = \frac{1}{4\pi\epsilon_0} \frac{p \cos \theta}{r^2}$



Field of a dipole:

$$E_r = \frac{1}{4\pi\epsilon_0} \frac{2p \cos \theta}{r^3}, \quad E_\theta = \frac{1}{4\pi\epsilon_0} \frac{p \sin \theta}{r^3}$$



Torque on a dipole placed in  $\vec{E}$ :  $\vec{\tau} = \vec{p} \times \vec{E}$

Pot. energy of a dipole placed in  $\vec{E}$ :  $U = -\vec{p} \cdot \vec{E}$


## 5.2: Gauss's Law and its Applications

Electric flux:  $\phi = \oint \vec{E} \cdot d\vec{S}$

Gauss's law:  $\oint \vec{E} \cdot d\vec{S} = q_{in}/\epsilon_0$


Field of a uniformly charged ring on its axis:

$$E_P = \frac{1}{4\pi\epsilon_0} \frac{qx}{(a^2+x^2)^{3/2}}$$




E and V of a uniformly charged sphere:

$$E = \begin{cases} \frac{1}{4\pi\epsilon_0} \frac{Qr}{R^3}, & \text{for } r < R \\ \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}, & \text{for } r \geq R \end{cases}$$




$$V = \begin{cases} \frac{Q}{8\pi\epsilon_0 R} \left(3 - \frac{r^2}{R^2}\right), & \text{for } r < R \\ \frac{1}{4\pi\epsilon_0} \frac{Q}{r}, & \text{for } r \geq R \end{cases}$$




E and V of a uniformly charged spherical shell:

$$E = \begin{cases} 0, & \text{for } r < R \\ \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}, & \text{for } r \geq R \end{cases}$$



$$V = \begin{cases} \frac{1}{4\pi\epsilon_0} \frac{Q}{R}, & \text{for } r < R \\ \frac{1}{4\pi\epsilon_0} \frac{Q}{r}, & \text{for } r \geq R \end{cases}$$



Field of a line charge:  $E = \frac{\lambda}{2\pi\epsilon_0 r}$


Field of an infinite sheet:  $E = \frac{\sigma}{2\epsilon_0}$

Field in the vicinity of conducting surface:  $E = \frac{\sigma}{\epsilon_0}$


## 5.3: Capacitors

Capacitance:  $C = q/V$


Parallel plate capacitor:  $C = \epsilon_0 A/d$




Spherical capacitor:  $C = \frac{4\pi\epsilon_0 r_1 r_2}{r_2 - r_1}$




Cylindrical capacitor:  $C = \frac{2\pi\epsilon_0 l}{\ln(r_2/r_1)}$



Capacitors in parallel:  $C_{eq} = C_1 + C_2$



Capacitors in series:  $\frac{1}{C_{eq}} = \frac{1}{C_1} + \frac{1}{C_2}$



Force between plates of a parallel plate capacitor:

$$F = \frac{Q^2}{2A\epsilon_0}$$

Energy stored in capacitor:  $U = \frac{1}{2} CV^2 = \frac{Q^2}{2C} = \frac{1}{2} QV$

Energy density in electric field E:  $U/V = \frac{1}{2} \epsilon_0 E^2$

Capacitor with dielectric:  $C = \frac{\epsilon_0 K A}{d}$

## 5.4: Current electricity

Current density:  $j = i/A = \sigma E$

Drift speed:  $v_d = \frac{1}{2} \frac{eE}{m} \tau = \frac{i}{neA}$


Resistance of a wire:  $R = \rho l/A$ , where  $\rho = 1/\sigma$

Temp. dependence of resistance:  $R = R_0(1 + \alpha \Delta T)$


Ohm's law:  $V = iR$

**Kirchhoff's Laws:** (i) *The Junction Law:* The algebraic sum of all the currents directed towards a node is zero i.e.,  $\sum_{\text{node}} I_i = 0$ . (ii) *The Loop Law:* The algebraic sum of all the potential differences along a closed loop in a circuit is zero i.e.,  $\sum_{\text{loop}} \Delta V_i = 0$ .


Resistors in parallel:  $\frac{1}{R_{eq}} = \frac{1}{R_1} + \frac{1}{R_2}$



Resistors in series:  $R_{eq} = R_1 + R_2$



Wheatstone bridge:




Balanced if  $R_1/R_2 = R_3/R_4$ .

Electric Power:  $P = V^2/R = I^2 R = IV$




**Galvanometer as an Ammeter:**




$$i_g G = (i - i_g) S$$

**Galvanometer as a Voltmeter:**




$$V_{AB} = i_g (R + G)$$

**Charging of capacitors:**



$$q(t) = CV \left[ 1 - e^{-\frac{t}{RC}} \right]$$


**Discharging of capacitors:**  $q(t) = q_0 e^{-\frac{t}{RC}}$



**Time constant in RC circuit:**  $\tau = RC$

**Peltier effect:**  $\text{emf } e = \frac{\Delta H}{\Delta Q} = \frac{\text{Peltier heat}}{\text{charge transferred}}$

**Seebeck effect:**



1. Thermo-emf:  $e = aT + \frac{1}{2}bT^2$
2. Thermoelectric power:  $de/dt = a + bT$ .
3. Neutral temp.:  $T_n = -a/b$ .
4. Inversion temp.:  $T_i = -2a/b$ .

**Thomson effect:**  $\text{emf } e = \frac{\Delta H}{\Delta Q} = \frac{\text{Thomson heat}}{\text{charge transferred}} = \sigma \Delta T$ .

**Faraday's law of electrolysis:** The mass deposited is


$$m = Zit = \frac{1}{F} Eit$$

where  $i$  is current,  $t$  is time,  $Z$  is electrochemical equivalent,  $E$  is chemical equivalent, and  $F = 96485 \text{ C/g}$  is Faraday constant.

**Energy of a magnetic dipole placed in  $\vec{B}$ :**


$$U = -\vec{\mu} \cdot \vec{B}$$

**Hall effect:**  $V_w = \frac{Bi}{ned}$




## 5.6: Magnetic Field due to Current

**Biot-Savart law:**  $d\vec{B} = \frac{\mu_0}{4\pi} \frac{i d\vec{l} \times \vec{r}}{r^3}$



**Field due to a straight conductor:**




$$B = \frac{\mu_0 i}{4\pi d} (\cos \theta_1 - \cos \theta_2)$$

**Field due to an infinite straight wire:**  $B = \frac{\mu_0 i}{2\pi d}$

**Force between parallel wires:**  $\frac{dF}{dl} = \frac{\mu_0 i_1 i_2}{2\pi d}$




**Field on the axis of a ring:**



$$B_P = \frac{\mu_0 i a^2}{2(a^2 + d^2)^{3/2}}$$


**Field at the centre of an arc:**  $B = \frac{\mu_0 i \theta}{4\pi a}$




**Field at the centre of a ring:**  $B = \frac{\mu_0 i}{2a}$

**Ampere's law:**  $\oint \vec{B} \cdot d\vec{l} = \mu_0 I_{in}$


**Field inside a solenoid:**  $B = \mu_0 n i$ ,  $n = \frac{N}{l}$



**Field inside a toroid:**  $B = \frac{\mu_0 N i}{2\pi r}$




**Field of a bar magnet:**



$$B_1 = \frac{\mu_0}{4\pi} \frac{2M}{d^3}, \quad B_2 = \frac{\mu_0}{4\pi} \frac{M}{d^3}$$

**Angle of dip:**  $B_h = B \cos \delta$



**Tangent galvanometer:**  $B_h \tan \theta = \frac{\mu_0 n i}{2r}$ ,  $i = K \tan \theta$

**Moving coil galvanometer:**  $n i A B = k \theta$ ,  $i = \frac{k}{n A B} \theta$


**Time period of magnetometer:**  $T = 2\pi \sqrt{\frac{I}{M B_h}}$

**Permeability:**  $\vec{B} = \mu \vec{H}$


## 5.5: Magnetism

**Lorentz force on a moving charge:**  $\vec{F} = q\vec{v} \times \vec{B} + q\vec{E}$

**Charged particle in a uniform magnetic field:**




**Force on a current carrying wire:**



$$\vec{F} = i \vec{l} \times \vec{B}$$

**Magnetic moment of a current loop (dipole):**



**Torque on a magnetic dipole placed in  $\vec{B}$ :**  $\vec{\tau} = \vec{\mu} \times \vec{B}$


## 5.7: Electromagnetic Induction

**Magnetic flux:**  $\phi = \oint \vec{B} \cdot d\vec{S}$

**Faraday's law:**  $e = -\frac{d\phi}{dt}$

**Lenz's Law:** Induced current create a  $B$ -field that opposes the change in magnetic flux.


**Motional emf:**  $e = Blv$




**Self inductance:**  $\phi = Li$ ,  $e = -L\frac{di}{dt}$

**Self inductance of a solenoid:**  $L = \mu_0 n^2 (\pi r^2 l)$

**Growth of current in LR circuit:**  $i = \frac{e}{R} [1 - e^{-t/L/R}]$



**Decay of current in LR circuit:**  $i = i_0 e^{-t/L/R}$



**Time constant of LR circuit:**  $\tau = L/R$


**Energy stored in an inductor:**  $U = \frac{1}{2} Li^2$

**Energy density of B field:**  $u = \frac{U}{V} = \frac{B^2}{2\mu_0}$

**Mutual inductance:**  $\phi = Mi$ ,  $e = -M\frac{di}{dt}$

**EMF induced in a rotating coil:**  $e = NAB\omega \sin \omega t$


**Alternating current:**



$$i = i_0 \sin(\omega t + \phi), \quad T = 2\pi/\omega$$

**Average current in AC:**  $\bar{i} = \frac{1}{T} \int_0^T i dt = 0$

**RMS current:**  $i_{\text{rms}} = \left[ \frac{1}{T} \int_0^T i^2 dt \right]^{1/2} = \frac{i_0}{\sqrt{2}}$




**Energy:**  $E = i_{\text{rms}}^2 RT$

**Capacitive reactance:**  $X_c = \frac{1}{\omega C}$

**Inductive reactance:**  $X_L = \omega L$


**Impedance:**  $Z = e_0/i_0$

**RC circuit:**




$$Z = \sqrt{R^2 + (1/\omega C)^2}, \quad \tan \phi = \frac{1}{\omega C R}$$

**LR circuit:**



$$Z = \sqrt{R^2 + \omega^2 L^2}, \quad \tan \phi = \frac{\omega L}{R}$$

**LCR Circuit:**




$$Z = \sqrt{R^2 + \left(\frac{1}{\omega C} - \omega L\right)^2}, \quad \tan \phi = \frac{\frac{1}{\omega C} - \omega L}{R}$$

$$\nu_{\text{resonance}} = \frac{1}{2\pi} \sqrt{\frac{1}{LC}}$$

**Power factor:**  $P = e_{\text{rms}} i_{\text{rms}} \cos \phi$

**Transformer:**  $\frac{N_1}{N_2} = \frac{e_1}{e_2}$ ,  $e_1 i_1 = e_2 i_2$



**Speed of the EM waves in vacuum:**  $c = 1/\sqrt{\mu_0 \epsilon_0}$

## 6 Modern Physics

### 6.1: Photo-electric effect


Photon's energy:  $E = h\nu = hc/\lambda$

Photon's momentum:  $p = h/\lambda = E/c$

Max. KE of ejected photo-electron:  $K_{\max} = h\nu - \phi$

Threshold freq. in photo-electric effect:  $\nu_0 = \phi/h$

Stopping potential:  $V_o = \frac{hc}{e} \left( \frac{1}{\lambda} \right) - \frac{\phi}{e}$



de Broglie wavelength:  $\lambda = h/p$

### 6.2: The Atom

Energy in  $n$ th Bohr's orbit:

$$E_n = -\frac{mZ^2e^4}{8\epsilon_0^2h^2n^2}, \quad E_n = -\frac{13.6Z^2}{n^2} \text{ eV}$$

Radius of the  $n$ th Bohr's orbit:

$$r_n = \frac{\epsilon_0 h^2 n^2}{\pi m Z e^2}, \quad r_n = \frac{n^2 a_0}{Z}, \quad a_0 = 0.529 \text{ \AA}$$

Quantization of the angular momentum:  $l = \frac{n\hbar}{2\pi}$


Photon energy in state transition:  $E_2 - E_1 = h\nu$



Wavelength of emitted radiation: for a transition from  $n$ th to  $m$ th state:

$$\frac{1}{\lambda} = RZ^2 \left[ \frac{1}{n^2} - \frac{1}{m^2} \right]$$

X-ray spectrum:  $\lambda_{\min} = \frac{hc}{eV}$



Moseley's law:  $\sqrt{\nu} = a(Z - b)$

X-ray diffraction:  $2d \sin \theta = n\lambda$

Heisenberg uncertainty principle:


$$\Delta p \Delta x \geq h/(2\pi), \quad \Delta E \Delta t \geq h/(2\pi)$$

### 6.3: The Nucleus

Nuclear radius:  $R = R_0 A^{1/3}$ ,  $R_0 \approx 1.1 \times 10^{-15} \text{ m}$

Decay rate:  $\frac{dN}{dt} = -\lambda N$

Population at time  $t$ :  $N = N_0 e^{-\lambda t}$



Half life:  $t_{1/2} = 0.693/\lambda$

Average life:  $t_{\text{av}} = 1/\lambda$

Population after  $n$  half lives:  $N = N_0/2^n$

Mass defect:  $\Delta m = [Zm_p + (A - Z)m_n] - M$


Binding energy:  $B = [Zm_p + (A - Z)m_n - M] c^2$

Q-value:  $Q = U_i - U_f$


Energy released in nuclear reaction:  $\Delta E = \Delta mc^2$   
where  $\Delta m = m_{\text{reactants}} - m_{\text{products}}$ .

### 6.4: Vacuum tubes and Semiconductors

Half Wave Rectifier:



Full Wave Rectifier:



Triode Valve:





Plate resistance of a triode:  $r_p = \left. \frac{\Delta V_p}{\Delta i_p} \right|_{\Delta V_g=0}$

Transconductance of a triode:  $g_m = \left. \frac{\Delta i_p}{\Delta V_g} \right|_{\Delta V_p=0}$

Amplification by a triode:  $\mu = - \left. \frac{\Delta V_p}{\Delta V_g} \right|_{\Delta i_p=0}$

Relation between  $r_p$ ,  $\mu$ , and  $g_m$ :  $\mu = r_p \times g_m$

Current in a transistor:  $I_e = I_b + I_c$



$\alpha$  and  $\beta$  parameters of a transistor:  $\alpha = \frac{I_c}{I_e}$ ,  $\beta = \frac{I_c}{I_b}$

Transconductance:  $g_m = \frac{\Delta I_c}{\Delta V_{be}}$

Logic Gates:

		AND	OR	NAND	NOR	XOR
A	B	AB	A+B	$\overline{AB}$	$\overline{A+B}$	$A\overline{B} + \overline{A}B$
0	0	0	0	1	1	0
0	1	0	1	1	0	1
1	0	0	1	1	0	1
1	1	1	1	0	0	0