

Resonance[®]
Educating for better tomorrow

JEE (Main) PAPER-1 (B.E./B. TECH.)

2023

COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

Date: 31 January, 2023 (SHIFT-1) | **TIME :** (9.00 a.m. to 12.00 p.m)
Duration: 3 Hours | **Max. Marks:** 300

SUBJECT: MATHEMATICS

Resonance Eduventures Ltd.
Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005
Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222
To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029
Toll Free : 1800 258 5555 | **7340010333** | [f](https://www.facebook.com/ResonanceEd) [i](https://www.instagram.com/ResonanceEd) [in](https://www.linkedin.com/company/resonance-eduventures) www.youtube.com/resonance [B](https://www.blog.resonance.ac.in) www.resonance.ac.in

This solution was download from Resonance JEE (Main) 2023 Solution portal

PART : MATHEMATICS

1. $|\vec{a}| = \sqrt{6}$, $|\vec{b}| = \sqrt{14}$ and $|\vec{a} \times \vec{b}| = \sqrt{48}$ then find $(\vec{a} \cdot \vec{b})^2$

Ans. (36)

Sol. $(\vec{a} \cdot \vec{b})^2 + (\vec{a} \times \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2$

$$(\vec{a} \cdot \vec{b})^2 + 48 = 14 \times 6$$

$$(\vec{a} \cdot \vec{b})^2 = 84 - 48 = 36$$

2. A bag contains 6 balls. 2 balls are drawn randomly if both are black then find probability that bag contains at least 5 black balls.

(1) $\left(\frac{5}{7}\right)$ (2) $\left(\frac{1}{2}\right)$ (3) $\left(\frac{1}{7}\right)$ (4) $\left(\frac{3}{7}\right)$

Ans. (1)

Sol. Total possibility = 2B + 4 others
 or
 = 3B + 3 others
 or
 = 4B + 2 others
 or
 = 5B + 1 others
 or
 = 6B + 0 others

Now

$$\text{required probability} = \frac{{}^5C_2 + {}^6C_2}{{}^6C_2 + {}^6C_2}$$

$$= \frac{{}^2C_2 + {}^3C_2 + {}^4C_2 + {}^5C_2 + {}^6C_2}{{}^6C_2 + {}^6C_2}$$

$$= \frac{10 + 15}{1 + 3 + 6 + 10 + 15} = \frac{25}{35} = \frac{5}{7}$$

3. Find remainder when 5^{99} is divided by 11

Ans. (9)

Sol. $5^{99} = 625 \times 5^{95}$
 $= 625 \times (3124 + 1)^{19}$
 $= 625 \times (11\lambda + 1)^{19}$
 Remainder = 625 or 9

4. Sum of four consecutive terms of a G.P. is 126 and their product is 1296 then find sum of common ratio of all G.P.

Ans. (3)

Sol. G.P. $\Rightarrow \frac{a}{r^3}, \frac{a}{r}, ar, ar^3$

given : $\frac{a}{r^3} \cdot \frac{a}{r} \cdot ar \cdot ar^3 = 1296 \Rightarrow a^4 = 1296 \Rightarrow a = 6$

and $\frac{a}{r^3} + \frac{a}{r} + ar + ar^3 = 126 \Rightarrow \frac{1}{r^3} + \frac{1}{r} + r + r^3 = 21$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/resonanceedu](https://www.facebook.com/resonanceedu) | twitter.com/ResonanceEdu | www.youtube.com/resonance | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 1

$$\left(r^3 + \frac{1}{r^3}\right) + \left(r + \frac{1}{r}\right) = 21$$

$$\Rightarrow \left(r + \frac{1}{r}\right)^3 - 3r \cdot \frac{1}{r} \left(r + \frac{1}{r}\right) + \left(r + \frac{1}{r}\right) = 21$$

$$t^3 - 2t - 21 = 0$$

$$(t-3)(t^2 + 3t + 7) = 0,$$

$$t = 3$$

$$r + \frac{1}{r} = 3$$

$$r^2 - 3r + 1 = 0 \begin{cases} r_1 \\ r_2 \end{cases}$$

Sum = 3

5. Find number of four digit number which are divisible by 3 or 11 and less than and equal to 2800.

Ans. (709)

Sol. Number of four digit numbers which are divisible by 3 is 600
 1002....., 2799

Number of four digit numbers which are divisible by 11 is 163

Number of four digit numbers which are divisible by 33 is 54

so required number is = 600+ 163 -54= 709

6. For ellipse $\frac{x^2}{4} + \frac{y^2}{b^2} = 1, 0 < b < 2$ maximum distance of normal at point P from its centre is 1 then e

equal to

$\sqrt{3}$

$\sqrt{3}$

$\sqrt{3}$

3

- (1) $\frac{1}{2}$ (2) $\frac{1}{3}$ (3) $\frac{1}{4}$ (4) $\frac{1}{4}$

Ans. (1)

Sol. Normal at $(2\cos \theta, b \sin \theta)$ is
 $2x \sec \theta - by \operatorname{cosec} \theta = 4 - b^2$

$$\text{Now, } 1 = \left(\frac{4 - b^2}{\sqrt{4 \sec^2 \theta + b^2 \operatorname{cosec}^2 \theta}} \right)_{\max.}$$

$$1 = \frac{4 - b^2}{2 + b}$$

$$2 + b = 4 - b^2$$

$$b^2 + b - 2 = 0$$

$$b = -2 \text{ or } 1$$

$$\text{so } b = 1$$

$$\text{Now } e = \sqrt{1 - \frac{1}{4}} = \frac{\sqrt{3}}{2}$$

7. If a_1, a_2, \dots, a_n is an A.P such that $a_5 = 2a_7$ and $a_{11} = 18$ then find

$$12 \left[\frac{1}{\sqrt{a_{10}} + \sqrt{a_{11}}} + \frac{1}{\sqrt{a_{11}} + \sqrt{a_{12}}} + \frac{1}{\sqrt{a_{12}} + \sqrt{a_{13}}} + \dots + \frac{1}{\sqrt{a_{17}} + \sqrt{a_{18}}} \right]$$

(1) 8

(2) 4

(3) 10

(4) 6

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No. : +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | facebook.com/ResonanceEdu | twitter.com/ResonanceEdu | www.youtube.com/resonance | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 2

Ans. (1)

Sol. Given $a + 4d = 2(a + 6d)$

$$a + 8d = 0 \dots\dots(1)$$

$$\text{and } a_{11} = 18 \Rightarrow a + 10d = 18 \dots\dots(2)$$

From (1) and (2)

$$-8d + 10d = 18 \Rightarrow d = 9 \text{ and } a = -72$$

$$a_{10} = a + 9d = -72 + 81 = 9$$

$$a_{18} = a + 17d = -72 + 153 = 81$$

Now

$$12 \left[\frac{1}{\sqrt{a_{10}} + \sqrt{a_{11}}} + \frac{1}{\sqrt{a_{11}} + \sqrt{a_{12}}} + \dots + \frac{1}{\sqrt{a_{17}} + \sqrt{a_{18}}} \right]$$

$$12 \left[\frac{\sqrt{a_{10}} - \sqrt{a_{11}}}{-d} + \frac{\sqrt{a_{11}} - \sqrt{a_{12}}}{-d} + \dots + \frac{\sqrt{a_{17}} - \sqrt{a_{18}}}{-d} \right]$$

$$12 \left[\frac{\sqrt{a_{10}} - \sqrt{a_{18}}}{-d} \right] = 12 \left[\frac{3 - 9}{-9} \right] = 12 \times \frac{6}{9} = 8$$

8. Five digit numbers are formed using 0, 2, 3, 4, 7, 9 (repetition is allowed) and arranged in increasing order then the rank of 42937 is

Ans. (3005)

Sol. Number starting with 2 & 3 = 2×6^4

$$\text{Number starting with 40} = 6^3$$

$$\text{Number starting with 420, 422, 423, 424}$$

$$427 = 5 \times 6^2$$

$$\text{Similarly starting with 4290 \& } 4292 = 2 \times 6$$

$$\text{Number starting with 4293 before 42937} = 6 - 1 = 5$$

$$\text{Total} = 2 \times 6^4 + 6^3 + 5 \times 6^2 + 2 \times 6 + 5 = 3005$$

9. If $f(x) + \int_3^{e^f(x)} dt = \sqrt{x+1}$; $f(3) = 2$ and $x \geq 3$ then find $12f(8)$

(1) 11

(2) 17

(3) 1

(4) 7

Ans. (2)

Sol. $f'(x) + \frac{f(x)}{x} = \frac{1}{2\sqrt{x+1}}$

$$\frac{dy}{dx} + \frac{y}{x} = \frac{1}{2\sqrt{x+1}}$$

$$\int \frac{1}{x} dx = \ln|x| + C$$

$$y \cdot x = \int x \frac{1}{2\sqrt{x+1}} dx + c \text{ put } x+1 = t^2, dx = 2t dt$$

$$y \cdot x = \frac{1}{2} \int \frac{(t^2-1)2t dt}{t} + c$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | f | t | www.youtube.com/reswatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 3

Resonance® | JEE MAIN-2023 | DATE : 31-01-2023 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

$$y \cdot x = \frac{1}{2} \left(\frac{2}{3} t^3 - 2t \right) + c$$

$$y \cdot x = \frac{(\sqrt{x+1})^3}{3} - \sqrt{x+1} + c$$

$$\text{now } f(3) = 2$$

$$6 = \frac{(2)^3}{3} - 2 + c$$

$$8 - \frac{8}{3} = c$$

$$\frac{16}{3} = c$$

$$\therefore f(x) = \frac{1}{x} \left(\frac{(\sqrt{x+1})^3}{3} - \sqrt{x+1} + \frac{16}{3} \right)$$

$$\therefore 12f(8) = \frac{12}{8} \left(\frac{27}{3} - 3 + \frac{16}{3} \right)$$

$$= \frac{3}{2} \left(6 + \frac{16}{3} \right) = 17$$

10. Find the shortest distance between lines $\frac{x-1}{3} = \frac{y-2}{0} = \frac{z-3}{-1}$ & $\frac{x-1}{1} = \frac{y-1}{1} = \frac{4-z}{1}$

(1) 3

(2) 2

(3) 1

(4) 4

Ans. (3)

Sol. $\frac{x-1}{3} = \frac{y-2}{0} = \frac{z-3}{-1}$ & $\frac{x-1}{1} = \frac{y-1}{1} = \frac{4-z}{1}$

$$\text{S.D.} = \left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b} \times \vec{d})}{(\vec{b} \times \vec{d})} \right|$$

$$\vec{b} \times \vec{d} = \begin{vmatrix} i & j & k \\ 3 & 0 & -1 \\ 1 & 1 & -1 \end{vmatrix} = i(1) - j(-2) + k(3)$$

$$\text{S.D.} = \left| \begin{vmatrix} 0 & 1 & -1 \\ 3 & 0 & -1 \\ 1 & 1 & -1 \end{vmatrix} \right|$$

$$= |0(1) - 1(-2) - 1(3)|$$

$$= 1$$

11. A Circle $x^2 + y^2 - 4x - 6y + 11 = 0$ on the tangent to it at (3, 2) is rolled up by 4 units upwards. If M and

N are foot of perpendiculars of centres on x-axis, then area of quadrilateral by centres, M and N is

(1) 20

(2) 10

(3) 30

(4) 40

Ans (1)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222


To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | f | t | www.youtube.com/reswatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 4

Sol. tangent; $x - y - 1 = 0$
 since $\angle C_1 C_2 D = 45^\circ$
 $C_2 D = C_1 D = 4$



$$\text{Area} = \frac{1}{2} (3 + 7) \times 4 = 20 \text{ units.}$$

12. A wire of ℓ length is divided into 2 parts of length ℓ_1 and ℓ_2 . From wire of length ℓ_1 a square is formed with area 'a₁' and with wire of length ℓ_2 a circle is formed with area a₂. If $2a_1 + 3a_2$ is minimum then find

$$\frac{\pi \ell_1}{\ell_2}$$

Ans. (6)

Sol. $\ell = 4x + 2\pi r$, where $x =$ side of square and $r =$ radius of circle

now let $2a_1 + 3a_2 = A$

$$A = 2x^2 + 3\pi r^2$$

$$= 2x^2 + 3\pi \left(\frac{\ell - 4x}{2\pi} \right)^2 = 2x^2 + \frac{3}{4\pi} (\ell - 4x)^2$$

$$\frac{dA}{dx} = 4x - \frac{2 \times 3 \times 4}{4\pi} (\ell - 4x)$$

for maxima minima

$$\frac{dA}{dx} = 0$$

$$\therefore x = \frac{2 \times 3}{4\pi} (\ell - 4\pi)$$

$$4\pi x = 6\ell - 24\pi$$

$$x = \frac{6\ell}{4\pi + 24}$$

$$\text{now } 4x = \ell_1 = \frac{6\ell}{\pi + 6} \quad \text{also } \ell_2 = 2\pi r$$

$$= \ell - 4x = \ell - \frac{6\ell}{\pi + 6} = \frac{\pi\ell + 6\ell - 6\ell}{\pi + 6} = \frac{\pi\ell}{\pi + 6}$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | f | t | www.youtube.com/reswatch | b | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 5

$$\text{now } \frac{\pi \ell_1}{\ell_2} = \frac{\pi \times \frac{6\ell}{\pi + 6}}{\frac{\pi \ell}{\pi + 6}} = 6$$

13. Let $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & -1 \\ 0 & 12 & -3 \end{bmatrix}$ then sum of diagonal element of $(A + I)^{11}$ is

(1) 4097 (2) 4095 (3) 6465 (4) 2049

Ans. (1)

Sol. $A^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & -1 \\ 0 & 12 & -3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & -1 \\ 0 & 12 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & -1 \\ 0 & 12 & -3 \end{bmatrix} = A$

Now $(A + I)^{11} = {}^{11}C_0 A^{11} + {}^{11}C_1 A^{10} + \dots + {}^{11}C_{11} I$
 $= A ({}^{11}C_0 + {}^{11}C_1 + \dots + {}^{11}C_{10}) + I$
 $= A (2^{11} - 1) + I$
 Trace of $(A + I)^{11}$
 $= 2^{11} + 4(2^{11} - 1) + 1 + (-3)(2^{11} - 1) + 1$
 $= 2 \times 2^{11} + 1$
 $= 2^{12} + 1$

14. If $\sin^{-1}(\alpha/17) + \cos^{-1}(4/5) - \tan^{-1}(\frac{77}{36}) = 0$ then find $\sin^{-1}(\sin \alpha) + \cos^{-1}(\cos \alpha)$

- (1) π (2) 2π (3) 3π (4) 4π

Ans. (1)

Sol. $\sin^{-1} \frac{\alpha}{17} = \tan^{-1} \frac{77}{36} - \cos^{-1} \frac{4}{5}$
 $\sin^{-1} \frac{\alpha}{17} = \tan^{-1} \frac{77}{36} - \sin^{-1} \frac{3}{5}$
 $\frac{\alpha}{17} = \sin \left(\tan^{-1} \frac{77}{36} \right) \cos \sin^{-1} \left(\frac{3}{5} \right) - \cos \tan^{-1} \left(\frac{77}{36} \right) \sin \sin^{-1} \left(\frac{3}{5} \right)$
 $\frac{\alpha}{17} = \frac{77}{85} \times \frac{4}{5} - \frac{36}{85} \times \frac{3}{5}$
 $= \frac{308 - 108}{425} = \frac{200}{425} = \frac{8}{17}$
 $\alpha = 8$
 Now $\sin^{-1} \sin \alpha + \cos^{-1} \cos \alpha$
 $= \sin^{-1} \sin 8 + \cos^{-1} \cos 8$
 $= -8 + 3\pi + 8 - 2\pi$
 $= \pi$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/resonanceedu](https://www.facebook.com/resonanceedu) | twitter.com/resonanceedu | www.youtube.com/reswatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 6

15. If $B = \ln(1-x)$ and $P(x) = \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \dots + \frac{x^{50}}{50} \right)$, then $\int_0^x \frac{1}{1-t} dt$ equals

- (1) $-(B + P(x))$ (2) $-B + P(x)$
 (3) $B - P(x)$ (4) $B + P(x)$

Ans. (3)

Sol. $\int_0^x \frac{1}{1-t} dt$
 $= \int_0^x (1 + t + t^2 + \dots + \infty) dt = \left(\frac{t^2}{2} + \frac{t^3}{3} + \dots \right)_0^x = \frac{x^2}{2} + \frac{x^3}{3} + \dots$
 $\ln(1-x) - P(x) = B - P(x)$

16. If $\sqrt{x^2 - 4x + 3} + \sqrt{x^2 - 9} = \sqrt{4x^2 - 14x + 6}$ then find real roots.

Ans. (3)

Sol. $\sqrt{(x-1)(x-3)} + \sqrt{(x-3)(x+3)} = \sqrt{2(2x-1)(x-3)}$
 $x-3 = 0$ and $\sqrt{x-1} + \sqrt{x+3} = \sqrt{2(2x-1)}$ (1)
 Squaring both side ; $x-1 + x+3 + 2\sqrt{(x-1)(x+3)} = 4x-2$

$$2\sqrt{(x-1)(x+3)} = 2x - 4$$

$$\sqrt{x^2 + 3x - x - 3} = x - 2 \dots (2)$$

$$\text{Squaring both side ; } x^2 + 2x - 3 = x^2 - 4x + 4$$

$$6x = 7 \Rightarrow x = \frac{7}{6}$$

At $x = \frac{7}{6}$ equation (2) positive = negative

Rejected

\therefore real root $x = 3$

17. Curve $C_1 : |z| = 4$ and C_2 : the locus of $z + \frac{1}{z}$ (where z satisfy the curve c_1) then C_1 & C_2

(1) Intersect at 2 points

(2) Intersect at 4 points

(3) do not intersect

(4) None of these

Ans. (2)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | www.resonance.ac.in | www.facebook.com/resonance | www.youtube.com/resonance | www.instagram.com/resonance | www.linkedin.com/resonance

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 7

Resonance® | JEE MAIN-2023 | DATE : 31-01-2023 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

Sol. $|z| = 4$ is a circle and $z + \frac{1}{z}$ is an ellipse $\frac{x^2}{\left(\frac{17}{4}\right)^2} + \frac{y^2}{\left(\frac{15}{4}\right)^2} = 1$



18.
$$\int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \frac{2+3\sin x}{\sin x(1+\cos x)} dx$$

(1) $\frac{10}{3} + \sqrt{3} - \frac{1}{2} \log 3$

(2) $\frac{20}{3} + \sqrt{3} - \log 3$

(3) $\frac{10}{3} - \sqrt{3} + \frac{1}{2} \log 3$

(4) $\frac{20}{3} - \sqrt{3} + \log 3$

Ans. (3)

Sol.
$$= \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \frac{2(1-\cos x)}{\sin x(1-\cos^2 x)} dx + \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \frac{3}{1+\cos x} dx$$

$$= \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \frac{2}{\sin^3 x} dx - 2 \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \cot x \cdot \cos^2 x dx + 3 \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \sec^2 \frac{x}{2} dx$$

$$= 2 \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \sqrt{1+\cot^2 x} \cdot \cos^2 x dx - 2 \int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \cot x \cdot \cos^2 x dx + 3 \left[\tan \frac{x}{2} \right]_{\frac{\pi}{3}}^{\frac{2\pi}{3}}$$

Let $\cot x = t$

$$= -2 \int_{\frac{1}{\sqrt{3}}}^0 \sqrt{1+t^2} dt + 2 \int_{\frac{1}{\sqrt{3}}}^0 t dt + 3 \left[1 - \frac{1}{\sqrt{3}} \right]$$

$$= 2 \left[\frac{1}{2} \sqrt{1+t^2} + \frac{1}{2} \log |t + \sqrt{1+t^2}| \right]_{\frac{1}{\sqrt{3}}}^0 - \left[\frac{t^2}{2} \right]_{\frac{1}{\sqrt{3}}}^0 + 3 - \sqrt{3}$$

$$= \frac{2}{3} + \log \sqrt{3} - \frac{1}{3} + 3 - \sqrt{3} = \frac{10}{3} - \sqrt{3} + \frac{1}{2} \log 3$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/resonanceedu](https://www.facebook.com/resonanceedu) | twitter.com/resonanceedu | www.youtube.com/resonance | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 8



JEE MAIN-2023 | DATE : 31-01-2023 (SHIFT-1) | PAPER-1 | MEMORY BASED | MATHEMATICS

19. The direction ratio of two lines which are parallel are given by $\langle 2, 1, -1 \rangle$ and $\langle \alpha + \beta, 1 + \beta, 2 \rangle$. Then the value of $|2\alpha + 3\beta|$ is _____.

Ans. (11)

Sol. $\frac{\alpha + \beta}{2} = \frac{1 + \beta}{1} = \frac{2}{-1}$

$\alpha = -1, \quad \beta = -3$

$|2\alpha + 3\beta| = 11$

20. Domain of the function $y = \frac{[x]}{1+x^2}$ is $[2, 6)$ then Range of y is

(1) $\left(\frac{5}{37}, \frac{2}{5}\right]$

(2) $\left(\frac{5}{26}, \frac{2}{5}\right]$

(3) $\left(\frac{4}{26}, \frac{2}{5}\right]$

(4) $\left(\frac{5}{26}, \frac{2}{5}\right]$

Ans. (1)

Sol. $y = \frac{[x]}{1+x^2} = \begin{cases} \frac{2}{1+x^2} & ; 2 \leq x < 3 \\ \frac{3}{1+x^2} & ; 3 \leq x < 4 \\ \frac{4}{1+x^2} & ; 4 \leq x < 5 \\ \frac{5}{1+x^2} & ; 5 \leq x < 6 \end{cases}$

Since, $y = \frac{[x]}{1+x^2}$ is decreasing.

for range, $f(2) = \frac{2}{5} = 0.4$

$f(3) = \frac{3}{10} = 0.3$

$f(4) = \frac{4}{17} = 0.23$

$f(5) = \frac{5}{26} = 0.19$

$f(6) = \frac{5}{37} = 0.13$

So range of $y \in \left(\frac{5}{37}, \frac{2}{5}\right]$

21. Let $\vec{a} = 2\hat{i} + \hat{j} - \hat{k}$ and \vec{b}, \vec{c} are non-zero vectors such that $\vec{b} \cdot \vec{c} = 0$ and $|\vec{a} + \vec{b} + \vec{c}| = |\vec{a} + \vec{b} - \vec{c}|$

Statement -1 : $|\vec{a} + \lambda \vec{c}| \geq |\vec{a}|$ for $\lambda \in \mathbb{R}$

Statement -2 : \vec{a} & \vec{c} are parallel, then

(1) Both correct

(2) Both wrong

(3) Only S-1 correct

(4) Only S₂ correct

Ans. (3)

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/resonanceedu](https://www.facebook.com/resonanceedu) | twitter.com/resonanceedu | www.youtube.com/resonance | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 9

Sol. $|\vec{a} + \vec{b} + \vec{c}|^2 = |\vec{a} + \vec{b} - \vec{c}|^2$

$$4\vec{b} \cdot \vec{c} + 4\vec{a} \cdot \vec{c} = 0 \Rightarrow \vec{a} \cdot \vec{c} = 0$$

Statement -1 $|\vec{a} + \lambda \vec{c}|^2 \geq |\vec{a}|^2$

$$a^2 + \lambda^2 c^2 + 2\vec{a} \cdot \vec{c} \lambda \geq a^2$$

$$\lambda^2 c^2 \geq 0 \text{ which is true}$$

Statement -1 is true

Statement -2 \vec{a} & \vec{c} are perpendicular. So, statement 2 is false

22. $y = f(x)$ is a parabola with focus $(-\frac{1}{2}, 0)$ and directrix $y = -\frac{1}{2}$.

Given that $\tan^{-1} \sqrt{f(x)} + \sin^{-1} \sqrt{f(x)+1} = \frac{\pi}{2}$. Then number of solution for x is

Ans. (2)

Sol. Equation of parabola $\sqrt{(x + \frac{1}{2})^2 + y^2} = |y + \frac{1}{2}|$

$$x^2 + \frac{1}{4} + x + y^2 = y^2 + \frac{1}{4} + y$$

$$x^2 + x = y$$

$$(x + \frac{1}{2})^2 = (y + \frac{1}{4})$$

$$Y = f(x) = (x + \frac{1}{2})^2 - \frac{1}{4}$$

Now $\tan^{-1} \sqrt{f(x)} + \sin^{-1} \sqrt{f(x)+1} = \frac{\pi}{2}$

$$\cos^{-1} \frac{1}{\sqrt{1+f(x)}} + \sin^{-1} \sqrt{f(x)+1} = \frac{\pi}{2}$$

So $\sqrt{f(x)+1} = \frac{1}{\sqrt{1+f(x)}}$

$$f(x) + 1 = 1$$

$$f(x) = 0$$

$$(x + \frac{1}{2})^2 = \pm \frac{1}{2}$$

$$x = 0, -1$$

Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | facebook.com/Resonanceedu | twitter.com/Resonanceedu | www.youtube.com/reswater | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 10



TO KNOW MORE

Call: 0744-2777777,
8441024095

TARGET: JEE (ADV.) 2023

Polish your subject knowledge with the guidance of
Top Notch Sr. Faculty of Resonance

SPARK

15 WEEKS COMPAC COURSE

OFFLINE / ONLINE

CLASS
STARTS

6th FEBRUARY
2023

ACADEMIC FEATURES

- Course Duration: **15 Weeks**.
- Total No. of Lectures: **234** (P: 78 | C: 78 | M: 78)
- Duration of One Lecture: **1.5 hrs.** (90 Minutes)
- Classroom Teaching Hours: **351 Hrs.**
- Testing Duration: **60 Hrs.**
- Total Academic Hours: **411 Hrs.**

Course Features

- Study Material
- Quick Classes
- Back up support of recorded lectures
- Port/ Full Syllabus Test Series

Facilities for Offline Students

- IT-Enabled Computer Lab
- Self Study Rooms for Boys & Girls

AIR **6**

SAHARSH VAHNEYA

JEE (Adv.) 2022

TARGET: JEE (Main) 2023

Boost your Percentile with

PERCENTILE BOOSTER COURSE

8 WEEKS COMPAC COURSE

OFFLINE / ONLINE

CLASS
STARTS

6th FEBRUARY
2023

COURSE FEATURES

- Complete Course Coverage
- 25 Chapter wise Test
- Regular Practice through 35 Daily Online Practice Test
- 5 Full Syllabus Test
- 3 Joint Preparatory Test
- Approx 2000 practice Ques.
- 113 Teaching hours
- 99 Testing Hours
- Regular Test discussion classes for concept clearance
- Back up support of recorded lectures

AIR **37**

VEDAN VEJALA

JEE (Main) 2022

Resonance[®]
Educating for better tomorrow

JEE (ADVANCED) 2022
RESULT

RESONites ने फिर लहराया सफलता का परचम

STUDENTS FROM CLASSROOM PROGRAM (OFFLINE/ ONLINE)

AIR

6

KARTHIKEYA
POLISETTY

Roll No.: 21925115

AIR-1
GEN-EWS

AIR

8

DHEERAJ
KURUKUNDA

Roll No.: 21920714

Students
in TOP-100
All India
Ranks
(AIRs)



AIR-11
DEVANSHU MALI
Roll No.: 21219044



AIR-15
ARHIF ANAND
Roll No.: 21929195



AIR-35
SANSKAR SHRIVASTAVA
Roll No.: 21929155



AIR-50
ANURAG GARG
Roll No.: 21220122



AIR-54
SOUMITRA O. NAVIK
Roll No.: 21220984



AIR-58
KARISHK SHARMA
Roll No.: 21230454

ADMISSIONS OPEN FOR ACADEMIC SESSION 2023-24

TARGET: JEE (Adv.) 2024

for Class XII Passed Student



VISHESH COURSE

MODE: OFFLINE / ONLINE

CLASS STARTS
10th & 17th April

TARGET: JEE (Main) 2024

for Class XII Passed Student



ABHYAAS COURSE

MODE: OFFLINE / ONLINE

CLASS STARTS
10th & 24th April

SCHOLARSHIP ON THE BASIS OF JEE (MAIN) 2023 %ILE / AIR

Resonance Eduventures Limited

REGISTERED & CORPORATE OFFICE: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Rajasthan) - 324005
Tel. No.: 0744-2777777, 2777700 | CIN: U80302RJ2007PLC024029

Social Media Connect

