

**Resonance®**  
Educating for better tomorrow

# JEE (Main) PAPER-1 (B.E./B. TECH.)

## 2023

### COMPUTER BASED TEST (CBT) Memory Based Questions & Solutions

Date: 01 February, 2023 (SHIFT-2) | TIME : (3.00 p.m. to 6.00 p.m)  
Duration: 3 Hours | Max. Marks: 300

**SUBJECT: MATHEMATICS**

**Resonance Eduventures Ltd.**  
Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005  
Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222  
To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029  
Toll Free : 1800 258 5555 | 7340010333 | facebook.com/ResonanceEdu | twitter.com/ResonanceEdu | www.youtube.com/resowatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

**PART : MATHEMATICS**

1. Given functional relation be  $f(x) + f\left(\frac{1}{1-x}\right) = 1 + x, \forall x \in \mathbb{R} - \{0,1\}$ , then the value of  $f(2)$  is.

- (1)  $\frac{9}{4}$                       (2)  $\frac{7}{4}$                       (3) 0                      (4) None of these

Ans. (1)  
Sol. ,  $x = 2 \Rightarrow f(2) + f(-1) = 3$  .....(1)  
..... 1 ..... (1) ..... 3

$$x = \frac{1}{2} \Rightarrow 1\left(\frac{1}{2}\right) + 1(1) = \frac{3}{2} \quad \dots\dots\dots(1)$$

$$x = -1 \Rightarrow f(-1) + f\left(\frac{1}{2}\right) = 0 \quad \dots\dots\dots(3)$$

$$(2) \dots\dots (3) \Rightarrow f(2) - f(-1) = \frac{3}{2} \quad \dots\dots\dots(4)$$

$$(1) + (4) \Rightarrow 2f(2) = \frac{9}{2} \Rightarrow f(2) = \frac{9}{4}$$

2. If  $y = x^x$  then value of  $y''(2) - 2y'(2)$  is  
 (1)  $2 - 4(\ln 2)^2$       (2)  $4(\ln 2)^2 - 2$       (3)  $(\ln 2)^2 - 4$       (4)  $4(\ln 2)^2 + 2$

Ans. (2)

Sol.  $y = x^x$

$$\ln y = x \ln x$$

$$\frac{1}{y} \cdot y' = 1 + \ln x$$

$$y' = y(1 + \ln x) \quad \dots\dots\dots(1)$$

$$y' = x^x(1 + \ln x)$$

at  $x = 2$  we have  $y = 4$

$$\text{So } y'(2) = 4(1 + \ln 2) \quad \dots\dots\dots(2)$$

$$\text{and } y'' = y'(1 + \ln x) + \frac{y}{x}$$

$$y''(2) = y'(1 + \ln 2) + 2$$

$$y''(2) - y'(2) = y'(\ln 2) + 2$$

$$\begin{aligned} y''(2) - 2y'(2) &= (\ln 2 - 1)y'(2) + 2 \\ &= 4(\ln 2 - 1)(\ln 2 + 1) + 2 \\ &= 4(\ln 2)^2 - 2 \end{aligned}$$

3. If the term independent of  $x$  (coefficient of  $x^0$ ) in the expansion of  $\left(x^{\frac{2}{3}} + \frac{\alpha}{x^3}\right)^{22}$  is 7315 then the value of

$|\alpha|$  is-

- (1) 2      (2) 1      (3) 6      (4) 5

Ans. (2)

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No. : +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555    7340010333    f contact@resonance.edu    t twitter.com/ResonanceEdu    y www.youtube.com/resonance    b blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 1

Sol.  $T_{r+1} = {}^{22}C_r \left(\frac{\alpha}{x^3}\right)^r \left(x^{\frac{2}{3}}\right)^{22-r}$

$$= {}^{22}C_r \cdot \alpha^r \cdot x^{\frac{22-r}{3}}$$

for  $x^0 \Rightarrow r = 4$

$$\text{coefficient of } x^0 = {}^{22}C_4 \cdot \alpha^4 = 7315$$

$$\Rightarrow \frac{22 \cdot 21 \cdot 20 \cdot 19}{4 \cdot 3 \cdot 2} \alpha^4 = 7315$$

$$\Rightarrow (11 \times 7 \times 5 \times 19) \alpha^4 = 7315$$

$$\Rightarrow \alpha^4 = 1$$

$$\Rightarrow |\alpha| = 1$$

4. The number of integral solution of equation  $x + y + z = 21$  where  $x \geq 1, y \geq 3, z \geq 5$   
 (1)  ${}^{14}C_7$       (2)  ${}^{14}C_3$       (3)  ${}^{14}C_2$       (4) None of these

Ans. (3)

Sol.  $x \geq 1, y \geq 3, z \geq 5 \Rightarrow x - 1 \geq 0, y - 3 \geq 0, z - 5 \geq 0$

$$\text{Let } x - 1 = X, y - 3 = Y, z - 5 = Z$$

$$\text{So } x + y + z = 21$$

$$X + 1 + Y + 3 + Z + 5 = 21$$

$$X + Y + Z = 12 \quad X \geq 0, Y \geq 0, Z \geq 0$$

$$\text{No. of integral solution} = {}^{12+3-1}C_2 = {}^{14}C_2$$

5. The value of  $\sum_{n=1}^{\infty} \frac{(2n^2 + 2n + 4)}{(n+1)}$  is

(1)  $3e + \frac{3}{2e} - 4$       (2)  $e + \frac{3}{2e} - 4$       (3)  $3e + \frac{3}{e} + 4$       (4)  $e + \frac{1}{2e} - 4$

Ans. (1)

Sol. 
$$\sum_{n=1}^{\infty} \frac{2n^2 + 2n + 4}{2n!}$$

$$= \frac{1}{2} \sum_{n=1}^{\infty} \frac{2n(2n-1) + 6n + 8}{2n!}$$

$$= \frac{1}{2} \sum_{n=1}^{\infty} \left\{ \frac{1}{(2n-2)!} + \frac{3}{(2n-1)!} + \frac{8}{2n!} \right\}$$

$$= \frac{1}{2} \left\{ \left( \frac{e + e^{-1}}{2} \right) + 3 \left( \frac{e - e^{-1}}{2} \right) + 8 \left( \frac{e + e^{-1}}{2} - 1 \right) \right\}$$

$$= \frac{1}{2} (6e + 3e^{-1} - 8)$$

$$= 3e + \frac{3}{2e} - 4$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029  
Toll Free : 1800 258 5555 | 7340010333 | facebook.com/ResonanceEdu | twitter.com/ResonanceEdu | www.youtube.com/reswatch | blog.resonance.ac.in

This solution was downloaded from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 2

6. If a matrix  $A = \frac{1}{2} \begin{bmatrix} 1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{bmatrix}$  then

(1)  $A^{30} - A^{25} = I - A$       (2)  $A^{30} = A^{25}$       (3)  $A^{30} - A^{25} = A - I$       (4)  $A^{30} - A^{25} = 2I - A$

Ans. (1)

Sol.  $A = \begin{bmatrix} \frac{1}{2} & \frac{-\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix} = \begin{bmatrix} \cos 60^\circ & -\sin 60^\circ \\ \sin 60^\circ & \cos 60^\circ \end{bmatrix} \dots\dots\dots(1)$

$\therefore A(\alpha) \cdot A(\beta) = A(\alpha + \beta)$   
So by above.

$$A^{30} \left( \frac{\pi}{3} \right) = \begin{bmatrix} \cos 30 \left( \frac{\pi}{3} \right) & -\sin 30 \left( \frac{\pi}{3} \right) \\ \sin 30 \left( \frac{\pi}{3} \right) & \cos 30 \left( \frac{\pi}{3} \right) \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

$$A^{25} \left( \frac{\pi}{3} \right) = \begin{bmatrix} \cos 25 \left( \frac{\pi}{3} \right) & -\sin 25 \left( \frac{\pi}{3} \right) \\ \sin 25 \left( \frac{\pi}{3} \right) & \cos 25 \left( \frac{\pi}{3} \right) \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & \frac{-\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix} = A$$

$$A^{30} - A^{25} = I - A$$

7. The solution set of equation  $2 \tan^{-1} \left( \frac{1-x}{1+x} \right) = \cos^{-1} \left( \frac{1-x^2}{1+x^2} \right)$  for  $0 < x < 1$  is 'S' then

- (1) Number of elements in S which are smaller than  $\frac{1}{2}$  is 1
- (2) Number of elements in S which are greater than  $\frac{1}{2}$  is 1
- (3) Number of elements in S which are smaller than  $-\frac{5}{2}$  is 1
- (4) Number of elements in S which are greater than 1 is 1

Ans. (1)

Sol. Put  $x = \tan \theta$        $\theta \in \left( 0, \frac{\pi}{4} \right)$

$$2 \tan^{-1} \left( \frac{1 - \tan \theta}{1 + \tan \theta} \right) = \cos^{-1} \left( \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right)$$

$$2 \tan^{-1} \left[ \tan \left( \frac{\pi}{4} - \theta \right) \right] = \cos^{-1} [\cos(2\theta)]$$

$$\Rightarrow 2 \left( \frac{\pi}{4} - \theta \right) = 2\theta \Rightarrow \theta = \frac{\pi}{8}$$

$$\Rightarrow x = \tan \frac{\pi}{8}$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | f | t | www.youtube.com/reswatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 3


| JEE MAIN-2023 | DATE : 01-02-2023 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

8. The sum of absolute maximum and absolute minimum value of function  $f(x) = |x^2 - 5x + 6| - 3x + 2$  in interval  $[-1, 3]$  is

Ans. (24)

$$\text{Sol. } f(x) = \begin{cases} x^2 - 5x + 6 - 3x + 2 & , x \in (-\infty, 2) \cup [3, \infty) \\ -(x^2 - 5x + 6) - 3x + 2 & , x \in [2, 3] \end{cases}$$

$$\Rightarrow f(x) = \begin{cases} x^2 - 8x + 8 & , x \in (-\infty, 2) \cup [3, \infty) \\ -x^2 + 2x - 4 & , x \in [2, 3] \end{cases}$$



$$\text{absolute maximum} = |f(-1)| = |(-1)^2 - 8(-1) + 8| = 17$$

$$\text{absolute minimum} = |f(3)| = 7$$

$$\text{sum} = 17 + 7 = 24$$

9. Three different Arithmetic progression have first term 2, 3, 7, and common differences 3, 4, 5 respectively if the last terms of sequences are 359, 239 and 197 respectively then the sum of all common terms of the three progression is.

Ans. (321)

$$\text{Sol. } S_1 = \{2, 5, 8, 11, 14, \dots, 359\}$$

$$S_2 = \{3, 7, 11, 15, \dots, 239\}$$

$$S_3 = \{7, 12, 17, 22, \dots, 197\}$$

common AP

$$S = \{47, 107, 167\}$$

$$\text{Hence sum of common AP} = 321$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | f | t | www.youtube.com/reswatch | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 4

10.  $9 = x_1 < x_2 < x_3 < \dots < x_7$  are in arithmetic progression with common difference  $d$ . If the standard deviation of the seven terms is 4 and  $\bar{x}$  is the mean then the value of  $\bar{x} + x_6$  is

Ans. (34)

Sol. Mean  $\Rightarrow \bar{x} = \frac{\sum_{i=1}^7 x_i}{7} = \frac{7[2a+6d]}{7} = a+3d = x_4$

Variance =  $\frac{\sum_{i=1}^7 (x_i - \bar{x})^2}{7} = (4)^2 \Rightarrow \frac{\sum_{i=1}^7 (x_i - x_4)^2}{7} = 16$

$\Rightarrow \frac{(3d)^2 + (2d)^2 + d^2 + 0 + d^2 + (2d)^2 + (3d)^2}{7} = 16$

$= 4d^2 = 16 \Rightarrow d = 2$

$\Rightarrow \bar{x} = 9 + 3(2) = 15$

$\& x_6 = a + 5d = 9 + 5(2) = 19 \Rightarrow \bar{x} + x_6 = 34$

11. Which of the following is tautology

- (1)  $p \rightarrow (\sim p \wedge q)$       (2)  $p \rightarrow (p \vee q)$       (3)  $p \rightarrow (\sim p \vee q)$       (4)  $p \rightarrow (\sim p \wedge \sim q)$

Ans. (2)

Sol. A)  $p \rightarrow (\sim p \wedge q) \equiv \sim p \vee (\sim p \wedge q) \neq T$

B)  $p \rightarrow (p \vee q) \equiv \sim p \vee (p \vee q) \equiv (\sim p \vee p) \vee q$   
 $\equiv T \vee q$   
 $\equiv T \vee q$

C)  $p \rightarrow (\sim p \vee q) \equiv \sim p \vee (\sim p \vee q) \neq T$  (for  $p = T, q = F$ )


D)  $p \rightarrow (\sim p \wedge q) \equiv \sim p \vee (\sim p \wedge q) \neq T$  (for  $p = T$ )

12. The area bounded by curves  $xy < 8, y < x^2$  and  $y > 1$  is

- (1)  $4\ln 2 - \frac{14}{3}$       (2)  $4\ln 2 + \frac{20}{3}$       (3)  $8\ln 4 - \frac{14}{3}$       (4)  $8\ln 4 - \frac{20}{3}$

Ans. (3)

Sol.  $A = \int_1^4 \left( \frac{8}{y} - \sqrt{y} \right) dy$



$= \left[ 8\ln y - \frac{y^{3/2}}{3/2} \right]_1^4 = 8(\ln 4 - \ln 1) - \frac{2}{3}(4^{3/2} - 1)$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | f | t | www.youtube.com/reswatch | b | blog.resonance.ac.in

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 5

$= 8\ln 4 - \frac{16}{3} + \frac{2}{3} = 8\ln 4 - \frac{14}{3}$  Ans. C

13. The value of  $\int_{-\pi/4}^{\pi/4} \frac{x + \frac{\pi}{4}}{2 - \cos 2x} dx$  is

- (1)  $\frac{\pi^2}{6\sqrt{3}}$       (2)  $\frac{\pi^2}{12}$       (3)  $\frac{\pi^2}{3\sqrt{3}}$       (4)  $\frac{\pi^2}{12\sqrt{3}}$

Ans. (1)



$$\Rightarrow x_0 = \pm 3\sqrt{2}$$

$$\text{Point} \rightarrow \left(3\sqrt{2}, -\frac{3}{\sqrt{2}}\right) \text{ or } \left(-3\sqrt{2}, \frac{3}{\sqrt{2}}\right)$$

$$\Rightarrow \sqrt{2}|x_0 - y_0| = \sqrt{2}\left|3\sqrt{2} - \left(-\frac{3}{\sqrt{2}}\right)\right| = 9$$

16. How many six digit numbers are formed by digits 4, 5 and 9 which are divisible by 6 is.

Ans. (81)

Sol. Unit digit must be 4 since number should be divisible by 2.

Four out of remaining five places, each has 3 options and remaining one place will have only one option

so total number of six digit numbers =  $3 \cdot 3 \cdot 3 \cdot 3 \cdot 1 = 81$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [www.facebook.com/resonanceedu](https://www.facebook.com/resonanceedu) | [www.instagram.com/resonanceedu](https://www.instagram.com/resonanceedu) | [www.youtube.com/resonanceedu](https://www.youtube.com/resonanceedu) | [blog.resonance.ac.in](https://www.resonance.ac.in/blog)

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 7

Resonance® | JEE MAIN-2023 | DATE : 01-02-2023 (SHIFT-2) | PAPER-1 | MEMORY BASED | MATHEMATICS

17. A vector ( $\vec{r}$ ) satisfy  $\vec{r} \times \vec{a} = \vec{c} \times \vec{a}$  and  $\vec{r} \cdot \vec{b} = 0$

Where  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$

$\vec{b} = 2\hat{i} - \hat{j} + \hat{k}$

$\vec{c} = 3\hat{i} + \hat{k}$

then value of  $|\vec{r}|$  is

(1)  $\frac{5}{2}$

(2)  $\frac{5}{\sqrt{3}}$

(3)  $\frac{3\sqrt{3}}{2}$

(4)  $\frac{5\sqrt{3}}{2}$

Ans. (4)

Sol.  $\vec{r} \times \vec{a} = \vec{c} \times \vec{a}$

$$\Rightarrow (\vec{r} - \vec{c}) \times \vec{a} = 0$$

$$\Rightarrow (\vec{r} - \vec{c}) = \lambda \vec{a}$$

$$\Rightarrow \vec{r} = \vec{c} + \lambda \vec{a}$$

$$\Rightarrow \vec{r} \cdot \vec{b} = \vec{c} \cdot \vec{b} + \lambda \vec{a} \cdot \vec{b}$$

$$\Rightarrow 0 = (6+1) + \lambda(2-1+1)$$

$$\Rightarrow \lambda = -\frac{7}{2}$$

$$\text{Hence } \vec{r} = \vec{c} - \frac{7}{2} \vec{a}$$

$$= (3\hat{i} + \hat{k}) - \frac{7}{2}(\hat{i} + \hat{j} + \hat{k})$$

$$= -\frac{1}{2}\hat{i} - \frac{7}{2}\hat{j} - \frac{5}{2}\hat{k}$$

$$\Rightarrow |\vec{r}| = \sqrt{\frac{1}{4} + \frac{49}{4} + \frac{25}{4}} = \frac{5\sqrt{3}}{2}$$

18. Two unbiased dice are thrown simultaneously. A is the event such that the number on the first die is less than second die. B is the event such that number on the first die is even and number on the second die is odd. C is the event such that first die shows odd number and second die shows even number then

(1)  $n((A \cup B) \cap C) = 6$

(2) A and B are mutually exclusive events

(3) A & B independent event

(4)  $n(A) = 18, n(B) = 6, n(C) = 6$

Ans. (1)

Sol. A (I < II)      B (EO)      C(OE)

$n(A) = 15$        $n(B) = 9$        $n(C) = 9$

$n(A \cap B) = 6$        $n(A \cap C) = 6$        $n(B \cap C) = 6$

$$\begin{aligned}n(A \cap B) &= 5 & n(A \cup B) &= 9 & n(B \cap C) &= 0 \\n(A \cap B \cap C) &= 0 \\n((A \cup B) \cap C) &= n(A \cap C) + n(B \cap C) - n(A \cap B \cap C) = 6\end{aligned}$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/resonanceedu](https://www.facebook.com/resonanceedu) | [twitter.com/resonanceedu](https://twitter.com/resonanceedu) | [www.youtube.com/resonance](https://www.youtube.com/resonance) | [blog.resonance.ac.in](https://www.blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 8

19. If  $\frac{dy}{dx} = \frac{x^2 + 3y^2}{3x^2 + y^2}$ , satisfies  $y(1) = 0$ , then

$$(1) \frac{2x^2}{(x-y)^2} = n|x-y| + \frac{2x}{x-y}$$

$$(2) \frac{2x}{(x-y)^2} = n|x-y| + 1$$

$$(3) \frac{2x^2}{(x-y)^2} = n|x-y| + \frac{y}{x-y}$$

$$(4) \frac{2x}{(x-y)^2} = n|x-y| + \frac{y}{x-y}$$

Ans. (1)

Sol.  $\frac{dy}{dx} = \frac{x^2 + 3y^2}{3x^2 + y^2}$  is a homogeneous equation

$$\text{Put } y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$$

$$\Rightarrow v + x \frac{dv}{dx} = \frac{1 + 3v^2}{3 + v^2}$$

$$\Rightarrow \frac{3 + v^2}{1 + 3v^2 - 3v - v^3} dv = \frac{dx}{x}$$

$$\Rightarrow \int \frac{3 + v^2}{-v^3 + 3v^2 - 3v + 1} dv = \int \frac{dx}{x}$$

$$\Rightarrow \int \frac{3 + v^2}{(1-v)^3} dv = \int \frac{dx}{x}$$

$$\Rightarrow \int \left\{ \frac{1}{1-v} - \frac{2}{(1-v)^2} + \frac{4}{(1-v)^3} \right\} dv = \int \frac{dx}{x}$$

$$\Rightarrow -n|1-v| - \frac{2}{(1-v)} + \frac{2}{(1-v)^2} = n|x| + nc$$

$$\Rightarrow -n|x-y| - \frac{2x}{x-y} + \frac{2x^2}{(x-y)^2} = nc$$

Since  $y(1) = 0$

$$\Rightarrow nc = 0$$

$$\text{Hence } \frac{2x^2}{(x-y)^2} = n|x-y| + \frac{2x}{x-y}$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/resonanceedu](https://www.facebook.com/resonanceedu) | [twitter.com/resonanceedu](https://twitter.com/resonanceedu) | [www.youtube.com/resonance](https://www.youtube.com/resonance) | [blog.resonance.ac.in](https://www.blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2023 Solution portal

PAGE # 9



## TARGET: JEE (ADV.) 2023

Polish your subject knowledge with the guidance of  
Top Notch Sr. Faculty of Resonance

# SPARK

15 WEEKS COMPAC COURSE

OFFLINE / ONLINE

CLASS  
STARTS

6<sup>th</sup> FEBRUARY  
2023

### ACADEMIC FEATURES

- Course Duration: **15 Weeks**
- Total No. of Lectures: **234** (P: 78 | C: 78 | M: 78)
- Duration of One Lecture: **1.5 hrs.** (90 Minutes)
- Classroom Teaching Hours: **351 Hrs.**
- Testing Duration: **60 Hrs.**
- Total Academic Hours: **411 Hrs.**

### Course Features

- Study Material
- Back up support of recorded lectures
- 2000+ Classes
- Dist/ Full Syllabus Test Series

### Facilities for Offline Students

- In-house Computer Lab
- Self Study Rooms for Boys & Girls



## TARGET: JEE (Main) 2023

Boost your Percentile with

# PERCENTILE BOOSTER COURSE

8 WEEKS COMPAC COURSE

OFFLINE / ONLINE

CLASS  
STARTS

6<sup>th</sup> FEBRUARY  
2023

### COURSE FEATURES

- Complete Course Coverage
- 25 Chapter wise Test
- Regular Practice through 35 Daily Online Practice Test
- 5 Full Syllabus Test
- 3 Joint Preparatory Test
- Approx 2500 practice Ques.
- 113 Teaching hours
- 99 Testing Hours
- Regular Test discussion classes for concept clearance
- Back up support of recorded lectures



STUDENTS FROM CLASSROOM PROGRAM (OFFLINE/ ONLINE)

**AIR 6**



**KARTHIKEYA POLISETTY**  
Roll No.: 21925115

**AIR-1 GEN-EWS**

**AIR 8**



**DHEERAJ KURUKUNDA**  
Roll No.: 21920114

Students in TOP-100 All India Ranks (AIRs)



**AIR-11**  
DEWANGANJI MALI  
Roll No.: 2119044



**AIR-15**  
ASHRUTI ANAND  
Roll No.: 21925165



**AIR-35**  
SANCHIN SHRIVYA  
Roll No.: 21925155



**AIR-50**  
ANUBHAV GARG  
Roll No.: 21230192



**AIR-54**  
SULMITRA D. NAVIK  
Roll No.: 21230554



**AIR-58**  
KARISHK SHARMA  
Roll No.: 21230454

**ADMISSIONS OPEN FOR ACADEMIC SESSION 2023-24**

**TARGET: JEE (Adv.) 2024**

for Class XII Passed Student

**VISHESH COURSE**  
MODE: OFFLINE / ONLINE

CLASS STARTS  
10<sup>th</sup> & 17<sup>th</sup> April

**TARGET: JEE (Main) 2024**

for Class XII Passed Student

**ABHYAAS COURSE**  
MODE: OFFLINE / ONLINE

CLASS STARTS  
10<sup>th</sup> & 24<sup>th</sup> April

**SCHOLARSHIP ON THE BASIS OF JEE (MAIN) 2023 %ILE / AIR**

**Resonance Eduventures Limited**

REGISTERED & CORPORATE OFFICE: CG Tower, A-46 & 52, IPHA, Near City Mall, Jhalawar Road, Kota (Rajasthan) - 324005  
Tel. No.: 0744-2777777, 2777700 | CIN: U80302RJ2007PLC024029

**Social Media Connect**

 facebook.com/ResonanceEdu | 
  83007 41444 | 
  youtube.com/@ResonanceEdu | 
  in.linkedin.com/school/resonance-adventures-161/ | 
  Lma/OfficialResonance | 
  twitter.com/ResonanceEdu