

NARAYANA GRABS
THE LION'S SHARE IN JEE-ADV.2022

5 RANKS in OPEN CATEGORY
ONLY FROM NARAYANA
IN TOP 10 AIR




JEE MAIN (APRIL) 2023 (06-04-2023-FN)

Memory Based Question Paper
CHEMISTRY




CHEMISTRY

1. Predict expression for α in terms of K_{eq} and concentration C :



$$(1^*) \left(\frac{K_{eq}}{108C^4} \right)^{1/5} \quad (2) \left(\frac{K_{eq}}{5C^4} \right)^{1/5} \quad (3) \left(\frac{4K_{eq}}{5C^4} \right)^{1/5} \quad (4) \left(\frac{9K_{eq}}{5C^4} \right)^{1/5}$$



C




$$K_{eq} = \frac{(2C\alpha)^2(3C\alpha)^3}{C}$$

$$K_{eq} = 108C^4 \alpha^5$$

$$\alpha = \left(\frac{K_{eq}}{108C^4} \right)^{1/5}$$

2. Radius of first orbit of hydrogen atom is 51 pm. Determine the radius of 5th orbit of Li^{2+}

Ans. 425 pm



$$(r_H^{2+})_5 = (r_H)_1 \times \frac{n^2}{Z} = 51 \times \frac{5^2}{3} = 425 \text{ pm}$$

3. How many moles of $Ba_3(PO_4)_2$ will be formed by the reaction of 5 moles of $BaCl_2$ and 3 moles of $Na_3(PO_4)$.

Ans. $\frac{5}{3}$



$$\text{Moles of } Ba_3(PO_4)_2 = \frac{5}{3}$$

4. In which of the following pairs of elements electron gain enthalpy difference is highest ?

- (1) Cl, Ar (2) Cl, Ne (3) F, Ar (4) F, Ne

Ans. (2)

Sol. Chlorine has most negative ΔH_{eg} (-349 kJ/mole) whereas Neon has most positive ΔH_{eg} (116 kJ/mole)

5. In an ionic solid element Y crystallises in ccp lattice and element X occupy $\frac{1}{3}$ rd of tetrahedral void.

Find formula of ionic solid.

Ans. X_2Y_3

Sol. For 1 unit cell,

No. of particles

$$X \quad \frac{1}{3} \times 8$$

$$Y \quad 4$$

$$\therefore \text{Formula of Ionic solid} = X_{8/3}Y_4 = X_2Y_3$$

6. The value of $\log_{10}K$ for a reaction $A \rightleftharpoons B$ is

(Given $\Delta H_{298K}^\circ = -54.67$ kJmol⁻¹)

$$\Delta S_{298K}^\circ = 10 \text{ kJmol}^{-1}$$

and $R = 8.314$ JK⁻¹mol⁻¹

$$2.303 \times 8.314 \times 298 = 5705$$

Ans. 10

Sol. $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$

$$= -54.07 \times 1000 - 298 \times 10$$

$$= -57050$$

$$\Delta G^\circ = -2.303 RT \log_{10}K$$

$$\log K = 10$$

7. Determine the amount of urea (NH_2CONH_2) to be added in 1000 g of water to decrease its vapour pressure by 25%.

Sol. $\frac{P^\circ - P_s}{P^\circ} = \frac{n}{N+n} = \frac{1}{4}$

$$\Rightarrow 4n = N + n$$

$$n = \frac{N}{3} = \left(\frac{1000}{18} \right) \times \frac{1}{3}$$

$$\therefore \text{Amount of urea is } \frac{(1000)}{18 \times 3} \times 60 = \frac{10000}{9} \text{ gm}$$

$$\approx 1111.1 \text{ gram}$$


8. Which of the following slows down the process of setting of the cement ?

Ans. Gypsum

9. Number of ambidentate ligands in given complex $[M(en)(SCN)_4]$:

Ans. 4

Sol. SCN^- is an ambidentate ligand S & N both are donor atom.



(A) Redox reaction

(C) Displacement reaction

(B) Combination reaction


(D) Decomposition reaction

(1*) A & B

(2) B only

(3) A & D

(4) B & D



It is a redox, displacement reaction.

11. A \Rightarrow Spin only magnetic moment of $[Fe(CN)_6]^{3-}$ is 1.73 B.M. and $[Fe(H_2O)_6]^{3+}$ is 5.92 B.M.

R \Rightarrow In both cases Fe have +3 oxidation state

Ans. Both A & R are correct but R is not the correct explanation

Sol. $[Fe(CN)_6]^{3-} : Fe^{+3} : 3d^5$ with S.F.L

$$\Rightarrow n = 1$$

$$\text{Magnetic moment} = 1.73 \text{ B.M}$$

$[Fe(H_2O)_6]^{3+} : Fe^{+3} : 3d^5$ with W.F.L

$$\Rightarrow n = 5$$

$$\text{Magnetic moment} = 5.92 \text{ B.M}$$

12. Assertion: Radius of H^+ is 1.5×10^{-3} pm

Reason: H^+ cannot exist independently

Sol. Both assertion and reason are correct but reason is not a correct explanation of assertion.

13. Oxidation number of Mo in Ammonophosphomolybdate

Ans. 6

Sol. $(\text{NH}_4)_3\text{PMo}_{12}\text{O}_{40}$ or $(\text{NH}_4)_3\text{PO}_4 \cdot 12\text{MoO}_3$





$$+3 + 5 + 12x - 80 = 0$$

$$12x = 80 - 8$$

$$12x = 72$$

$$x = 6$$

14. Which of following are reducing and oxidising agent respectively.



Ans. (1)

Sol. $\text{Eu}^{2+} \longrightarrow \text{Eu}^{3+} + e^-$

$\text{Eu}^{2+} \longrightarrow$ Good reducing agent

$e^- + \text{Ce}^{4+} \longrightarrow \text{Ce}^{3+}$

Ce^{4+} is a good oxidising agent

15. Column-I

Column-II

(P) N_2O_5

(i) N-N bond

(Q) N_2O

(ii) N-O-N bond





(R) N_2O_4

(iii) N=N / N≡N bond

(S) NO_2

(iv) N=O bond


Ans. P - (ii), Q - (iii), R - (i), S - (iv)







16. Polymer which is named as orlon


- (1) Polyamide (2) Polyacrylonitrile
 (3) Polycarbamate (4) Polyethene

Ans. (2)

17.  Major product

- (1)  (2)  (3)  (4) 

Ans. (2)



18. Column I

- (i) Vitamin A
 (ii) Vitamin C (Ascorbic acid)
 (iii) Riboflavin
 (iv) Thiamine

- (1) i→c, ii→d, iii→a, iv→b
 (3) i→d, ii→c, iii→b, iv→a

Column II

- (a) Beri-beri
 (b) Cheilosis
 (c) Xerophthalmia
 (d) Scurvy

- (2) i→c, ii→d, iii→b, iv→a
 (4) i→c, ii→b, iii→d, iv→a

Ans. (2)

19. Photochemical smog found mainly in

- (1) Industrial area (2) Marshy place
 (3) Hilly area of Himachal (4) Cold humid climate

Ans. (1)

20. Column I (Chemical reactions)

- (i) Glucose \rightarrow CO₂ + Ethanol
 (ii) Sucrose \rightarrow Glucose + Fructose
 (iii) Starch \rightarrow Maltose
 (iv) Protein \rightarrow Amino acids
 (1) i \rightarrow c, ii \rightarrow d, iii \rightarrow b, iv \rightarrow a
 (3) i \rightarrow c, ii \rightarrow d, iii \rightarrow a, iv \rightarrow b


Column II (Enzymes used)

- (a) Pepsin
 (b) Diastase
 (c) Zymase
 (d) Invertase
 (2) i \rightarrow d, ii \rightarrow c, iii \rightarrow b, iv \rightarrow a
 (4) i \rightarrow c, ii \rightarrow b, iii \rightarrow d, iv \rightarrow a

Ans. (1)

21. How many bromo products are formed when ethane is reacted with excess of Br₂ on heating?


Ans. (9)




22. Match the following with the correct name of reaction



(P) Gattermann Koch reaction




(Q) Hell Volhard Zelinsky



(R) Iodoform reaction

- (1) (I) \rightarrow (Q), (II) \rightarrow (R), (III) \rightarrow (P)
 (2) (I) \rightarrow (R), (II) \rightarrow (Q), (III) \rightarrow (P)
 (3) (I) \rightarrow (Q), (II) \rightarrow (P), (III) \rightarrow (R)
 (4) (I) \rightarrow (P), (II) \rightarrow (Q), (III) \rightarrow (R)


Ans. (1)











Which of the following statement is correct ?

- (1) Acetic acid solvent can take in above reaction.
- (2) NaI is soluble in acetone but NaBr is precipitate in acetone
- (3) NaI is precipitated in acetone but NaBr is soluble in acetone
- (4) When acetone is taken in solvent transition state is highly polar

Ans. (2)



Product (P) and (Q) are respectively

- (1)  and 
- (2)  and 
- (3)  and 
- (4)  and 

Ans. (3)