

# QUESTIONS & SOLUTIONS

Reproduced from Memory Retention

 06 APRIL, 2023

 9:00 AM to 12:00 Noon

SHIFT - 1

Duration : 3 Hours

Maximum Marks : 300

## SUBJECT - CHEMISTRY

**LEAGUE OF TOPPERS (Since 2020)**  
TOP 100 AIRs IN JEE ADVANCED

AIR  
**5**  
MAYANK  
MOTWANI  
Roll No. : 20771637  
JEE Adv, 2022



AIR  
**15**  
DHANANJAY  
KEJRIWAL  
Roll No. : 19821428  
JEE Adv, 2020



AIR  
**29**  
SANKALP  
PARASHAR  
Roll No. : 20975666  
JEE Adv, 2020



AIR  
**30**  
AARYAN  
GUPTA  
Roll No. : 20975800  
JEE Adv, 2020



AIR  
**67**  
ANKAN  
SARKAR  
Roll No. : 20771420  
JEE Adv, 2021



Admission Announcement for **JEE Advanced** (For Session 2023-24)

VIKAAS  
**For Class X to XI**  
Moving Students  
**TARGET 2025**

VISHWAAS  
**For Class XI to XII**  
Moving Students  
**TARGET 2024**


VISHESH  
**For Class XII**  
Passed Students  
**TARGET 2024**

Starting From : **12 & 19 APRIL '23**


Avail Scholarship up to **90%** through **R-NET** on **EVERY SUNDAY**

**CHEMISTRY**

1. Predict expression for  $\alpha$  in terms of  $K_{eq}$  and concentration C :



(1\*)  $\left(\frac{K_{eq}}{108C^4}\right)^{1/5}$       (2)  $\left(\frac{K_{eq}}{5C^4}\right)^{1/5}$       (3)  $\left(\frac{4K_{eq}}{5C^4}\right)^{1/5}$       (4)  $\left(\frac{9K_{eq}}{5C^4}\right)^{1/5}$



C



$$K_{eq} = \frac{(2C\alpha)^2(3C\alpha)^3}{C}$$

$$K_{eq} = 108C^4 \alpha^5$$

$$\alpha = \left(\frac{K_{eq}}{108C^4}\right)^{1/5}$$

2. Radius of first orbit of hydrogen atom is 51 pm. Determine the radius of 5<sup>th</sup> orbit of Li<sup>2+</sup>

**Ans.** 425 pm

**Sol.**  $r_H = 51$  pm

$$(r_H^{2+})_5 = (r_H)_1 \times \frac{n^2}{Z} = 51 \times \frac{5^2}{3} = 425 \text{ pm}$$

3. How many moles of  $Ba_3(PO_4)_2$  will be formed by the reaction of 5 moles of  $BaCl_2$  and 3 moles of  $Na_3(PO_4)$ .

**Ans.**  $\frac{5}{3}$



5 mole                  3 mole

$$\text{Moles of } Ba_3(PO_4)_2 = \frac{5}{3}$$

4. In which of the following pairs of elements electron gain enthalpy difference is highest ?

- (1) Cl, Ar                      (2) Cl, Ne                      (3) F, Ar                      (4) F, Ne

**Ans. (2)**

**Sol.** Chlorine has most negative  $\Delta H_{eg}$  ( $-349$  kJ/mole) whereas Neon has most positive  $\Delta H_{eg}$  (116 kJ/mole)

5. In an ionic solid element Y crystallises in ccp lattice and element X occupy  $\frac{1}{3}$ <sup>rd</sup> of tetrahedral void.

Find formula of ionic solid.

**Ans.**  $X_2Y_3$

**Sol.** For 1 unit cell,

No. of particles

$$X \quad \frac{1}{3} \times 8$$

$$Y \quad 4$$

$$\therefore \text{Formula of Ionic solid} = X_{8/3}Y_4 = X_2Y_3$$

6. The value of  $\log_{10}K$  for a reaction  $A \rightleftharpoons B$  is

(Given  $\Delta H^\circ_{298K} = -54.67$  kJmol<sup>-1</sup>

$$\Delta S^\circ_{298K} = 10$$
 kJmol<sup>-1</sup>

and  $R = 8.314$  JK<sup>-1</sup>mol<sup>-1</sup>

$$2.303 \times 8.314 \times 298 = 5705)$$

**Ans. 10**

**Sol.**  $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$

$$= -54.07 \times 1000 - 298 \times 10$$

$$= -57050$$

$$\Delta G^\circ = -2.303 RT \log_{10}K$$

$$\log K = 10$$

7. Determine the amount of urea ( $NH_2CONH_2$ ) to be added in 1000 g of water to decrease its vapour pressure by 25%.

**Sol.** 
$$\frac{P^\circ - P_s}{P^\circ} = \frac{n}{N+n} = \frac{1}{4}$$

$$\Rightarrow 4n = N + n$$

$$n = \frac{N}{3} = \left( \frac{1000}{18} \right) \times \frac{1}{3}$$

$$\therefore \text{Amount of urea is } \frac{(1000)}{18 \times 3} \times 60 = \frac{10000}{9} \text{ gm}$$

$$\approx 1111.1 \text{ gram}$$


8. Which of the following slows down the process of setting of the cement ?

Ans. Gypsum

9. Number of ambidentate ligands in given complex  $[M(en)(SCN)_4]$  :

Ans. 4

Sol.  $SCN^-$  is an ambidentate ligand S & N both are donor atom.



(A) Redox reaction

(C) Displacement reaction

(B) Combination reaction


(D) Decomposition reaction

(1\*) A & B

(2) B only

(3) A & D

(4) B & D



It is a redox, displacement reaction.

11. A  $\Rightarrow$  Spin only magnetic moment of  $[Fe(CN)_6]^{-3}$  is 1.73 B.M. and  $[Fe(H_2O)_6]^{+3}$  is 5.92 B.M.

R  $\Rightarrow$  In both cases Fe have +3 oxidation state

Ans. Both A & R are correct but R is not the correct explanation

Sol.  $[Fe(CN)_6]^{-3} : Fe^{+3} : 3d^5$  with S.F.L

$$\Rightarrow n = 1$$

$$\text{Magnetic moment} = 1.73 \text{ B.M}$$

$[Fe(H_2O)_6]^{+3} : Fe^{+3} : 3d^5$  with W.F.L

$$\Rightarrow n = 5$$

$$\text{Magnetic moment} = 5.92 \text{ B.M}$$

12. Assertion: Radius of  $H^+$  is  $1.5 \times 10^{-3}$  pm

Reason:  $H^+$  cannot exist independently

Sol. Both assertion and reason are correct but reason is not a correct explanation of assertion.

13. Oxidation number of Mo in Ammonophosphomolybdate

Ans. 6

Sol.  $(\text{NH}_4)_3\text{PMo}_{12}\text{O}_{40}$  or  $(\text{NH}_4)_3\text{PO}_4 \cdot 12\text{MoO}_3$

$$+3 + 5 + 12x - 80 = 0$$

$$12x = 80 - 8$$

$$12x = 72$$

$$x = 6$$

14. Which of following are reducing and oxidising agent respectively.

(1)  $\text{Eu}^{+2}$ ,  $\text{Ce}^{+4}$

(2)  $\text{Ce}^{+3}$ ,  $\text{Ce}^{+4}$

(3)  $\text{Eu}^{+4}$ ,  $\text{Eu}^{+2}$

(4)  $\text{Tb}^{+2}$ ,  $\text{Ce}^{2+}$

Ans. (1)

Sol.  $\text{Eu}^{2+} \longrightarrow \text{Eu}^{3+} + e^-$

$\text{Eu}^{2+} \longrightarrow$  Good reducing agent

$e^- + \text{Ce}^{4+} \longrightarrow \text{Ce}^{3+}$

$\text{Ce}^{4+}$  is a good oxidising agent

15. Column-I

Column-II

(P)  $\text{N}_2\text{O}_5$

(i) N-N bond

(Q)  $\text{N}_2\text{O}$

(ii) N-O-N bond




(R)  $\text{N}_2\text{O}_4$

(iii) N=N / N≡N bond

(S)  $\text{NO}_2$

(iv) N=O bond


Ans. P - (ii), Q - (iii), R - (i), S - (iv)







16. Polymer which is named as orlon


- (1) Polyamide (2) Polyacrylonitrile  
(3) Polycarbamate (4) Polyethene

Ans. (2)

17.  Major product

- (1)  (2)  (3)  (4) 

Ans. (2)



18. Column I

- (i) Vitamin A  
(ii) Vitamin C (Ascorbic acid)  
(iii) Riboflavin  
(iv) Thiamine

- (1) i→c, ii→d, iii→a, iv→b  
(3) i→d, ii→c, iii→b, iv→a

Column II

- (a) Beri-beri  
(b) Cheilosis  
(c) Xerophthalmia  
(d) Scurvy

- (2) i→c, ii→d, iii→b, iv→a  
(4) i→c, ii→b, iii→d, iv→a

Ans. (2)

19. Photochemical smog found mainly in

- (1) Industrial area (2) Marshy place  
(3) Hilly area of Himachal (4) Cold humid climate


Ans. (1)

- | 20. Column I (Chemical reactions)  | Column II (Enzymes used)   |
|--|--|
| (i) Glucose $\rightarrow$ CO <sub>2</sub> + Ethanol                                | (a) Pepsin   |
| (ii) Sucrose $\rightarrow$ Glucose + Fructose                                      | (b) Diastase   |
| (iii) Starch $\rightarrow$ Maltose   | (c) Zymase   |
| (iv) Protein $\rightarrow$ Amino acids   | (d) Invertase  |
| (1) i $\rightarrow$ c, ii $\rightarrow$ d, iii $\rightarrow$ b, iv $\rightarrow$ a | (2) i $\rightarrow$ d, ii $\rightarrow$ c, iii $\rightarrow$ b, iv $\rightarrow$ a |
| (3) i $\rightarrow$ c, ii $\rightarrow$ d, iii $\rightarrow$ a, iv $\rightarrow$ b | (4) i $\rightarrow$ c, ii $\rightarrow$ b, iii $\rightarrow$ d, iv $\rightarrow$ a |

Ans. (1)

21. How many bromo products are formed when ethane is reacted with excess of Br<sub>2</sub> on heating?

Ans. (9)




22. Match the following with the correct name of reaction



- (1) (I)  $\rightarrow$  (Q), (II)  $\rightarrow$  (R), (III)  $\rightarrow$  (P)  
 (2) (I)  $\rightarrow$  (R), (II)  $\rightarrow$  (Q), (III)  $\rightarrow$  (P)  
 (3) (I)  $\rightarrow$  (Q), (II)  $\rightarrow$  (P), (III)  $\rightarrow$  (R)  
 (4) (I)  $\rightarrow$  (P), (II)  $\rightarrow$  (Q), (III)  $\rightarrow$  (R)


Ans. (1)











Which of the following statement is correct ?

- (1) Acetic acid solvent can take in above reaction.
- (2) NaI is soluble in acetone but NaBr is precipitate in acetone
- (3) NaI is precipitated in acetone but NaBr is soluble in acetone
- (4) When acetone is taken in solvent transition state is highly polar

Ans. (2)



Product (P) and (Q) are respectively

- (1)  and 
- (2)  and 
- (3)  and 
- (4)  and 

Ans. (3)



# SATYAM CHAKRAVORTY

(Classroom) →→

selected for



**ASIAN PACIFIC MATHEMATICS  
OLYMPIAD (APMO) 2023**



**IMOTC 2023 Camp** (Conducted by HBCSE)



**BRONZE MEDAL IN INOI**

(Indian National Olympiad in Informatics 2023)



Success **Delivered** to the **Deserving**



**RELIABLE INSTITUTE** : A-10, Road No.1, IPIA, Kota-324005 (Rajasthan), India

Tel. : 0744-3535544, 2665544 | Website : [www.reliablekota.com](http://www.reliablekota.com) | E-mail : [info@reliablekota.com](mailto:info@reliablekota.com)