


CHEMISTRY

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer:




- (P) Number of intermediates = 2
 (Q) Number of transition states = 3
 (R) Reaction is endothermic
 Correct statement is
 (1) P & Q only
 (2) P & R only
 (3) Q & R only
 (4) P, Q, R

Answer (1)

Sol. 3-step reaction

Number of transition states = 3
 Number of intermediates = 2
 Reaction is exothermic
 As $\Delta H < 0$

2. Which of the following compound is most acidic?



Answer (2)

Sol. is most acidic due to -I effect of -NO₂ group.

3. Which of the following is most basic

- (1) Tl₂O₃
 (2) Tl₂O
 (3) Cr₂O₃
 (4) B₂O₃

Answer (2)

Sol. Tl⁺ oxide is more basic than Tl³⁺ Cr₂O₃ is amphoteric

4. Which of the following element is not present in Nessler's reagent?

- (1) K
 (2) Hg
 (3) N
 (4) I

Answer (3)


Sol. Nessler's reagent is alkaline solution of K₂Hgl₄

5. Which of the following is not obtained on electrolysis of brine solution


- (1) NaOH
 (2) H₂ gas
 (3) Cl₂ gas
 (4) Na

Answer (4)

Sol. Anode



Cathode




Na metal is not obtained on electrolysis of brine.

6. BeCl₂ exists as in solid state, vapour phase and at high temperature of the order of 1200 K in that order.

- (1) Polymer, Dimer and Monomer
 (2) Dimer, Polymer and Monomer
 (3) Monomer, Dimer and Polymer
 (4) Polymer, Monomer and Dimer

Answer (1)

Sol. BeCl_2 has a linear polymeric chain structure with Be-atom undergoing sp^3 hybridisation. In the vapour phase BeCl_2 tends to form a chloro-bridged dimer,



which dissociates into the linear monomer at high temperature of the order of 1200 K.

7. Which of the following has highest hydration energy.

- (1) Be^{+2}
- (2) Mg^{+2}
- (3) Ca^{+2}
- (4) Ba^{+2}

Answer (1)


Sol. Hydration energy decreases down the group in the 2nd group metal cation.

8. Oxidation state of Mn in KMnO_4 changes by 3 units in which medium?

- (1) Strongly acidic
- (2) Strongly basic
- (3) Aqueous neutral
- (4) Weakly acidic

Answer (3)

Sol. KMnO_4 in aqueous neutral medium reduces to MnO_2 .



\therefore Oxidation state of Mn in KMnO_4 changes from +7 to +4 i.e., by 3 units.


9. IUPAC name of the compound $\text{K}_3[\text{Co}(\text{C}_2\text{O}_4)_3]$ is

- (1) Potassium trioxalatocobalt (III)
- (2) Potassium trioxalatocobaltate (III)
- (3) Potassium cobalttrioxalate (II)
- (4) Potassium oxalatocobaltate (III)





Answer (2)

Sol. IUPAC name of $\text{K}_3[\text{Co}(\text{C}_2\text{O}_4)_3]$ is Potassium trioxalatocobaltate (III).

10. Consider the following reaction




Select the P.

- (1) 
- (2) 
- (3) 
- (4) 

Answer (1)

Sol.



11. During detection of Lead.





Formation of which of following compound is not used as confirmatory test.

- (1) PbSO_4
- (2) $\text{Pb}(\text{NO}_3)_2$
- (3) PbCrO_4
- (4) PbI_2



Answer (2)

Sol. PbSO_4	-	White ppt
PbCrO_4	-	Yellow ppt
PbI_2	-	Yellow ppt
$\text{Pb}(\text{NO}_3)_2$	-	Soluble

12. Identify the final product (B) formed in the following sequence of reactions.



Answer (3)



Sol.

13. Consider the following:

- D.D.T.
- Aldrin
- Sodium arsenite
- Sodium chlorate

How many of these are pesticides?

- 1
- 2
- 3
- 4

Answer (2)

Sol. D.D.T. and Aldrin are pesticides while sodium arsenite and sodium chlorate are herbicides.

14. **Amino Acid** **Letter code**

- | | |
|------------------|------|
| A. Alanine | P. N |
| B. Asparagine | Q. A |
| C. Aspartic acid | R. R |
| D. Arginine | S. D |

- A - Q; B - S; C - P; D - R
- A - Q; B - S; C - R; D - P
- A - S; B - P; C - R; D - Q
- A - S; B - P; C - P; D - R

Answer (1)


- | | | |
|---------------|---|---|
| Sol. Alanine | - | A |
| Arginine | - | R |
| Aspartic acid | - | D |
| Asparagine | - | N |

-
-
-
-
-
-
-
-
-
-


SECTION - B

Numerical Value Type Questions: This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g., 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

21. The number of compounds that give iodoform test



Answer (02.00)



give iodoform test.

22. If a_0 is the radius of H-atom de-Broglie wavelength of e^- in 3rd orbit of Li^{2+} ion is $x\pi a_0$. Find out x.

Answer (02.00)

Sol. $r_3 = \frac{a_0 \times (3)^2}{(3)} = 3a_0$

$$2\pi r = 3\lambda$$

$$2\pi(3a_0) = 3\lambda$$

$$\Rightarrow \lambda = 2\pi a_0$$

$$x = 2$$

23. How many of the following will have same relative lowering in vapour pressure?

- (A) 1 M NaCl
- (B) 1 M Urea
- (C) 1.5 M AlCl_3
- (D) 2 M Na_2SO_4

Answer (02.00)

Sol. $\frac{\Delta P}{P_{\text{solvent}}} = i(x_{\text{solute}})$

i.M should be same

- (A) $1 \times 2 = 2$
- (B) $1 \times 1 = 1$
- (C) $1.5 \times 4 = 6$
- (D) $2 \times 3 = 6$

(C) & (D) will have same RLVP

24. We are given with 7 type of lattice.

- A. Cubic
- B. tetragonal


- C. Orthorhombic
- D. Hexagonal
- E. Rhombohedral
- F. Monoclinic
- G. Triclinic

How many of them can have BCC unit cell?

Answer (03.00)

Sol. Cubic, tetragonal and orthorhombic can have BCC unit cell.

25. How many of the given molecules are square planar in shape?



Answer (04.00)

- Sol.** XeF_4 : square planar
 SF_4 : see saw
 $[\text{Ni}(\text{CO})_4]$: tetrahedral
 $[\text{Ni}(\text{CN})_4]^{2-}$: square planar
 $[\text{NiCl}_4]^{2-}$: tetrahedral
 $[\text{FeCl}_4]^{2-}$: tetrahedral
 $[\text{Cu}(\text{NH}_3)_4]^{2+}$: square planar
 $[\text{PdCl}_4]^{2-}$: square planar

26. Volume of HBr (0.02 M) (in ml) needed to completely neutralise $\text{Ba}(\text{OH})_2$ (0.01 M, 10 ml)

Answer (10)

Sol. mEq of HBr = mEq of $\text{Ba}(\text{OH})_2$

$$0.02 \times V = 0.01 \times 10 \times 2$$

$$V = \frac{0.02 \times 10}{0.02} = 10 \text{ ml}$$

- 27.
- 28.
- 29.
- 30.