

**44**  
YEARS  
OF EXCELLENCE



**NARAYANA**  
EDUCATIONAL INSTITUTIONS

**NARAYANA GRABS**  
**THE LION'S SHARE IN JEE-ADV.2022**

**5** RANKS in OPEN CATEGORY  
ONLY FROM NARAYANA  
IN TOP 10 AIR



**JEE MAIN (APRIL) 2023 (06-04-2023-AN)**

*Memory Based Question Paper*

**PHYSICS**



Toll Free: 1800 102 3344

 SUBSCRIBE /TheNarayanaGroup

PHYSICS

SECTION - A

**Multiple Choice Questions:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

**Choose the correct answer:**

1. An object starts moving with an initial speed 10 m/s and acceleration 2 m/s<sup>2</sup> along positive x-direction. The time taken to attain 60 m/s speed is  
 (1) 25 s (2) 20 s  
 (3) 30 s (4) 15 s

**Answer (1)**

**Sol.**  $v = u + at$   
 $60 = 10 + (2)t$   
 $t = 25 \text{ s}$

2. Potential energy of an electron is defined as  $U = \frac{1}{2}mw^2x^2$  and follows Bohr's law. Radius of orbital as function of  $n$  depends on ( $w$  is some constant)

- (1)  $n^2$  (2)  $\frac{1}{\sqrt{n}}$   
 (3)  $\sqrt{n}$  (4)  $n^{2/3}$

**Answer (3)**

**Sol.**  $U = \frac{1}{2}mw^2x^2$

$mvx = \frac{nh}{2\pi}$

$\frac{mv^2}{x} = mw^2x$

$\Rightarrow v = wx$

$\Rightarrow x^2 \propto n$

or  $x \propto \sqrt{n}$

3. If  $W$  is the weight on the surface of earth then weight of same body at a height  $\frac{R_e}{4}$  above the surface of earth is equal to ( $R_e$  :- Radius of earth)

- (1)  $\frac{4}{5}W$  (2)  $\frac{16}{25}W$   
 (3)  $\frac{25}{16}W$  (4)  $\frac{5}{4}W$


**Answer (2)**

**Sol.**  $g' = \frac{g \times R^2}{25 R^2} = \frac{16g}{25}$

$W' = \frac{16W}{25}$

4. A proton is projected with speed  $v$  in magnetic field  $B$  of magnitude 1 T if angle between velocity and magnetic field is  $60^\circ$  as shown below. Kinetic energy of proton is 2 eV (Mass of proton =  $1.67 \times 10^{-27}$  kg,  $e = 1.6 \times 10^{-19}$  C).

The pitch of the path of proton is approximately



- (1)  $6.28 \times 10^{-2} \text{ m}$  (2)  $6.28 \times 10^{-4} \text{ m}$   
 (3)  $3.14 \times 10^{-2} \text{ m}$  (4)  $3.14 \times 10^{-4} \text{ m}$

**Answer (2)**

**Sol.**  $R = \left( \frac{mv \sin 60^\circ}{qB} \right)$

$T = \left( \frac{2\pi m}{qB} \right)$

$\text{K.E} = \frac{1}{2}mv^2$

$\text{Pitch} = v \cos \theta \times \left( \frac{2\pi m}{qB} \right)$   $\sqrt{\frac{2K}{m}} = v$


$= \sqrt{\frac{2K}{m}} \times \frac{1}{2} \times \frac{2\pi m}{qB}$

$= \frac{\pi}{eB} \cdot \sqrt{2Km} = \frac{3.14}{1.6 \times 10^{-19} \times 1} \times \sqrt{2 \times 2 \times (1.6)^2 \times 10^{-46}}$

$= \frac{3.14}{1.6 \times 10^{-19}} \times 2 \times 1.6 \times 10^{-23}$

$= 6.28 \times 10^{-4} \text{ m}$

5. An electric dipole is shown in the figure. If it is displaced angularly by a small angle with respect to electric field, then angular frequency of oscillation is given by



(1)  $\sqrt{\frac{6qE}{ml}}$


(2)  $\sqrt{\frac{3qE}{ml}}$

(3)  $\sqrt{\frac{2qE}{ml}}$

(4)  $\sqrt{\frac{qE}{ml}}$



11. An object A is released from a height  $h$  such that the ratio of its speed before striking the ground and after striking the ground is 4 : 1. If loss of kinetic energy is  $\frac{x}{4}\%$  then value of  $x$  is



- (1) 225                      (2) 50  
(3) 375                      (4) 25

**Answer (3)**

**Sol.**  $\frac{V_{\text{before}}}{V_{\text{after}}} = \frac{4}{1}$

$$\frac{KE_{\text{before}}}{KE_{\text{after}}} = \frac{16}{1}$$

$$\frac{\Delta KE}{KE_{\text{before}}} = \frac{15}{16}$$

$$= \frac{15}{16} \times 100\%$$

$$= \frac{375}{4}\%$$

12. **Assertion (A)** : When tooth paste is pressed, it follows Pascal's principle.  
**Reason (R)** : When pressure is applied on a fluid it is distributed constantly throughout the fluid in all direction and on the wall of the container.
- (1) A is correct and R is the correct explanation of A  
(2) A is correct and R is wrong explanation of A  
(3) A is correct, R is wrong  
(4) Both A and R are false

**Answer (1)**

**Sol.** Reason (R) is the Pascal's principle and which gives correct explanation of A.

13. **Assertion (A)** : In forward biased p-n junction, diffusion current is from p-region to n-region.  
**Reason (R)** : Diffusion takes place due to concentration gradient.
- (1) Both (A) and (R) are true, (R) is the correct explanation of (A)  
(2) Both (A) and (R) are true, (R) is not the correct explanation of (A)  
(3) (A) is true, (R) is false  
(4) Both (A) and (R) are false

**Answer (1)**

**Sol.** Diffusion takes place due to concentration gradient.

14. Radius of first orbit in H-atom is  $a_0$ . Then, de Broglie wavelength of electron in the third orbit is
- (1)  $3\pi a_0$                       (2)  $6\pi a_0$   
(3)  $9\pi a_0$                       (4)  $12\pi a_0$

**Answer (2)**

**Sol.**  $\lambda = \frac{4}{mv}$

$$= \frac{2\pi r}{n}$$

$$= \frac{2\pi a_0 n^2}{n}$$

$$= 2\pi a_0 (n)$$


$$= 6\pi a_0$$

15. Choose the incorrect statement from the given statements.
- (A) Planets revolve around the Sun with constant linear speed.  
(B) Energy of planet in elliptical orbit is constant.  
(C) Satellite in circular motion have constant energy.  
(D) Body falling towards the Earth results in negligible displacement of the Earth.
- (1) (A)                      (2) (B)  
(3) (C)                      (4) (D)

**Answer (1)**

**Sol.** Planet revolves around the Sun in elliptical orbit with variable speed.

16. A particle moves from A to B via C with uniform speed of  $\pi$  m/s. Average velocity during the journey is equal to



- (1)  $\sqrt{3}$  m/s  
(2)  $\frac{\sqrt{3}}{2}$  m/s  
(3)  $\frac{3\sqrt{3}}{2}$  m/s  
(4) 2 m/s

**Answer (3)**

**Sol.** Displacement

$$= 2R \times \sin 60^\circ = \sqrt{3} R$$

$$\text{Time} = \frac{2\pi R}{3} \times \frac{1}{\pi} = \frac{2R}{3} \text{ seconds}$$

$\Rightarrow$  Average velocity

$$= \frac{\sqrt{3}R \times 3}{2R} = \frac{3\sqrt{3}}{2} \text{ m/s}$$

17. The temperature of body drops from  $60^\circ\text{C}$  to  $40^\circ\text{C}$  in 7 min. The surrounding temperature is  $10^\circ\text{C}$ . The temperature of body drops from  $40^\circ\text{C}$  to  $T^\circ\text{C}$  in 7 min. Find the value of  $T$

- (1)  $16^\circ\text{C}$  (2)  $20^\circ\text{C}$   
(3)  $28^\circ\text{C}$  (4)  $36^\circ\text{C}$

**Answer (3)**

**Sol.**  $\frac{60 - 40}{7} = K(50 - 10)$

$$\frac{40 - T}{7} = K \left( \frac{40 + T}{2} - 10 \right)$$


$$\Rightarrow \frac{20}{40 - T} = \left( \frac{40 + T}{T + 20} \right)$$

$$\Rightarrow T + 20 = 160 - 4T$$

$$\Rightarrow 5T = 140$$

$$T = \frac{140}{5} = 28^\circ\text{C}$$

18. In the given circuit, find the current passing through  $20 \Omega$ .



- (1) 0.45 A (2) 0.23 A  
(3) 0.40 A (4) 0.78 A

**Answer (3)**

**Sol.** Let  $x$  be the potential at the node

$$\frac{30 - x}{60} + \frac{2 - x}{20} + \frac{12 - x}{30} = 0$$

$$\frac{1}{2} + \frac{1}{10} + \frac{4}{10} = \frac{x}{60} + \frac{x}{20} + \frac{x}{30}$$

$$\frac{10}{10} = \frac{2x + 6x + 4x}{120}$$

$$120 = 12x$$

$$x = 10 \text{ V}$$

$$i = \frac{10 - 2}{20} = \frac{8}{20} = 0.4 \text{ A}$$

19. Average energy density of an EM wave with electric field amplitude  $E_0$  and magnetic field amplitude  $B_0$  is equal to

- (1)  $\frac{1}{2} \epsilon_0 E_0^2$  (2)  $\frac{B_0^2}{\mu_0}$   
(3)  $\epsilon_0 E_0^2$  (4)  $\frac{1}{2} \mu_0 E_0^2$

**Answer (1)**

**Sol.** Total energy =  $\frac{1}{4} \epsilon_0 E_0^2 + \frac{1}{4} \frac{B^2}{\mu_0}$

$$\text{and } \epsilon_0 E_0^2 = \frac{B^2}{\mu_0}$$

$$\Rightarrow \text{total energy} = \frac{1}{2} \epsilon_0 E_0^2$$

20.

**SECTION - B**

**Numerical Value Type Questions:** This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g., 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

21. A solid sphere and a ring have equal masses and equal radius of gyration. If both are rotating about the axis passing through the centre of mass, then the ratio of radius is  $\sqrt{\frac{x}{2}}$  then find the value of  $x$ .

**Answer (5)**

**Sol.**  $\frac{2}{5} mR_1^2 = mK_1^2$  and  $mR_2^2 = K_2^2$

$$K_1 = \sqrt{\frac{2}{5}} R_1 \quad K_2 = R_2$$

$$K_2 = R_2 \quad K_1 = K_2$$

$$\sqrt{\frac{2}{5}} R_1 = R_2 \Rightarrow \frac{R_1}{R_2} = \sqrt{\frac{5}{2}}$$

22.  
23.  
24.  
25.  
26.  
27.  
28.  
29.  
30.