

**44**  
YEARS  
OF EXCELLENCE



**NARAYANA**  
EDUCATIONAL INSTITUTIONS

**NARAYANA GRABS**  
**THE LION'S SHARE IN JEE-ADV.2022**

**5** RANKS in OPEN CATEGORY  
ONLY FROM NARAYANA  
IN TOP 10 AIR



JEE MAIN (APRIL) 2023 (08-04-2023-FN)

*Memory Based Question Paper*  
**MATHEMATICS**



Toll Free: 1800 102 3344

 SUBSCRIBE /TheNarayanaGroup

**MATHEMATICS**

**SECTION - A**

**Multiple Choice Questions:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

**Choose the correct answer:**

1. Shortest distance between lines

$$\frac{x-5}{4} = \frac{y-3}{6} = \frac{z-2}{4} \text{ and } \frac{x-3}{7} = \frac{y-2}{5} = \frac{z-9}{6} \text{ is}$$

- (1)  $\frac{190}{37}$                       (2)  $\frac{190}{\sqrt{756}}$   
 (3)  $\frac{37}{190}$                       (4)  $\frac{756}{\sqrt{190}}$

**Answer (2)**

**Sol.**  $\vec{n}_1 \times \vec{n}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 6 & 4 \\ 7 & 5 & 6 \end{vmatrix}$

$$= 16\hat{i} + 4\hat{j} - 22\hat{k}$$

$$d = \frac{(\vec{a} - \vec{b}) \cdot (\vec{n}_1 \times \vec{n}_2)}{|\vec{n}_1 \times \vec{n}_2|}$$

$$= \frac{(2\hat{i} + \hat{j} - 7\hat{k}) \cdot (16\hat{i} + 4\hat{j} - 22\hat{k})}{\sqrt{16^2 + 4^2 + (22)^2}}$$

$$= \frac{32 + 4 + 154}{\sqrt{256 + 16 + 484}}$$

$$= \frac{190}{\sqrt{756}}$$

2. Consider the word "INDEPENDENCE". The number of words such that all the vowels are together, is

- (1) 16800                      (2) 15800  
 (3) 17900                      (4) 14800

**Answer (1)**

**Sol.** Vowels: I E E E E

Consonants: N N N D D P C

I E E E E 3N, 2D, P, C


$$\begin{aligned} \text{Number of required words} &= \frac{8!}{3!2!} \times \frac{5!}{4!} \\ &= 16800 \end{aligned}$$

3. 7 boys and 5 girls are to be seated around a circular table such that no two girls sits together is

- (1)  $126(5!)^2$                       (2)  $720(5!)$   
 (3)  $720(6!)$                       (4) 720

**Answer (1)**

**Sol.**  $B_1, B_2, B_3, B_4, B_5, B_6, B_7$



Boys can be seated in  $(7 - 1)!$  ways =  $6!$

Now ways in which no two girls can be seated together is

$$6! \times {}^7C_5 \times 5!$$

$$6! \times \frac{7!}{5!2!} \times 5!$$

$$= 126(5!)^2$$

4. Consider the data :  $x, y, 10, 12, 4, 6, 8, 12$ . If mean is 9 and variance is 9.25, then the value of  $3x - 2y$  is ( $x > y$ )

- (1) 25                              (2) 1  
 (3) 24                              (4) 13

**Answer (1)**

**Sol.**  $g = \frac{52 + x + y}{8}$

$$\Rightarrow x + y = 20$$

$$9.25 = \frac{x^2 + y^2 + 100 + 144 + 16 + 36 + 64 + 144}{8} - 81$$

$$\Rightarrow 722 = x^2 + y^2 + 504$$

$$\Rightarrow x^2 + y^2 = 218$$

$$(x + y)^2 - 2xy = 218$$

$$\Rightarrow xy = 91$$

$$\therefore x = 13, y = 7$$

$$3x - 2y = 39 - 14$$

$$= 25$$

5. Coefficient independent of  $x$  in the expansion of

$$\left(3x^2 - \frac{1}{2x^5}\right)^7 \text{ is}$$

- (1)  $\frac{5103}{4}$   
 (2)  $\frac{5293}{6}$   
 (3)  $\frac{6715}{3}$   
 (4)  $\frac{7193}{4}$

**Answer (1)**

**Sol.**  $T_{r+1} = {}^7C_r (3x^2)^{7-r} \left(\frac{-1}{2x^5}\right)^r$

$$= {}^7C_r 3^{7-r} \left(\frac{-1}{2}\right)^r x^{14-7r}$$

$$\Rightarrow 14 - 7r = 0$$

$$\Rightarrow r = 2$$

$\therefore$  Coefficient of  $x^0$  is

$${}^7C_2 3^5 \times \frac{1}{4}$$

$$\frac{7 \times 6 \times 3^5}{2 \times 1 \times 4}$$

$$= \frac{5103}{4}$$

6. Dot product of two vectors is 12 and cross product is  $4\hat{i} + 6\hat{j} + 8\hat{k}$  find product of modulus of vectors

- (1)  $4\sqrt{35}$   
 (2)  $2\sqrt{65}$   
 (3)  $5\sqrt{37}$   
 (4)  $6\sqrt{37}$

**Answer (2)**

**Sol.** Let the vectors be  $\vec{a}$  and  $\vec{b}$

$$|(\vec{a} \times \vec{b})|^2 = |\vec{a}|^2 |\vec{b}|^2 - (\vec{a} \cdot \vec{b})^2$$

$$116 + 144 = (|\vec{a}| |\vec{b}|)^2$$

$$\Rightarrow |\vec{a}| |\vec{b}| = \sqrt{260}$$

7. If the coefficients of three consecutive terms in the expansion of  $(1+x)^n$  are in the ratio  $1 : 5 : 20$ , then the coefficient of the fourth term of the expansion is

- (1) 3654  
 (2) 3658  
 (3) 3600  
 (4) 1000

**Answer (1)**

**Sol.** Given  ${}^nC_{r-1} : {}^nC_r : {}^nC_{r+1} = 1 : 5 : 20$

$$\therefore \frac{n!}{(r-1)!(n-r+1)!} \times \frac{r!(n-r)!}{n!} = \frac{1}{5}$$

$$\frac{r}{n-r+1} = \frac{1}{5}$$

$$\Rightarrow n - r + 1 = 5r$$

$$n = 6r - 1 \quad \dots (i)$$

Now,

$$\frac{n!}{r!(n-r)!} \times \frac{(r+1)!(n-r-1)!}{n!} = \frac{5}{20}$$

$$\Rightarrow \frac{r+1}{n-r} = \frac{1}{4}$$

$$\Rightarrow 4r + 4 = n - r$$

$$n = 5r + 4 \quad \dots (ii)$$

By (i) and (ii)

$$5r + 4 = 6r - 1$$

$$\Rightarrow r = 5$$

and  $n = 29$

Now coefficient of fourth term

$$= {}^nC_3 = {}^{29}C_3 = 3654$$


8. The area under the curve of equations:  $x^2 \leq y$ ,  $y \leq 8 - x^2$  and  $y \leq 7$ , is

- (1)  $\frac{16}{3}$   
 (2) 18  
 (3) 20  
 (4)  $\frac{22}{3}$

**Answer (3)**



**Sol :**



Required area =  $2 \left[ \int_0^4 \sqrt{y} \, dy + \int_4^7 (\sqrt{8-y}) \, dy \right]$

$$= 2 \left[ \frac{2}{3} y^{3/2} \Big|_0^4 - \frac{(8-y)^{3/2}}{3/2} \Big|_4^7 \right]$$

$$= \frac{4}{3} (8 - (1-8))$$

$$= \frac{4}{3} (15) = 20 \text{ sq. units}$$

9.  $P = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}, Q = PAP^T$

$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ , then  $P^T Q^{2007} P = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Find  $2a + b +$

$3c - 4d$ .

(1) 2005

(2) 2007

(3) 2006

(4) 2008

**Answer (1)**

**Sol.**  $P \times P^T = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix} \begin{bmatrix} \frac{\sqrt{3}}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Similarly  $P^T P = I$

Now,  $Q^{2007} = (PAP^T)(PAP^T) \dots 2007 \text{ times}$   
 $= PA^{2007} P^T$

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$$

$$A^{2007} = \begin{bmatrix} 1 & 2007 \\ 0 & 1 \end{bmatrix}$$

$$P^T Q^{2007} P = P^T P A^{2007} P^T P = A^{2007} = \begin{bmatrix} 1 & 2007 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow a = 1, b = 2007, c = 0, d = 1$$

$$2a + b + 3c - 4d = 2 \times 1 + 2007 + 3 \times 0 - 4 \times 1 = 2005$$

10. A bolt manufacturing factory has three products A, B and C. 50% and 30% of the products are A and B type respectively and remaining are C type. Then probability that the product A is defective is 4%, that of B is 3% and that of C is 2%. A product is picked randomly and found to be defective, then the probability that it is type C.

(1)  $\frac{4}{33}$

(2)  $\frac{1}{33}$

(3)  $\frac{2}{33}$

(4)  $\frac{9}{33}$

**Answer (1)**

**Sol.** Product A is 50%, B is 30% and C is 20%

Let  $A_1$  is the event that product A is selected

$B_1$  is the event that product B is selected

$C_1$  is the event that product C is selected

and  $D$  is the event that product is defective

then,

$$P\left(\frac{D}{C_1}\right) = \frac{P(C_1)P\left(\frac{D}{C_1}\right)}{P(A_1)P\left(\frac{D}{A_1}\right) + P(B_1)P\left(\frac{D}{B_1}\right) + P(C_1)P\left(\frac{D}{C_1}\right)}$$

$$= \frac{\frac{20}{100} \times \frac{2}{100}}{\frac{50}{100} \times \frac{4}{100} + \frac{30}{100} \times \frac{3}{100} + \frac{20}{100} \times \frac{2}{100}}$$

$$= \frac{40}{200 + 90 + 40} = \frac{4}{33}$$

11. A has 5 elements and B has 2 elements. The number of subsets of  $A \times B$  such that the number of elements in subset is more than or equal to 3 and less than 6, is

(1) 602

(2) 484

(3) 582

(4) 704

**Answer (3)**

**Sol.**  $n(A) = 5, n(B) = 2$

$$\Rightarrow n(A \times B) = 10$$

$$\text{Number of subsets having 3 elements} = {}^{10}C_3$$

$$\text{Number of subsets having 4 elements} = {}^{10}C_4$$

$$\text{Number of subsets having 5 elements} = {}^{10}C_5$$

$$\therefore {}^{10}C_3 + {}^{10}C_4 + {}^{10}C_5$$

$$= 120 + 210 + 252$$

$$= 582$$

12. Check whether the function  $f(x) = \frac{(1+2^x)^7}{2^x}$  is

- (1) Even
- (2) Odd
- (3) Neither even nor odd
- (4) None of these

**Answer (3)**

**Sol.**  $f(x) = \frac{(1+2^x)^7}{2^x}$

$$f(-x) = \frac{(1+2^{-x})^7}{2^{-x}} = \frac{(2^x+1)^7}{2^{6x}}$$

$\therefore f(x)$  is neither even nor odd.

13. Let  $I(x) = \int \frac{(x+1) dx}{x(1+xe^x)^2}$ , then  $\lim_{x \rightarrow \infty} I(x) = 1$ . The value of  $I(1)$  is

- (1)  $\frac{1}{e+1} - \ln(e+1) + 1$
- (2)  $\frac{1}{e+1} - \ln(e+1)$
- (3)  $\frac{1}{e+1} - \ln(e+1) + 2$
- (4)  $\frac{1}{e+1} + 2$

**Answer (3)**

**Sol.**  $I(x) = \int \frac{(x+1) dx}{x(1+xe^x)^2}$

$$= \int \frac{e^x(x+1)}{xe^x(1+xe^x)^2} dx$$

Let  $1+xe^x = t$

$$\Rightarrow e^x(1+x) dx = dt$$

$$= \int \frac{dt}{(t-1)t^2} = -\ln t + \frac{1}{t} + \ln(t-1) + c$$

$$= -\ln(1+xe^x) + \frac{1}{x \cdot e^x + 1} + \ln(x \cdot e^x) + c$$

$$= \ln\left(\frac{xe^x}{1+xe^x}\right) + \frac{1}{xe^x+1} + c$$

$$\lim_{x \rightarrow \infty} I(x) = c = 1$$

$$\therefore I(x) = \ln\left(\frac{xe^x}{1+xe^x}\right) + \frac{1}{xe^x+1} + 1$$

$$I(1) = \ln\left(\frac{e}{1+e}\right) + \frac{1}{e+1} + 1$$

$$= 2 + \frac{1}{e+1} - \ln(1+e)$$

- 14.
- 15.
- 16.
- 17.
- 18.
- 19.
- 20.

---

**SECTION - B**

**Numerical Value Type Questions:** This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g., 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

21. If  $a_\alpha$  is the maximum value of  $a_n = \frac{n^3}{n^4 + 147}$ .

Then find  $\alpha$

**Answer (5)**

**Sol.**  $f(n) = \frac{n^3}{n^4 + 147}$

$$f'(n) = \frac{(3n^2)(n^4 + 147) - (n^3)(4n^3)}{(n^4 + 147)^2} = 0$$

$f(n) = 0$

$\Rightarrow n = \sqrt{21}$

$4 < \sqrt{21} < 5$

$a_5 > a_4$

$\therefore$  for  $n = 5$  the value is maximum

$\boxed{\alpha = 5}$

22. Maximum value  $n$  such that  $(66)!$  is divisible by  $3^n$

**Answer (31)**

**Sol.**  $\because 3$  is a prime number

$$\left[ \frac{66}{3} \right] + \left[ \frac{66}{3^2} \right] + \left[ \frac{66}{3^3} \right] + \left[ \frac{66}{3^4} \right] + \dots$$

$22 + 7 + 2 + 0 \dots$

$= 31$

$(66)! = (3)^{31} \dots$

Maximum value of  $n = 31$

23. If  $A = \begin{bmatrix} 2 & 1 & 0 \\ 1 & 2 & -1 \\ 0 & 1 & 2 \end{bmatrix}$  and  $|\text{adj}(\text{adj}(\text{adj}(A)))| = 16^n$  then

the value of  $n$  is

**Answer (06)**

**Sol.**  $|A| = 2(5) - 1(2) = 8$

$\therefore$  Now  $|\text{adj}(\text{adj}(\text{adj}(A)))| = |A|^{(n-1)^3}$   
 $= 8^8 = 16^6$

$\therefore \boxed{n = 6}$

24. The value of  $\frac{8}{\pi} \int_{\frac{\pi}{6}}^{\frac{5\pi}{6}} (8[\text{cosec}x] - 5[\cot x]) dx$  is

([.] represents greatest integer function) \_\_\_\_\_.

**Answer (56)**

**Sol.**  $\frac{8}{\pi} \int_{\frac{\pi}{6}}^{\frac{5\pi}{6}} 8[\text{cosec}x] dx - \frac{8}{\pi} \int_{\frac{\pi}{6}}^{\frac{5\pi}{6}} 5[\cot x] dx$

$$= \frac{8}{\pi} \times 8 \int_{\frac{\pi}{6}}^{\frac{5\pi}{6}} 1 \cdot dx - \frac{8}{\pi} \times 5 \left( \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} 1 dx + \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} 0 \cdot dx + \int_{\frac{\pi}{2}}^{\frac{3\pi}{4}} (-1) dx + \int_{\frac{3\pi}{4}}^{\frac{5\pi}{6}} (-2) dx \right)$$

$$= \frac{64}{\pi} \left( \frac{2\pi}{3} \right) - \frac{40}{\pi} \left( \left( \frac{\pi}{4} - \frac{\pi}{6} \right) + 0 - \left( \frac{3\pi}{4} - \frac{\pi}{2} \right) - 2 \left( \frac{5\pi}{6} - \frac{3\pi}{4} \right) \right)$$

$$= \frac{128}{3} - \frac{40}{\pi} \left( \frac{\pi}{12} - \frac{\pi}{4} - \frac{2\pi}{12} \right)$$

$$= \frac{128}{3} - 40 \left( \frac{1}{12} - \frac{1}{4} - \frac{1}{6} \right)$$

$$= \frac{128}{3} - 40 \left( \frac{1-3-2}{12} \right) = \frac{128}{3} - 40 \left( -\frac{1}{3} \right)$$

$$= \frac{168}{3}$$

$$= 56$$

25. If  $\lim_{x \rightarrow 0} \frac{1 - \cos^2 3x}{\cos^3 4x} \times \frac{\sin^3 4x}{(\log(1+2x))^5} = t$  then  $[t]$  is

(where [.] represents greatest integer fraction)

**Answer (18)**

**Sol.**  $\lim_{x \rightarrow 0} \frac{1 - \cos^2 3x}{\cos^3 4x} \times \frac{\sin^3 4x}{(\log(1+2x))^5}$

$$\lim_{x \rightarrow 0} \frac{\sin^2 3x \sin^3 4x}{\cos^3(4x) (\log(1+2x))^5}$$

$$\lim_{x \rightarrow 0} \frac{\frac{\sin^2 3x}{(3x)^2} \cdot \frac{\sin^3 4x}{(4x)^3} \cdot (3x)^2 \cdot (4x)^3}{\cos^3 4x \cdot \left( \frac{\log(1+2x)}{2x} \right)^5 (2x)^5}$$

$$= \frac{9 \times 64}{32} = 18$$

26.

27.

28.

29.

30.