

44
YEARS
OF EXCELLENCE



NARAYANA
EDUCATIONAL INSTITUTIONS

NARAYANA GRABS
THE LION'S SHARE IN JEE-ADV.2022

5 RANKS in OPEN CATEGORY
ONLY FROM NARAYANA
IN TOP 10 AIR



JEE MAIN (APRIL) 2023 (12-04-2023-FN)

Memory Based Question Paper
MATHEMATICS



Toll Free: 1800 102 3344

 SUBSCRIBE /TheNarayanaGroup

Sol. $\frac{dy}{dx} + \frac{x}{1+x^2}y = \frac{y^2}{1+x^2}$

$$\frac{1}{y^2} \frac{dy}{dx} + \frac{x}{1+x^2} \times \frac{1}{y} = \frac{1}{1+x^2}$$

Let $\frac{1}{y} = t$

$$-\frac{1}{y^2} \frac{dy}{dx} = \frac{dt}{dx}$$

$$\frac{-dt}{dx} + \left(\frac{x}{1+x^2}\right) dt = \frac{1}{1+x^2}$$

$$\frac{dt}{dx} - \left(\frac{x}{1+x^2}\right) dt = -\frac{1}{1+x^2}$$

$$IF = e^{-\int \frac{x}{1+x^2} dx} = e^{-\frac{1}{2} \log|1+x^2|} = \frac{1}{\sqrt{1+x^2}}$$

$$\frac{t}{\sqrt{1+x^2}} = -\int \frac{1}{(1+x^2)\sqrt{1+x^2}} dx$$

Let $x = \tan\theta$

$$dx = \sec^2\theta d\theta$$

$$I = \int \frac{\sec^2\theta}{\sec^2\theta \cdot \sec\theta} d\theta = \int \cos\theta = \sin\theta + C$$

$$\therefore \frac{1}{y\sqrt{1+x^2}} = -\frac{x}{\sqrt{1+x^2}} + C$$

$$\therefore y(1) = 1$$

$$\Rightarrow C = \sqrt{2}$$

$$\therefore \frac{1}{y\sqrt{1+x^2}} + \frac{x}{\sqrt{1+x^2}} = \sqrt{2}$$

$$1 + xy = \sqrt{2}y\sqrt{1+x^2}$$

Now

$$y(2\sqrt{2})$$

$$1 + 2\sqrt{2}y = 3\sqrt{2}y$$

$$\sqrt{2}y = 1$$

$$y = \frac{1}{\sqrt{2}}$$

4. For the expression $(1 - x)^{100}$. Then sum of coefficient of first 50 terms is

(1) ${}^{99}C_{49}$ (2) $-\frac{{}^{100}C_{50}}{2}$

(3) $-{}^{99}C_{49}$ (4) $-{}^{101}C_{50}$

Answer (2)

Sol. Sum of coefficient of first 50 terms

$$(t) = {}^{100}C_0 - {}^{100}C_1 + \dots + {}^{100}C_{49}$$

Now

$${}^{100}C_0 - {}^{100}C_1 + \dots + {}^{100}C_{100} = 0$$

$$2[{}^{100}C_0 - {}^{100}C_1 + \dots] + {}^{100}C_{50} = 0$$

$$\therefore t = -\frac{1}{2} {}^{100}C_{50}$$

5. Positive numbers a_1, a_2, \dots, a_5 are in geometric progression. Their mean and variance are $\frac{31}{10}$ and $\frac{m}{n}$ respectively. The mean of the

reciprocals is $\frac{31}{40}$, then $m + n$ is

(1) 209 (2) 211

(3) 113 (4) 429

Answer (2)

Sol. $a\left(\frac{1}{r^2} + \frac{1}{r} + 1 + r + r^2\right) = \frac{31}{2}$

$$\frac{1}{a}\left(\frac{1}{r^2} + \frac{1}{r} + 1 + r + r^2\right) = \frac{31}{8}$$

$$\Rightarrow a^2 = 4$$

$$\Rightarrow a = 2$$

$$\Rightarrow \frac{1}{r^2} + \frac{1}{r} + 1 + r + r^2 = \frac{31}{4}$$

$$\Rightarrow \left(r + \frac{1}{r}\right)^2 + \left(r + \frac{1}{r}\right) = \frac{31}{4} + 1 = \frac{35}{4}$$

$$4t^2 + 4t - 35 = 0$$

$$\Rightarrow t = \frac{5}{2}$$

$$\Rightarrow r = 2$$

$$\therefore \text{numbers are} = \frac{1}{2}, 1, 2, 4, 8$$

$$\begin{aligned} \therefore \sigma^2 &= \frac{1}{4} + 1 + 4 + 16 + 64 - \left(\frac{31}{10}\right)^2 \\ &= \frac{341}{20} - \frac{961}{100} \\ &= \frac{1705 - 961}{100} \\ &= \frac{744}{100} = \frac{186}{25} \end{aligned}$$

$$\begin{aligned} \therefore m + n &= 186 + 25 \\ &= 211 \end{aligned}$$

6. If $\Delta(k) = \begin{vmatrix} 1 & 2k-1 & 2k \\ n & n^2 & n(n+1) \\ \cos^2 n & \cos^2(n+1) & (n+2) \end{vmatrix}$, then

$$\sum_{k=1}^n \Delta(k) =$$

- (1) n (2) 1
 (3) $\frac{n^2}{2}$ (4) 0

Answer (4)


Sol. $\sum_{k=1}^n \Delta(k) = \begin{vmatrix} n & n^2 & n(n+1) \\ n & n^2 & n(n+1) \\ \cos^2 n & \cos^2(n+1) & (n+2) \end{vmatrix} = 0$

7. Given A, B, C represents angles of a ΔAB and $\cos A + 2 \cos B + \cos C = 2$ and $AB = 3$ and $BC = 7$ then $\cos A - \cos C$ is

- (1) $-\frac{10}{7}$ (2) $\frac{10}{7}$
 (3) $\frac{5}{7}$ (4) $-\frac{5}{7}$

Answer (1)

Sol.



$$\cos A + 2 \cos B + \cos C = 2.$$

$$\frac{9 + b^2 - 49}{6b} + 2 \left(\frac{49 + 9 - b^2}{42} \right) + \left(\frac{49 + b^2 - 9}{14b} \right) = 2$$

$$\frac{b^2 - 40}{6b} + \frac{58 - b^2}{21} + \frac{40 + b^2}{14b} = 2$$

$$\Rightarrow b = -4 \text{ or } 4 \text{ or } 5$$

b cannot be -4 and 4

$$\Rightarrow b = 5.$$

Now,

$$\cos A - \cos C$$

$$\frac{9 + 25 - 49}{2 \times 3 \times 5} - \frac{49 + 25 - 9}{2 \times 7 \times 5}$$

$$= \frac{1}{2} - \frac{13}{14} = \frac{-20}{14} = \frac{-10}{7}$$

8. Let $x^2 + \sqrt{6}x + 4 = 0$ be any quadratic equation and α, β are the roots of that equation then

$$\frac{\alpha^{34}\beta^{24} + \alpha^{32}\beta^{26} + 2\alpha^{33}\beta^{25}}{\alpha^{31}\beta^{20} + \alpha^{28}\beta^{23} + 3\alpha^{30}\beta^{21} + 3\alpha^{29}\beta^{22}}$$
 is

- (1) $-\frac{2^7}{3}\sqrt{6}$ (2) $\frac{2^7}{3}\sqrt{6}$
 (3) $-\frac{2^8}{3}\sqrt{6}$ (4) $\frac{2^8}{3}\sqrt{6}$

Answer (1)

Sol.

$$x^2 + \sqrt{6}x + 4 = 0 \begin{cases} \alpha \\ \beta \end{cases}$$

$$\therefore \alpha + \beta = -\sqrt{6}, \alpha\beta = 4$$

Now $\frac{\alpha^{34}\beta^{24} + \alpha^{32}\beta^{26} + 2\alpha^{33}\beta^{25}}{\alpha^{31}\beta^{20} + \alpha^{28}\beta^{23} + 3\alpha^{30}\beta^{21} + 3\alpha^{29}\beta^{22}}$

$$= \frac{\alpha^{32}\beta^{24} [\alpha^2 + \beta^2 + 2\alpha\beta]}{\alpha^{28}\beta^{20} [\alpha^3 + \beta^3 + 3\alpha^2\beta + 3\alpha\beta^2]}$$

$$= (\alpha\beta)^4 \frac{[(\alpha + \beta)^2]}{(\alpha + \beta)^3} = \frac{4^4}{-\sqrt{6}} = \frac{-2^7}{3}\sqrt{6}$$

9. If a plane $4x - 3y + z = 2$ is rotated by an angle of $\frac{\pi}{2}$ at intersection point of another plane $3x + 11z - 4y = 12$, then $P(2, 3, 4)$ is at what distance from resultant plane?

- (1) $\frac{250}{\sqrt{63245}}$ (2) $\frac{885}{\sqrt{66345}}$
 (3) $\frac{925}{\sqrt{66215}}$ (4) $\frac{24}{\sqrt{11235}}$

Answer (2)

Sol : Equation of required plane:

$$4x - 3y + z - 2 + \lambda(3x - 4y + 11z - 12) = 0$$

If is perpendicular to $4x - 3y + z = 2$

$$\therefore (4 + 3\lambda) \cdot 4 + (-3 - 4\lambda)(-3) + (1 + 11\lambda) \cdot 1 = 0$$

$$\Rightarrow 16 + 12\lambda + 9 + 12\lambda + 1 + 11\lambda = 0$$

$$\Rightarrow 35\lambda + 26 = 0$$

$$\Rightarrow \lambda = -\frac{26}{35}$$

$$\therefore x(4 + 3\lambda) - y(3 + 4\lambda) + z(1 + 11\lambda) - 2 - 12\lambda = 0$$

$$\Rightarrow \frac{-62x}{35} - y\left(\frac{1}{35}\right) + z\left(\frac{-250}{35}\right) + \left(\frac{242}{35}\right) = 0$$

$$\Rightarrow 62x + y + 250z = 242$$

Distance from (2, 3, 4)

$$= \frac{124 + 3 + 1000 - 242}{\sqrt{66345}}$$

$$= \frac{885}{\sqrt{66345}}$$

10. A circle with centre $z_0 = \frac{1}{2} + \frac{3i}{2}$ exist in an argand plane. A point $z_1 = 1 + i$ and z_2 lies outside the circle, such that $|z_0 - z_1| |z_0 - z_2| = 1$. Then the largest value of $|z_2|$ is

- (1) $\sqrt{5} - \sqrt{2}$ (2) $\sqrt{\frac{5}{2}} - \sqrt{2}$
 (3) $\sqrt{\frac{5}{2}}$ (4) $\sqrt{\frac{5}{2}} + \sqrt{2}$

Answer (4)


Sol. $\left| z - \frac{1}{2} - \frac{3i}{2} \right| = r \rightarrow$ Circle

Now,

$$|z_0 - z_1| |z_0 - z_2| = 1$$

$$\frac{1}{\sqrt{2}} |z_0 - z_2| = 1$$

$$|z_0 - z_2| = \sqrt{2}$$



$$\begin{aligned} \text{Max } |z_2| &= \sqrt{\frac{1}{4} + \frac{9}{4}} + \sqrt{2} \\ &= \sqrt{\frac{10}{4}} + \sqrt{2} \end{aligned}$$

$$= \left(\sqrt{\frac{5}{2}} + \sqrt{2} \right) \text{ unit}$$

11. Let $\vec{a} = \lambda\hat{i} + \hat{j} - \hat{k}$, $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$ and \vec{c} is a vector such that $(\vec{a} + \vec{b} + \vec{c}) \times \vec{c} = 0$ and $\vec{a} \cdot \vec{c} = -17$, $\vec{b} \cdot \vec{c} = -20$. Find $|\vec{c} \times (\lambda\hat{i} + \hat{j} + \hat{k})|^2$ given ($\lambda > 0$)
- (1) 46
 (2) 61
 (3) 48
 (4) 51

Answer (1)

Sol. $k(\vec{a} + \vec{b}) = \vec{c}$

$$\vec{a} \cdot \vec{c} = -17, \vec{b} \cdot \vec{c} = -20$$

$$k(\lambda^2 + 3\lambda - 1) = -17, k(3\lambda + 11) = -20$$

$$\Rightarrow \lambda = -\frac{69}{20}, 3$$

$$\lambda = 3, k = -1$$

$$\vec{c} = -1(\vec{a} + \vec{b})$$

$$= -((\lambda + 3)\hat{i} + \hat{k}) = -6\hat{i} - \hat{k}$$

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -6 & 0 & -1 \\ 3 & 1 & 0 \end{vmatrix} = \hat{i}(1) - \hat{j}(3) + \hat{k}(-6) = \hat{i} - 3\hat{j} - 6\hat{k}$$

$$|\vec{c} \times (\lambda\hat{i} + \hat{j} + \hat{k})|^2 = 46$$

12.
 13.
 14.
 15.
 16.
 17.
 18.
 19.
 20.

SECTION - B

Numerical Value Type Questions: This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g., 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

21. If $\frac{{}^n C_n}{n+1} + \frac{{}^n C_{n-1}}{n} + \dots + \frac{{}^n C_1}{2} + {}^n C_0 = \frac{255}{8}$. Then

value of n is

Answer (07)

Sol. $\int_0^1 (1+x)^n dx = \int_0^1 ({}^n C_0 + {}^n C_1 x + {}^n C_2 x^2 + \dots + {}^n C_n x^n) dx$

$$\frac{(1+x)^{n+1}}{n+1} \Big|_0^1 = {}^n C_0 x + {}^n C_1 \frac{x^2}{2} + {}^n C_2 \frac{x^3}{3} + \dots + \frac{{}^n C_n x^{n+1}}{n+1} \Big|_0^1$$

$$\frac{2^{n+1}}{n+1} - \frac{1}{n+1} = {}^n C_0 + \frac{{}^n C_1}{2} + \frac{{}^n C_2}{3} + \dots + \frac{{}^n C_n}{n+1}$$

Now

$$\frac{2^{n+1} - 1}{n+1} = \frac{255}{8}$$

$$\Rightarrow n = 7$$

22. If the value of $\int_{-0.15}^{0.15} |100x^2 - 1| dx = \frac{k}{3000}$, then the

value of k is _____.

Answer (575)

Sol. $I = 2 \int_0^{0.15} |100x^2 - 1| dx$

$$= 2 \left[\int_0^{0.1} -(100x^2 - 1) dx + \int_{0.1}^{0.15} (100x^2 - 1) dx \right]$$

$$= 2 \left[\left[x - \frac{100x^3}{3} \right]_0^{0.1} + \left[\frac{100x^3}{3} - x \right]_{0.1}^{0.15} \right]$$

$$= \frac{575}{3000}$$

$$\Rightarrow k = 575$$

23. $N > 40000$, where N is divisible by 5. How many such 5 digits numbers using 0, 1, 3, 5, 7, 9?

Answer (120)

Sol. Case I : Number starts with 5

$$\begin{array}{cccc} \underline{5} & & & \underline{0} \\ \downarrow & \downarrow & \downarrow & \\ 4 \text{ ways} & 3 \text{ ways} & 2 \text{ ways} & = 4 \times 3 \times 2 = 24 \end{array}$$

Case II : Number starts with 7

$$\begin{array}{cccc} \underline{7} & & & \\ \downarrow & \downarrow & \downarrow & \downarrow \\ 4 & 3 & 2 & 2 = 4 \times 3 \times 2 \times 2 = 48 \end{array}$$

Case III : Number starts with 9

$$\begin{array}{cccc} \underline{9} & & & \\ \downarrow & \downarrow & \downarrow & \downarrow \\ 4 & 3 & 2 & 2 = 4 \times 3 \times 2 \times 2 = 48 \end{array}$$

Total ways = 120

24. Three numbers a, b, c are in A.P. and they are used to make a 9-digits number using each digit thrice, such that at least 3 consecutive digits are in A.P. then number of such numbers is

Answer (1260)

Sol. $\boxed{a} \boxed{b} \boxed{c}$ or $\boxed{b} \boxed{c} \boxed{a}$

So, total number $\frac{{}^7 C_1 \times 2 \times 6!}{2! 2! 2!} = \frac{7!}{4}$

$$= 7 \times 6 \times 5 \times 3 \times 2 = 1260$$

$$a\hat{i} + \hat{j} + k$$

25. If $\hat{i} + b\hat{j} + k$ are co-planar then the value of

$$\hat{i} + \hat{j} + ck$$

$$\frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} \text{ is}$$

Answer (01)

Sol. $\begin{vmatrix} a & 1 & 1 \\ 1 & b & 1 \\ 1 & 1 & c \end{vmatrix} = 0$

$$R_1 \rightarrow R_1 - R_2, R_2 \rightarrow R_2 - R_3$$

$$\begin{vmatrix} a-1 & 1-b & 0 \\ 0 & b-1 & 1-c \\ 1 & 1 & c \end{vmatrix} = 0$$

$$(a-1)[c(b-1) - (1-c)] + 1[(1-b)(1-c)] = 0$$

$$c(a-1)(b-1) - (a-1)(1-c) + (1-b)(1-c) = 0$$

Multiply and divide by $(1-a)(1-b)(1-c)$

$$-\frac{1-c-1}{1-c} + \frac{1}{1-b} + \frac{1}{1-a} = 0$$

$$-1 + \frac{1}{1-c} + \frac{1}{1-b} + \frac{1}{1-a} = 0$$

$$\therefore \frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 1$$

26. $f(x) = \lfloor x \rfloor + \sqrt{x - \lfloor x \rfloor}$. The number of points of discontinuity of $f(x)$ in $[-2, 1]$ is.

Answer (2)

Sol. $f(x) = \lfloor x \rfloor + \sqrt{\{x\}}$

$x = -2$

$f(-2) = 2$

$f(-2^+) = 2 + 0 = 2$

$x = -1$

$f(-1) = 1 + 0 = 1$

$f(-1^-) = 2 + 1 = 3$

\therefore discontinuous at $x = -1$

$x = 0$

$f(0) = 0$

$f(0^-) = 1 + 1 = 2$

\therefore discontinuous at $x = 0$

$x = 1$

$f(1) = 1$

$f(1^-) = 0 + 1 = 1$

\therefore discontinuous at $x = -1$ and at $x = 0$

\therefore 2 points of discontinuity


27. Given $9x^2 + 4y^2 = 36$ and a point $P\left(\frac{2\sqrt{3}}{\sqrt{7}}, \frac{6}{\sqrt{7}}\right)$ lie

on ellipse. PQ is a diameter of ellipse and RS is a diameter which is perpendicular to PQ . If

$\frac{1}{PQ^2} + \frac{1}{RS^2} = \frac{p}{m}$ in simplest form then $p + m$ is

Answer (157)

Sol. $r_1 = \sqrt{\frac{48}{7}}$



$$\frac{r_1^2 \cos^2 \theta}{4} + \frac{r_1^2 \sin^2 \theta}{9} = 1$$

$$\frac{\cos^2 \theta}{4} + \frac{\sin^2 \theta}{9} = \frac{7}{48} \quad \dots(i)$$

$$\frac{r_2^2 \sin^2 \theta}{4} + \frac{r_2^2 \cos^2 \theta}{9} = 1$$

$$\frac{\sin^2 \theta}{4} + \frac{\cos^2 \theta}{9} = \frac{1}{r_2^2}$$

From (i), $\frac{1}{r_2^2} = \frac{1}{4} + \frac{1}{9} - \frac{7}{48} = \frac{31}{144}$

$$\frac{1}{PQ^2} + \frac{1}{RS^2} = \frac{1}{4} \left(\frac{1}{r_1^2} + \frac{1}{r_2^2} \right)$$

$$= \frac{1}{4} \left(\frac{7}{48} + \frac{31}{144} \right) = \frac{13}{144}$$

28.

29.

30.