

# Consortium of Medical Engineering and Dental Colleges of Karnataka (COMEDK-2009)

## CHEMISTRY

62. Which one of the following is an unsaturated fatty acid?  
a) Palmitic acid  
b) Lauric acid  
 c) Linolenic acid  
d) Myristic acid
63. When chlorine is passed through boiling toluene we get  
a) o-Chloro toluene  
b) p-Chloro toluene  
 c) Mixture of o & p-Chloro toluene  
d) Benzyl chloride
64. The standard temperature used in thermo chemical calculations is  
a) 273 K  
 b) 298 K  
c) 297 K  
d) 303 K
65. Which of the following is an intensive property?  
a) Enthalpy  
b) Entropy  
 c) Density  
d) Mass
66. Schiff's reagent contains  
a) Rochelle salt  
b) Resorcinol  
 c) Rosaniline  
d)  $\alpha$  naphthol
67. The formula of chromyl chloride is  
a) CrCl  
b) CrCl<sub>3</sub>  
c) CrOCl<sub>2</sub>  
 d) CrO<sub>2</sub>Cl<sub>2</sub>
68. Horn silver is  
a) Oxide ore  
b) Sulfide ore  
 c) Halide ore  
d) Carbonate ore

Physics and Chemistry

Ver. C

69. Tetrahedral structure is formed by

- a)  $sp^3$  hybridization
- b)  $sp^3$  hybridization
- c)  $dsp^3$  hybridization
- d)  $d^2sp^3$  hybridization

$sp^3$  hybridization

70.  $NO^+$  ligand is

- a) nitronium
- b) nitrosyl
- c) nitrosonium
- d) nitro

nitrosonium


71. Cationic Complex is

- a) hexa amino platinum chloride
- b) potassium ferro cyanide
- c) sodium argento cyanide
- d) nickel carbonyl

hexa amino platinum chloride

72.  $2p_x$  atomic orbital undergoes linear combination with

- a)  $2p_x$  orbital
- b)  $2p_x$  orbital
- c) Both  $2p_x$  and  $2p_z$  orbitals
- d)  $2p_z$  orbital




73. In a first order reaction, molar concentration of a reactant decreases from 0.1 to 0.01 in 100 seconds, The rate constant of the reaction is

- a) 2.3030
- b) 0.02303
- c) 0.2303
- d) 0.002303

$$k = \frac{2.303}{t} \log \frac{a}{a-x} = \frac{2.303}{100} \log \frac{0.1}{0.01} = 0.02303$$

74. In which one of the following equilibria, pressure has no effect

- a)  $PCl_5 \rightleftharpoons PCl_3 + Cl_2$
- b)  $2NH_3 \rightleftharpoons N_2 + 3H_2$
- c)  $2SO_2 + O_2 \rightleftharpoons 2SO_3$
- d)  $N_2 + O_2 \rightleftharpoons 2NO$



75. Conductivity of a solution is not affected by

- a) Addition of water
- b) Process of heating
- c) Addition of acetic acid
- d) Addition of ethanol

Addition of ethanol

Space for calculation / rough work

$$k = \frac{2.303}{100} \log \frac{0.1}{0.01}$$


$$= 2.303 \times 0.02 = 0.02303$$

Physics and Chemistry

Ver. C

76. The lowering in vapour pressure is maximum for  
 a) 0.1M urea  
 b) 0.1M NaCl  
 c) 0.1M MgCl<sub>2</sub>  
 d) 0.1M K<sub>4</sub>[Fe(CN)<sub>6</sub>]


77. Bromo ethane and isopropyl chloride with metallic sodium in ether forms  
 a) Pentane  
 b) 2-methyl butane  
 c) 3-methyl butane  
 d) 2:3 dimethyl butane




78. To dry ammonia gas the drying agent used is  
 a) Con. H<sub>2</sub>SO<sub>4</sub>  
 b) P<sub>2</sub>O<sub>5</sub>  
 c) soda lime  
 d) anhydrous CaCl<sub>2</sub>

The moisture present in ammonia can't be dried by conc. H<sub>2</sub>SO<sub>4</sub>, anhydrous CaCl<sub>2</sub> and P<sub>2</sub>O<sub>5</sub>.

79. The metal hydroxide which is soluble in excess of ammonium hydroxide is  
 a) Fe(OH)<sub>2</sub>  
 b) Fe(OH)<sub>3</sub>  
 c) Cu(OH)<sub>2</sub>  
 d) Al(OH)<sub>3</sub>



80. Potassium dichromate can be converted to potassium chromate by adding  
 a) KOH  
 b) Con. H<sub>2</sub>SO<sub>4</sub>  
 c) NH<sub>4</sub>OH  
 d) acetic acid



81. 0.5g of an acid is neutralized by 40cc of 0.125N NaOH. The equivalent mass of the acid is  
 a) 50  
 b) 100  
 c) 40  
 d) 80

100 ; Eqr weight of NaOH = 40

82. 5 liters of NaOH solution of pH 12 contains  
 a) 200g  
 b) 0.2g  
 c) 20g  
 d) 2g

pOH = 2 ; [OH<sup>-</sup>] = 1 x 10<sup>-2</sup> M  
 Weight (NaOH) = 40g  
 In 5 liters = 5 x 40 x 1 x 10<sup>-2</sup> = 2g

Space for calculation / rough work

$$E = \frac{0.5}{0.125 \times 40} = \frac{0.5 \times 1000}{1000} = 1000$$

$$E = \frac{0.5 \times 1000}{0.125 \times 40} = \frac{500}{5} = 100$$

$$10^{-2} = \frac{40}{40 \times 5}$$

$$M = \frac{100 \times 1000}{1000} = 100$$

83. 50cc of oxalic acid is oxidized by 25cc of 0.20 N  $\text{KMnO}_4$ . The mass of oxalic acid present in 500cc of the solution is

- a) 3.15g
- b) 31.5g
- c) 6.3g
- d) 63g

84. Pure water is neutral because

- a)  $\text{PH} = 7$
- b) Litmus has no effect
- c) It is free from dissolved salts
- d)  $\text{PH} = 0$

85. In the titration of Mohr salt against  $\text{KMnO}_4$ , the indicator used is

- a) diphenyl amine
- b)  $\text{KMnO}_4$
- c) phenolphthalein
- d) Methyl orange

86. The relationship between half life of a reaction and the order of reaction is

- a)  $t_{\frac{1}{2}} \propto a^{(n+1)}$
- b)  $t_{\frac{1}{2}} \propto \frac{1}{a^{(n+2)}}$
- c)  $t_{\frac{1}{2}} \propto \frac{1}{a^n}$
- d)  $t_{\frac{1}{2}} \propto \frac{1}{a^{(n-1)}}$

87. 6gm of urea is dissolved in 90g of water. Relative lowering of vapour pressure is

- a) 0.02
- b) 0.2
- c) 0.002
- d) 0.04

88. 6.84g of sucrose is dissolved in 200g of water. The molality of the solution is

- a) 0.2M
- b) 0.3M
- c) 0.1M
- d) 0.02M

When common salt is added to a saturated solution of soap, soap is precipitated. This is based on the principle of

- a)  Common ion effect
- b) Principle of solubility product
- c) Adsorption from solution
- d) Peptisation

Common ion effect

Highest osmotic pressure is shown by a solution of

- a)  0.1M Aluminium sulfate
- b) 0.1M Potassium Nitrate
- c) 0.1M Magnesium Chloride
- d) 0.1M Barium Chloride

0.1M Aluminium sulfate

50% of a first order reaction is completed in 30min. The velocity constant of the reaction is

- a) 0.231
- b) 2.31
- c) 0.00231
- d)  0.0231

$t_{1/2} = 30 \text{ min}$

$k = \frac{0.693}{30} = 0.0231$

The ebullioscopic constant is the elevation in boiling point produced by

- a) 1Molar solution
- b)  1Molal solution
- c) 1N solution
- d) 10% solution

1 Molal solution

The mass of glucose to be dissolved in 50g of water to get 0.3 Molal solution is

- a) 27g
- b) 0.27g
- c)  2.7g
- d) 5.4g

25ml of 0.08N Mohr salt solution is Oxidised by 20ml of  $K_2Cr_2O_7$  in acid medium. The Mass of Mohr salt present in 500cc is

- a) 3.96g
- b)  19.6g
- c) 39.6g
- d) 39.2g

19.6g

A reaction is spontaneous at all temperature when

- a)   $\Delta H$  is -ve and  $\Delta S$  is +ve
- b)  $\Delta H$  is +ve and  $\Delta S$  is -ve
- c) Both  $\Delta H$  &  $\Delta S$  are -ve
- d) Both  $\Delta H$  &  $\Delta S$  are +ve

$\Delta H$  is -ve and  $\Delta S$  is +ve

$Al_2(SO_4)_3$  P.N.K

$m_2 = \frac{9.309}{30} \log$

$k = \frac{0.693}{30}$

$\Delta G = -T \Delta S = -0.0231$

Space for calculation / rough work

$0.3 = \frac{3}{80} \times 20$

$3 = 400 \times 0.3$   
 $= \frac{120}{1000}$

$0.3 = \frac{m}{16}$

$m = \frac{0.3 \times 16}{1000} \times 1000$

25.0g

$25 \times 0.08 = 20 \times \frac{m}{16}$

48

32

198

96. The coordination number of sodium chloride is

- a) 4
- b) 8
- c) 6
- d) 12

97. Conjugate acid of  $NH_2^-$  is

- a)  $NH_3$
- b)  $NH_4^+$
- c)  $N^{3-}$
- d)  $NH_2^+$



98. Highest molar conductivity is given by

- a) 0.005 M NaCl
- b) 0.1 M NaCl
- c) 0.05 M NaCl
- d) 0.01 M NaCl

Molar Conductivity is defined as the conductivity of an electrolyte solution divided by molar concentration.

99. In the detection of III group basic radicals  $NH_4OH$  is added after  $NH_4Cl$  to

- a) increase in the ionization of  $NH_4OH$
- b) increase in the ionization of salt solution
- c) decrease in the ionization of salt solution
- d) decrease in the ionization of  $NH_4OH$

decrease in the ionization of  $NH_4OH$

100. Just before attaining the chemical equilibrium

- a) Rate of forward reaction decreases & Rate of backward reaction increases
- b) Rate of forward reaction increases & Rate of backward reaction decreases
- c) No change in the rates of forward & backward reactions.
- d) Rate of forward reaction equals the rate backward reaction.

101. Which one of the following shows highest magnetic moment?

- a)  $Fe^{2+}$
- b)  $Co^{2+}$
- c)  $Cr^{3+}$
- d)  $Ni^{2+}$


$Fe^{2+}$ ; becoz it has 4 unpaired electrons.

102. In 3d series as we move from scandium to zinc the paramagnetism

- a) increases
- b) decreases
- c) first increases to a maximum & then decreases
- d) first decreases to a minimum & then increases

"C"; Paramagnetism in the transition elements is caused by the presence of unpaired electrons in the  $d$  suborbital.


Space for calculation / rough work



$20 \times 2 = 40$   
 $10$   
 $100$   
 $22$   
 $108$   
 $22$   
 $200$   
 $6189 \times 1000$   
 $200 \times 1000$

103. The number of unpaired electrons in  $Fe^{3+}$  is

- a) 2
- b) 3
- c) 4
- d) 5



104. The IUPAC name of  $K_3[Fe(CN)_6]$  is

- a) Potassium ferri cyanide
- b) Potassium ferro cyanide
- c) Potassium Hexa cyano ferrate (II)
- d) Potassium Hexa cyano ferrate (III)

Potassium Hexacyano ferrate(II)

105. The adsorption of an inert gases on activated charcoal increases with

- a) decrease of pressure
- b) increase of temperature
- c) decrease of atomic mass
- d) decrease of temperature

decrease of temperature

106. Electrolysis of brine gives a mixture of

- a)  $H_2, Na, Cl_2$
- b)  $Cl_2, H_2, NaOH$
- c)  $H_2, O_2, NaOH$
- d)  $O_2, Cl_2, NaOH$

107. Sucrose is a non reducing sugar due to

- a) 1-2 linkage
- b) 1-4 linkage
- c) 1-5 linkage
- d) 1-6 linkage

1-2 linkage

108. Sulfur containing amino acid is

- a) alanine
- b) proline
- c) tyrosine
- d) cystein

109. Lysine is

- a) Neutral amino acid
- b) Acidic amino acid
- c) Basic amino acid
- d) Heterocyclic amino acid

Basic amino acid


Space for calculation / rough work

-6 = -4  
2


110. In the Molisch reagent, the substance used is  
 a)  $\beta$  naphthol in alcohol  
 b)  $\alpha$  naphthol in alcohol  
 c) Resorcinol in alcohol  
 d) Rosaniline in water

$\alpha$  naphthol in alcohol

111. In benzene, each carbon atom undergoes  
 a)  $sp$  hybridization  
 b)  $sp^2$  hybridization  
 c)  $sp^3$  hybridization  
 d)  $dsp^2$  hybridization



112. When vapours of isopropyl alcohol is passed over heated copper we get acetone. It is an example for  
 a) dehydration  
 b) dehalogenation  
 c) dehydrohalogenation  
 d) dehydrogenation




113. 
$$\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3-\text{N}-\text{CH}_3 \end{array}$$
 is the IUPAC name of

- a) tri methyl amine  
 b) 2 methyl ethanamine  
 c) N-N dimethyl methanamine  
 d) trimethyl ammonia

N-N dimethyl methanamine



114. When Benzaldehyde is condensed with acetic anhydride in presence of fused sodium acetate we get  
 a) Crotonic acid  
 b) Cinnamic acid  
 c) Aspartic acid  
 d) Salicylic acid



115. When a mixture of Calcium Benzoate & Calcium formate is dry distilled, we get  
 a) Formaldehyde  
 b) Acetaldehyde  
 c) Benzaldehyde  
 d) Salicylaldehyde

Benzaldehyde

Space for calculation / rough work



$71 \times 50 = 25$

$2 = \frac{20 \times 10}{50}$


$0.10 = \frac{2}{10}$



Physics and Chemistry


116. Which one of the following is strongly basic?

- a) Dimethyl amine
- b) Methyl amine
- c) Ammonia
- d) Aniline





117. Which one of the following is bi functional compound?

- a) Formic acid
- b) Acetic acid
- c) Benzoic acid
- d) Cinnamic acid



118. When phenol is treated with Chloro methane in presence of  $\text{AlCl}_3$ , we get

- a) o-cresol
- b) m-cresol
- c) p-cresol
- d) mixture of o & p-cresol



119. In the synthesis of ammonia  $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$

- a)  $K_p = K_c RT$
- b)  $K_p = K_c$
- c)  $K_p = K_c (RT)^{-2}$
- d)  $K_p = K_c (RT)^{-1}$

$$\Delta n = -2$$


$$K_p = K_c (RT)^{-2}$$

120. When the same amount of electricity is passed through solutions of silver nitrate and copper sulfate, 0.4g copper is deposited. The amount of silver deposited is

- a) 1.35g
- b) 2.7g
- c) 5.1g
- d) 5.4g

\*\*\*

Space for calculation / rough work



$$z_p = k A P^{an} S^c$$

$$an = 2 - 4$$

$$= -2$$