

PHYSICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer:

1. A force $F = -40x$ acts on a mass of 1 kg. x is the position of the mass. If maximum speed of the mass is 4 m/s, find the amplitude. All parameters are in SI units.

- (1) $\frac{1}{\sqrt{10}}$ m (2) $\frac{2}{\sqrt{10}}$ m
 (3) $\frac{3}{\sqrt{10}}$ m (4) $\frac{4}{\sqrt{10}}$ m


Answer (2)

Sol. $V_{\max} = A\omega = A\sqrt{\frac{k}{m}}$

$\Rightarrow 4 = A\sqrt{\frac{40}{1}}$


$\Rightarrow A = \frac{2}{\sqrt{10}}$ m

2. Consider 2 inclined plane of same height. 1st has a smooth surface and angle of inclination is 45° , other has a rough surface and angle of inclination is 60° . If ratio of time taken to slide on then its 'n'. Find coefficient of friction of rough inclined plane.



- (1) $\mu = 3n^2$
 (2) $\mu = \frac{3 - 2n^2}{\sqrt{3}}$
 (3) $\mu = \frac{3 - \sqrt{3}n^2}{2}$
 (4) $\mu = \frac{2n^2}{\sqrt{3}}$

Answer (2)



$a = g \sin \theta = \left(\frac{g}{\sqrt{2}}\right)$

$t_1 = \sqrt{\frac{2l_1}{a}}$

$= \sqrt{\frac{2\sqrt{2}h}{\left(\frac{g}{\sqrt{2}}\right)}}$


$= \sqrt{\frac{4h}{g}}$

$\frac{t_1}{t_2} = \sqrt{\frac{3 - \sqrt{3}\mu}{2}} = n$

$\Rightarrow 3 - \sqrt{3}\mu = 2n^2$

$\Rightarrow \sqrt{3}\mu = 3 - 2n^2$

$\Rightarrow \mu = \left(\frac{3 - 2n^2}{\sqrt{3}}\right)$



$a = g \sin \theta - \mu g \cos \theta$

$= \left(g \frac{\sqrt{3}}{2} - \frac{\mu g}{2}\right)$

$= g \left(\frac{\sqrt{3}}{2} - \frac{\mu}{2}\right)$

$t_2 = \sqrt{\frac{2l_2}{a}}$

$= \sqrt{\frac{2 \times 2h}{\sqrt{3}g \left(\frac{\sqrt{3}}{2} - \frac{\mu}{2}\right)}}$

$= \sqrt{\frac{8h}{g(3 - \sqrt{3}\mu)}}$

3. A particle undergoing uniform circular motion about origin. At certain instant $x = 2$ m and $v = -4\hat{j}$ m/s, find velocity and acceleration of particle when at $x = -2$ m.

(1) $\vec{v} = -4\hat{j}$ m/s

$\vec{a} = 8\hat{i}$ m/s²

(2) $\vec{v} = 4\hat{j}$ m/s

$\vec{a} = 8\hat{i}$ m/s²

(3) $\vec{v} = -4\hat{j}$ m/s


$\vec{a} = -8\hat{i}$ m/s²

(4) $\vec{v} = 4\hat{j}$ m/s

$\vec{a} = -8\hat{i}$ m/s²

Answer (2)

Sol. For uniform circular motion,




At $x = -2 \text{ m}$, $v = 4\hat{j}$

$$a = \left(\frac{v^2}{R}\right) \text{ towards the centre}$$

$$a = \left(\frac{4^2}{2}\right) = 8 \text{ m/s}^2$$

$$\vec{a} = 8 \text{ m/s}^2 (\hat{i})$$

4. A man pulls a block as shown:



Consider the following statements:

- (a) Work done by gravity on block is +ve
- (b) Work done by gravity on block is -ve
- (c) If man pulls block with constant speed, then tension in string equals weight of block.
- (d) None of the above

Which of the statement(s) is/are correct?


- (1) (b) & (c) only
- (2) (d) only
- (3) (a) & (c) only
- (4) (a) only

Answer (1)

Sol. Weight acts down and displacement is up \Rightarrow statement (b) is correct.

$$T - mg = ma$$

$$\Rightarrow \text{If } a = 0, T = mg$$



RMS current in circuit (a) is I_a while RMS current in circuit (b) is I_b then

- (1) $I_a > I_b$
- (2) $I_a < I_b$
- (3) $I_a = I_b$
- (4) None of the above

Answer (1)


Sol. $Z_a = 4 \Omega$ & $Z_b = \sqrt{4^2 + (5-3)^2} \Omega = \sqrt{20} \Omega$

$$\Rightarrow I_a = \frac{220}{4} \quad \& \quad I_b = \frac{220}{\sqrt{20}}$$

$$= 55 \text{ A} \quad \quad = \frac{110}{\sqrt{5}}$$

So $I_a > I_b$


6. Find truth table



A	B	X
0	0	1
0	1	1
1	0	1
1	1	0

A	B	X
0	0	1
0	1	0
1	0	0
1	1	1

Answer (4)



Sol.

$$X_1 = \bar{A}$$

$$X_3 = B \cdot \bar{A}$$

$$X_2 = \bar{B}$$

$$X_4 = (A\bar{B})$$

Sol. $E = \left(\frac{P^2}{2m}\right)$

where P is linear momentum, $E =$ kinetic energy, $m =$ mass of particle.

$P = \sqrt{2Em}$

$\lambda = \frac{h}{P} = \frac{h}{\sqrt{2Em}}$

13. In a region with electric field $30\hat{i}$ V/m a charge particle of charge $q = 2 \times 10^{-4}$ C is displaced slowly from (1, 2) to origin. The work done by the external agent is equal to

- (1) 1 mJ (2) 6 mJ
 (3) 2 mJ (4) 3 mJ


Answer (2)

Sol. $F = qE = 2 \times 10^{-4} \times 30$ N

Work done = $6 \times 10^{-3} \times (1)$ J = 6 mJ

14. Consider the following potentiometer circuit :

When switch S is open, length AJ is 300 cm. When switch S is closed, length AJ is 200 cm. If R 5 Ω , find internal resistance r of the cell.



- (1) 4 Ω (2) 2 Ω
 (3) 5 Ω (4) 2.5 Ω

Answer (4)

Sol. $C \times 300 = \varepsilon$... (1)

$C \times 200 = \frac{\varepsilon}{R+r} R$... (2)

$\Rightarrow \frac{300}{200} = \frac{R+r}{R}$

$\Rightarrow r = \frac{R}{2} = 2.5 \Omega$

15.
16.
17.
18.
19.
20.

SECTION - B

Numerical Value Type Questions: This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g. 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

21. At 300 K, RMS speed of an ideal gas molecules is

$\sqrt{\frac{\alpha + 5}{\alpha}}$ times the average speed of gas molecules, then value of α is equal to (take $\pi = \frac{22}{7}$)

Answer (28.00)

Sol. $v_{rms} = \sqrt{\frac{3RT}{M_0}}$

$v_{av} = \sqrt{\frac{8RT}{\pi M_0}}$

$\frac{v_{rms}}{v_{av}} = \sqrt{\frac{3\pi}{8}}$

$= \sqrt{\frac{3 \times 22}{8 \times 7}}$

$= \sqrt{\frac{33}{28}} = \sqrt{\frac{28+5}{28}}$

$\Rightarrow \alpha = 28$

22. An α -particle and a proton are accelerated through same potential difference. The ratio of de-Broglie wavelength of alpha particle to proton is equal to

$\frac{1}{\sqrt{x}}$. Value of x is (take $m_\alpha = 4m_{proton}$)

Answer (08.00)

