


**PHYSICS**

**SECTION - A**

**Multiple Choice Questions:** This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

**Choose the correct answer:**

1. Bob *P* is released from the position of rest at the moment shown. If it collides elastically with an identical bob *Q* hanging freely then velocity of *Q*, just after collision is ( $g = 10 \text{ m/s}^2$ )



- (1) 1 m/s                      (2) 4 m/s  
(3) 2 m/s                      (4) 8 m/s

**Answer (3)**

**Sol.** Velocity of *P* just before collision is  $= \sqrt{2gl}$   
 $= 2 \text{ m/sec}$

As collision is elastic and the mass of *P* and *Q* are equal therefore just after collision velocity of *P* is 0 and that of *Q* is 2 m/sec.

2. Choose the option showing the correct relation between Poisson's ratio ( $\sigma$ ), Bulk modulus (*B*) and modulus of rigidity (*G*).

- (1)  $\sigma = \frac{3B - 2G}{2G + 6B}$                       (2)  $\sigma = \frac{6B + 2G}{3B - 2G}$   
(3)  $\sigma = \frac{9BG}{3B + G}$                       (4)  $B = \frac{3\sigma - 3G}{6\sigma + 2G}$

**Answer (1)**

**Sol.**  $E = 2G(1 + \sigma)$                       ....(1)

$E = 3B(1 - 2\sigma)$                       ....(2)


$$1 = \frac{2G}{3B} \left( \frac{1 + \sigma}{1 - 2\sigma} \right)$$

$$\Rightarrow 3B - 6B\sigma = 2G + 2G\sigma$$

$$\Rightarrow 3B - 2G = \sigma (2G + 6B)$$

$$\sigma = \left( \frac{3B - 2G}{2G + 6B} \right)$$

3. Two conducting solid spheres (*A* & *B*) are placed at a very large distance with charge densities and radii as shown:



When the key *K* is closed, find the ratio of final charge densities.

- (1) 4 : 1                      (2) 1 : 2  
(3) 2 : 1                      (4) 1 : 4

**Answer (3)**


**Sol.** Final potential is same

$$\Rightarrow \frac{1}{4\pi\epsilon_0} \frac{Q_1}{R} = \frac{1}{4\pi\epsilon_0} \frac{Q_2}{2R} \quad \dots(1)$$


$$\text{Also, } Q_1 + Q_2 = \sigma \cdot 4\pi R^2 + \sigma \cdot 4\pi(2R)^2 \quad \dots(2)$$

$$\Rightarrow \frac{\sigma_1}{\sigma_2} = 2.$$

4. Position-time graph for a particle is parabolic and is as shown:



Choose the corresponding *v* - *t* graph



**Answer (2)**

**Sol.** Since  $x \propto t^2$

$$\Rightarrow v = \frac{dx}{dt} \propto t'$$

$\Rightarrow$  Option 2 is correct

5. For a system undergoing isothermal process, heat energy is supplied to the system. Choose the option showing correct statements
- (a) Internal energy will increase
  - (b) Internal energy will decrease
  - (c) Work done by system is positive
  - (d) Work done by system is negative
  - (e) Internal energy remains constant
- (1) (a), (c), (e)                      (2) (b), (d)  
 (3) (c), (e)                            (4) (a), (d), (e)

**Answer (3)**

**Sol.** For isothermal process,

$$dT = 0$$

so,  $dU = 0 \Rightarrow$  Internal energy remains same

$$dQ = dW$$

as  $dQ$  is positive,

so  $dW$  is positive

6. The heat passing through the cross-section of a conductor, varies with time ' $t$ ' as  $Q(t) = \alpha t - \beta t^2 + \gamma t^3$ . ( $\alpha$ ,  $\beta$  and  $\gamma$  are positive constants.) The minimum heat current through the conductor is

- (1)  $\alpha - \frac{\beta^2}{2\gamma}$                       (2)  $\alpha - \frac{\beta^2}{3\gamma}$   
 (3)  $\alpha - \frac{\beta^2}{\gamma}$                         (4)  $\alpha - \frac{3\beta^2}{\gamma}$

**Answer (2)**

**Sol.** Heat through cross section of rod

$$Q = \alpha t - \beta t^2 + \gamma t^3$$

$$\text{so heat current} = \frac{dQ}{dt}$$

$$\text{heat current} = \frac{dQ}{dt} = \alpha - 2\beta t + 3\gamma t^2$$

for heat current to be minimum

$$\frac{d^2Q}{dt^2} = -2\beta + 6\gamma t = 0$$

$$t = \frac{2\beta}{6\gamma} = \left( \frac{\beta}{3\gamma} \right)$$


so minimum heat current

$$\left. \frac{dQ}{dt} \right|_{\text{minimum}} = \alpha - 2\beta \times \frac{\beta}{3\gamma} + 3\gamma \times \frac{\beta^2}{9\gamma^2}$$

$$= \alpha - \frac{2\beta^2}{3\gamma} + \frac{\beta^2}{3\gamma}$$

$$= \left( \alpha - \frac{\beta^2}{3\gamma} \right)$$


7. Momentum-time graph of an object moving along a straight line is as shown in figure. If  $(P_2 - P_1) < P_1$  and  $(t_2 - t_1) = t_1 < (t_3 - t_2)$  then at which points among A, B and C the magnitude of force experienced by the object is maximum and minimum respectively.



- (1) A, B                                      (2) A, C  
 (3) B, C                                      (4) B, A

**Answer (2)**

**Sol.**



$$F_A = \frac{P_1}{t_1}$$

$$F_B = \frac{P_2 - P_1}{t_2 - t_1}$$

$$F_C = \frac{P_2 - P_1}{t_3 - t_2}$$


Therefore the maximum force is at A and minimum force is at C.

8. A particle moving in unidirectional motion travels half of the total distance with a constant speed of 15 m/s. Now first half of the journey time it travels at 10 m/s and second half of the remaining journey time it travels at 5 m/s. Average speed of the particle is

- (1) 12 m/s                                      (2) 10 m/s  
 (3) 7 m/s                                        (4) 9 m/s

**Answer (2)**

**Sol.**




$$v_{av} = \frac{2x}{\frac{x}{15} + 2t}$$

$$= \frac{2x}{\frac{x}{15} + \frac{2x}{10+5}}$$

$$= 10 \text{ m/sec}$$

9. A bullet strikes a stationary ball kept at a height as shown. After collision, range of bullet is 120 m and that of ball is 30 m. Find initial speed of bullet. Collision is along horizontal direction.

Take  $g = 10 \text{ m/s}^2$



- (1) 150 m/s                      (2) 90 m/s  
(3) 240 m/s                      (4) 360 m/s

**Answer (4)**

**Sol.**  $m_1v + m_2(0) = m_1v_1' + m_2v_2'$  ... (1)

$$\Delta t = \sqrt{\frac{2h}{g}} = 2s \quad \dots(2)$$

$$\Rightarrow v_1' = \frac{120 \text{ m}}{2s} = 60 \text{ m/s}$$


$$\& v_2' = \frac{30 \text{ m}}{2s} = 15 \text{ m/s}$$

$$\Rightarrow v = 360 \text{ m/s}$$

10. If an inductor with inductive reactance,  $X_L = R$  is connected in series with resistor  $R$  across an A.C voltage, power factor comes out to be  $P_1$ . Now, if a capacitor with capacitive reactance,  $X_C = R$  is also connected in series with inductor and resistor in the same circuit, power factor becomes  $P_2$ . Find  $\frac{P_1}{P_2}$

- (1)  $\sqrt{2} : 1$                       (2)  $1 : \sqrt{2}$   
(3)  $1 : 1$                       (4)  $1 : 2$

**Answer (2)**




$$Z = \sqrt{R^2 + R^2}$$

$$= \sqrt{2}R$$

$$P_1 = \cos\phi = \text{power factor} = \frac{R}{Z} = \left(\frac{1}{\sqrt{2}}\right)$$

When capacitor is also connected in series



The LCR circuit is in resonance stage

$$\text{So, } Z = \sqrt{R^2 + (X_L - X_C)^2}$$

$$Z = R$$

$$P_2 = \cos\phi = \text{power factor} = \frac{R}{Z} = \frac{R}{R} = 1$$

$$\text{So, } \frac{P_1}{P_2} = \frac{\left(\frac{1}{\sqrt{2}}\right)}{1} = \frac{1}{\sqrt{2}}$$

11. Electromagnetic wave beam of power 20 mW is incident on a perfectly absorbing body for 300 ns. The total momentum transferred by the beam to the body is equal to

- (1)  $2 \times 10^{-17} \text{ Ns}$                       (2)  $1 \times 10^{-17} \text{ Ns}$   
(3)  $3 \times 10^{-17} \text{ Ns}$                       (4)  $5 \times 10^{-17} \text{ Ns}$

**Answer (1)**

**Sol.** Total energy incident =  $Pt$

$$\text{So total initial momentum} = \frac{Pt}{c}$$

$$\text{Total final momentum} = 0$$

$$\text{Total momentum transferred} = \frac{Pt}{c}$$

$$= \frac{20 \times 10^{-3} \times 300 \times 10^{-9}}{3 \times 10^8}$$

$$= 2 \times 10^{-17} \text{ Ns}$$

12. The velocity of an electron in the seventh orbit of hydrogen-like atom is  $3.6 \times 10^6 \text{ m/s}$ . Find the velocity of the electron in the 3<sup>rd</sup> orbit.

- (1)  $4.2 \times 10^6 \text{ m/s}$                       (2)  $8.4 \times 10^6 \text{ m/s}$   
(3)  $2.1 \times 10^6 \text{ m/s}$                       (4)  $3.6 \times 10^6 \text{ m/s}$

**Answer (2)**

**Sol.** For hydrogen like atom,

$$v \propto \frac{1}{n}$$

$$\left(\frac{v_1}{v_2}\right) = \left(\frac{n_2}{n_1}\right)$$

$$\Rightarrow \frac{3.6 \times 10^6}{v_2} = \frac{3}{7}$$

$$\Rightarrow v_2 = \frac{7}{3} \times 3.6 \times 10^6$$

$$= 8.4 \times 10^6 \text{ m/s}$$

13. Electric field in a region is given by  $\vec{E} = \frac{a}{x^2} \hat{i} + \frac{b}{y^3} \hat{j}$ ,

where  $x$  &  $y$  are co-ordinates. Find SI units of  $a$  &  $b$ .

- (1)  $a - \text{Nm}^2\text{C}^{-1}$                       (2)  $a - \text{Nm}^3\text{C}^{-1}$   
 $b - \text{Nm}^3\text{C}^{-1}$                        $b - \text{Nm}^2\text{C}^{-1}$   
 (3)  $a - \text{NmC}^{-1}$                       (4)  $a - \text{Nm}^2\text{C}^{-1}$   
 $b - \text{Nm}^2\text{C}^{-1}$                        $b - \text{Nm}^2\text{C}^{-1}$

**Answer (1)**

**Sol.**  $E - \text{NC}^{-1}$

$$x^2 - \text{m}^2$$

$$y^3 - \text{m}^3$$

$$\Rightarrow a - \text{Nm}^2\text{C}^{-1}$$

$$\& \quad b - \text{Nm}^3\text{C}^{-1}$$

14. Coil A of radius 10 cm has  $N_A$  number of turns and  $I_A$  current is flowing through it. Coil B of radius 20 cm has  $N_B$  number of turns and  $I_B$  current is flowing through it. If magnetic dipole moment of both the coils is same then

- (1)  $I_A N_A = 4 I_B N_B$                       (2)  $I_A N_A = \frac{1}{4} I_B N_B$   
 (3)  $I_A N_A = 2 I_B N_B$                       (4)  $I_A N_A = \frac{1}{2} I_B N_B$

**Answer (1)**

**Sol.** Magnetic dipole moment  $\mu = NIA = NI\pi R^2$

$$\text{So } \frac{\mu_A}{\mu_B} = \frac{N_A I_A R_A^2}{N_B I_B R_B^2} = 1$$

$$\frac{N_A I_A (10^2)}{N_B I_B (20^2)} = 1$$

$$N_A I_A = 4 N_B I_B$$

15. An ideal gas undergoes a thermodynamic process following the relation  $PT^2 = \text{constant}$ . Assuming symbols have their usual meaning then volume expansion coefficient of the gas is equal to

- (1)  $\frac{2}{T}$                                       (2)  $\frac{3}{T}$   
 (3)  $\frac{1}{2T}$                                       (4)  $\frac{1}{T}$

**Answer (2)**

**Sol.** Volume expansion coefficient  $= \frac{dV}{VdT}$

For  $PT^2 = \text{constant}$




$$\text{Or } \frac{T^3}{V} = \text{constant}$$

$$\text{Or } \frac{dV}{dT} = (C) 3T^2$$

$$\text{Or } \frac{dV}{VdT} = \frac{3T^2}{T^3}$$

$$\frac{dV}{VdT} = \frac{3}{T}$$

16. Consider a combination of gates as shown :



**Answer (1)**

**Sol.**  $y = (A'B') = A + B$

$\Rightarrow$  OR gate


$\Rightarrow$  Option 1

17. For the given YDSE setup. Find the number of fringes by which the central maxima gets shifted from point O.

(Given  $d = 1 \text{ mm}$ )

$D = 1 \text{ m}$


$\lambda = 5000 \text{ \AA}$ )



- (1) 10                                      (2) 15  
 (3) 8                                      (4) 12

**Answer (1)**

Sol.



at central position, path difference, is,

$$(\mu - 1)t_1 - (\mu - 1)t_2$$

$$\Delta x = (\mu - 1)(t_1 - t_2)$$

$$\Delta x = \left(\frac{3}{2} - 1\right)(5.11 - 5.10) \text{ mm}$$

$$= \frac{1}{2} \times (0.01) \text{ mm}$$

$$= 0.005 \text{ mm}$$

$$= 5 \times 10^{-6} \text{ m}$$

$$\text{No. of fringes shifted} = \frac{\Delta x}{\lambda} = \frac{5 \times 10^{-6} \text{ m}}{5 \times 10^{-7} \text{ m}}$$

$$= 10$$

18.


19.

20.

**SECTION - B**

**Numerical Value Type Questions:** This section contains 10 questions. In Section B, attempt any five questions out of 10. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value (in decimal notation, truncated/rounded-off to the second decimal place; e.g. 06.25, 07.00, -00.33, -00.30, 30.27, -27.30) using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.

21. In a part of a circuit shown:



Find  $V_A - V_B$  in volts. It is given that current is decreasing at a rate of 1 ampere/s.

**Answer (18)**

Sol.  $V_A - iR - L \frac{di}{dt} - 12 = V_B$

$$\Rightarrow V_A - V_B = +18 \text{ volts}$$

22. A particle undergoing SHM follows the position-time equation given as  $x = A \sin\left(\omega t + \frac{\pi}{3}\right)$ . If the SHM motion has a time period of  $T$ , then velocity will be maximum at time  $t = \frac{T}{\beta}$  for first time after  $t = 0$ . Value of  $\beta$  is equal to

**Answer (03.00)**

Sol.  $x = A \sin\left(\omega t + \frac{\pi}{3}\right)$

$$\Rightarrow v = A\omega \cos\left(\omega t + \frac{\pi}{3}\right)$$

For maximum value of  $v$

$$\cos\left(\omega t + \frac{\pi}{3}\right) = \pm 1$$


$$\Rightarrow \omega t + \frac{\pi}{3} = \pi \text{ (for nearest value of } t)$$

$$\omega t = \frac{2\pi}{3}$$


$$t = \frac{T}{3}$$

So  $\beta = 3$

23. A block of mass 1 g is in equilibrium with the help of a current carrying square loop which is partially lying in constant magnetic field ( $B$ ) as shown. Resistance of the loop is  $10 \Omega$ . Find the voltage ( $V$ ) (in volts) of the battery in the loop.



**Answer (10.00)**



$$ilB = mg$$

$$i = \left( \frac{mg}{lB} \right) = \frac{(1 \times 10^{-3} \text{ kg}) \times (10 \text{ m/s}^2)}{(0.1 \text{ m}) \times (0.1 \text{ T})}$$

$$= 1 \times 10^{-3} \times 10^3$$

$$i = 1 \text{ A}$$

As resistance of loop =  $10 \Omega$

$$i = \frac{V}{R} = 1 \text{ A}$$

$$V = (1 \times 10) \text{ V} = 10 \text{ V}$$

24. Initial volume of 1 mole of a monoatomic gas is 2 litres. It is expanded isothermally to a volume of 6 litres. Change in internal energy is  $xR$ . Find  $x$ .

**Answer (00)**

Sol.  $\Delta U = nC_V \Delta T$

$$= nC_V(0) \quad (\because \text{isothermal})$$

$$\Rightarrow \Delta U = 0$$

25. An object is placed at a distance of 40 cm from the pole of a converging mirror. The image is formed at a distance of 120 cm from the mirror on the same side. If the focal length is measured with a scale where each 1 cm has 20 equal divisions. If the fractional error in the measurement of focal length

is  $\frac{1}{10k}$ . Find  $k$ .

**Answer (60.00)**

Sol.  $u = -40 \text{ cm}$

$$v = -120 \text{ cm}$$

$$\frac{1}{v} + \frac{1}{u} = \frac{1}{f}$$

$$\Rightarrow -\frac{1}{120} - \frac{1}{40} = \frac{1}{f}$$

$$\frac{1}{f} = \left( \frac{-1-3}{120} \right) = -\frac{4}{120}$$


$$f = -30 \text{ cm}$$

$$\text{Least count of scale} = \left( \frac{1}{20} \right) \text{ cm}$$

$$\text{Fractional error} = \left( \frac{1}{\frac{20}{30}} \right) = \left( \frac{1}{600} \right)$$


$$\text{as } \frac{1}{10k} = \frac{1}{600}$$

$$k = 60$$



In two circuit shown above the value of current  $I_1$  (in amperes) is equal to  $-\frac{y}{5}$  A. Value of  $y$  is equal to

**Answer (11.00)**



Using Kirchoff's law.

$$I_1 + I_3 - I_2 = -2 \quad \dots(i)$$

$$I_3 + 2I_2 = 5 \quad \dots(ii)$$

$$2I_2 - (I_3 - I_2) - (I_1 + I_3 - I_2) = 5 \quad \dots(iii)$$

$$\Rightarrow I_1 = -\frac{11}{5} \text{ A}$$

$$\Rightarrow y = 11$$

27. ??

28. ??

29. ??

30. ??