

3. If $A = \begin{bmatrix} \sqrt{2} & 1 \\ -1 & \sqrt{2} \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$, $C = ABA^T$ and $X = AC^2A^T$, then $|X|$ is equal to

- (1) 729 (2) 283
 (3) 27 (4) 23

Answer (1)

Sol. $|A| = 3$

$$|B| = 1$$

$$\Rightarrow |C| = |ABA^T| = |A||B|A^T = |A|^2|B| \\ = 9$$

$$\Rightarrow |X| = |A||C|^2|A^T|$$

$$= 3 \times 9^2 \times 3 = 9 \times 9^2 = 729$$

4. If 3, 7, 11, ..., 403 = AP_1
 2, 5, 8, ..., 401 = AP_2

Find sum of common term of AP_1 and AP_2

- (1) 3366 (2) 6699
 (3) 9999 (4) 6666

Answer (2)

Sol. 3, 7, 11, 15, 19, 23, 27, ... 403 = AP_1

2, 5, 8, 11, 14, 17, 20, 23, ... 401 = AP_2

so common terms A.P.

11, 23, 35, ..., 395

$$\Rightarrow 395 = 11 + (n-1) 12$$

$$\Rightarrow 395 - 11 = 12(n-1)$$

$$\frac{384}{12} = n-1$$

$$32 = n-1$$

$$n = 33$$

$$\text{Sum} = \frac{33}{2}[2 \times 11 + (32)12]$$

$$= \frac{33}{2}[22 + 384]$$

$$= 6699$$

5. $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{8\sqrt{2}\cos x}{(1+e^{\sin x})(1+\sin^4 x)} dx = a\pi + b\log(3+2\sqrt{2})$

then find $a + b$.

- (1) 4 (2) 6
 (3) 8 (4) 2

Answer (1)

Sol. $I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{8\sqrt{2}\cos x}{(1+e^{\sin x})(1+\sin^4 x)} dx$

$$= \int_0^{\frac{\pi}{2}} \left\{ \frac{8\sqrt{2}\cos x}{(1+e^{\sin x})(1+\sin^4 x)} + \frac{8\sqrt{2}\cos x}{(1+e^{-\sin x})(1+\sin^4 x)} \right\} dx \\ = 8\sqrt{2} \int_0^{\frac{\pi}{2}} \frac{\cos x}{1+\sin^4 x} dx$$

Let $\sin x = t$

$$I = 8\sqrt{2} \int_0^1 \frac{dt}{1+t^4} \\ = 4\sqrt{2} \int_0^1 \frac{\left(1+\frac{1}{t^2}\right) - \left(1-\frac{1}{t^2}\right)}{t^2 + \frac{1}{t^2}} dt \\ = 4\sqrt{2} \int_0^1 \frac{\left(1+\frac{1}{t^2}\right) dt}{t^2 + \frac{1}{t^2} + 2} - 4\sqrt{2} \int_0^1 \frac{\left(1-\frac{1}{t^2}\right) dt}{t^2 + \frac{1}{t^2} - 2} \\ = 4\sqrt{2} \cdot \frac{1}{\sqrt{2}} \left[\tan^{-1} \frac{t-1}{\sqrt{2}} \right]_0^1 - 4\sqrt{2} \cdot \frac{1}{2\sqrt{2}} \left[\log \left| \frac{t+\frac{1}{t}-\sqrt{2}}{t+\frac{1}{t}+\sqrt{2}} \right| \right]_0^1 \\ = 2\pi - 2 \log \left| \frac{2-\sqrt{2}}{2+\sqrt{2}} \right| \\ = 2\pi + 2 \log(3+2\sqrt{2})$$

$$\therefore a = b = 2$$

6. If $(t+1)dx = (2x + (t+1)^3)dt$ and $x(0) = 2$, then $x(1)$ is equal to

- (1) 5 (2) 12
 (3) 6 (4) 8

Answer (2)

Sol. $(t+1)dx = (2x + (t+1)^3)dt$

$$\therefore \frac{dx}{dt} - \frac{2x}{t+1} = (t+1)^2$$

$$\therefore \text{I.F.} = e^{\int -\frac{2}{t+1} dt} = \frac{1}{(t+1)^2}$$

\therefore Solution is

$$\frac{x}{(t+1)^2} = \int 1 dt$$

$$x = (t+c)(t+1)^2$$

$$\because x(0) = 2 \text{ then } c = 2$$

$$\therefore x = (t+2)(t+1)^2$$

$$\therefore x(1) = 12$$

7. Five people are distributed in four identical rooms. A room can also contain zero people. Find the number of ways to distribute them.

(1) 47

(2) 53

(3) 43

(4) 51

Answer (4)

Sol. Total ways to partition 5 into 4 parts are:

$$5 \ 0 \ 0 \ 0 \rightarrow 1$$

$$4 \ 1 \ 0 \ 0 \rightarrow \frac{5!}{4!} = 5$$

$$3 \ 2 \ 0 \ 0 \rightarrow \frac{5!}{3! \cdot 2!} = 10$$

$$3 \ 1 \ 1 \ 0 \rightarrow \frac{5!}{3! \cdot 2!} = 10$$

$$2 \ 2 \ 1 \ 0 \rightarrow \frac{5!}{2! \cdot 2! \cdot 2!} = 15$$

$$2 \ 1 \ 1 \ 1 \rightarrow \frac{5!}{2! \times 3!} = 10$$

51 → Total way

8. $5f(x) + 4f\left(\frac{1}{x}\right) = x^2 - 4$ and $y = 9f(x) \cdot x^2$. If y is strictly increasing function, find interval of x .

$$(1) \left(-\infty, -\frac{1}{\sqrt{5}}\right] \cup \left(-\frac{1}{\sqrt{5}}, 0\right)$$

$$(2) \left(\frac{-1}{\sqrt{5}}, 0\right) \cup \left(0, \frac{1}{\sqrt{5}}\right)$$

$$(3) \left(0, \frac{1}{\sqrt{5}}\right) \cup \left(\frac{1}{\sqrt{5}}, \infty\right)$$

$$(4) \left(-\sqrt{\frac{2}{5}}, 0\right) \cup \left(\sqrt{\frac{2}{5}}, \infty\right)$$

Answer (4)

Sol. $5f(x) + 4f\left(\frac{1}{x}\right) = x^2 - 4 \quad \dots(1)$

Replace x by $\frac{1}{x}$

$$5f\left(\frac{1}{x}\right) + 4f(x) = \frac{1}{x^2} - 4 \quad \dots(2)$$

$$5 \times \text{equation (1)} - 4 \times \text{equation (2)}$$


$$9f(x) = 5x^2 - \frac{4}{x^2} - 4$$

$$y = 9f(x) \cdot x^2 = \frac{5x^4 - 4 - 4x^2}{x^2} x^2$$

$$y = 5x^4 - 4 - 4x^2$$

$$y' = 20x^3 - 8x > 0$$

$$4x(5x^2 - 2) > 0$$



$$x \in \left(-\sqrt{\frac{2}{5}}, 0\right) \cup \left(\sqrt{\frac{2}{5}}, \infty\right)$$

9. If hyperbola $x^2 - y^2 \operatorname{cosec}^2 \theta = 5$ and ellipse $x^2 \operatorname{cosec}^2 \theta + y^2 = 5$ has eccentricity e_H and e_e respectively and $e_H = \sqrt{7}e_e$, then θ is equal to

$$(1) \frac{\pi}{3}$$

$$(2) \frac{\pi}{6}$$

$$(3) \frac{\pi}{2}$$

$$(4) \frac{\pi}{4}$$

Answer (1)

Sol. $x^2 - y^2 \operatorname{cosec}^2 \theta = 5 \Rightarrow \frac{x^2}{1} - \frac{y^2}{\sin^2 \theta} = 5$

$$x^2 \operatorname{cosec}^2 \theta + y^2 = 5 \Rightarrow \frac{x^2}{\sin^2 \theta} + \frac{y^2}{1} = 5$$

$$e_H = \sqrt{7}e_e$$

$$e_H = \sqrt{1 + \frac{\sin^2 \theta}{1}}$$

$$\text{and } e_e = \sqrt{1 - \frac{\sin^2 \theta}{1}}$$

$$\Rightarrow \sqrt{1 + \sin^2 \theta} = \sqrt{7} \sqrt{1 - \sin^2 \theta}$$

$$\Rightarrow 1 + \sin^2 \theta = 7 - 7 \sin^2 \theta$$

$$\Rightarrow 8 \sin^2 \theta = 6$$

$$\Rightarrow \sin \theta = \sqrt{\frac{3}{4}} = \frac{\sqrt{3}}{2}$$

$$\Rightarrow \theta = \frac{\pi}{3}$$

13. $f(x) = \begin{cases} e^{-x}, & x < 0 \\ \ln x, & x > 0 \end{cases}$


$$g(x) = \begin{cases} e^x, & x < 0 \\ x, & x > 0 \end{cases}$$

The $gof: A \rightarrow R$ is

- (1) Onto but not one-one
- (2) Into and many one
- (3) Onto and one-one
- (4) Into and one-one


Answer (2)

Sol.



$$gof(x) = \begin{cases} f(x), & f(x) < 0 \\ f(x), & f(x) > 0 \end{cases}$$

$$= \begin{cases} e^{\ln x} = x, & (0, 1) \\ e^{-x}, & (-\infty, 0) \\ \ln x, & (1, \infty) \end{cases}$$



$\therefore gof(x)$ is many one and into

14. If $\tan A = \frac{1}{\sqrt{x^2 + x + 1}}$, $\tan B = \frac{\sqrt{x}}{\sqrt{x^2 + x + 1}}$ and

$$\tan C = \frac{1}{\sqrt{x(x^2 + x + 1)}}, \text{ then } A + B =$$

- (1) 0
- (2) $\pi - C$
- (3) $\frac{\pi}{2} - C$
- (4) None

Answer (3)

Sol. $\tan B \times \tan C = \frac{\sqrt{x}}{\sqrt{x^2 + x + 1}} \times \frac{1}{\sqrt{x(x^2 + x + 1)}}$

$$= \frac{1}{x^2 + x + 1} = \tan^2 A$$

$$\tan^2 A = \tan B \tan C$$

It is only possible when $A = B = C$ at $x = 1$

$$\Rightarrow A = 30^\circ, B = 30^\circ, C = 30^\circ$$

$$\left[\tan A = \tan B = \tan C = \frac{1}{\sqrt{3}} \right]$$

$$\therefore A + B = \frac{\pi}{2} - C$$

15. $\lim_{x \rightarrow 0} \frac{\cos^{-1}(1 - \{x\}^2) \sin^{-1}(1 - \{x\})}{\{x\} - \{x\}^3}$, where $\{ \}$ is fractional part function.

If L.H.L = L and R.H.L = R , then the correct relation between L and R is

- (1) $\sqrt{2}R = 4L$
- (2) $\sqrt{2}L = 4R$
- (3) $R = L$
- (4) $R = 2L$

Answer (1)

Sol. $RHL \Rightarrow \lim_{x \rightarrow 0^+} \frac{\cos^{-1}(1 - x^2) \sin^{-1}(1 - x)}{x - x^3}$

$$\Rightarrow \lim_{x \rightarrow 0^+} \frac{\pi}{2} \cdot \frac{\cos^{-1}(1 - x^2)}{x}$$

$$\frac{\pi}{2} \lim_{x \rightarrow 0^+} \frac{-1}{\sqrt{(1 - (1 - x^2))^2}} (-2x)$$

$$= \frac{\pi}{2} \lim_{x \rightarrow 0^+} \frac{2x}{\sqrt{2x^2 - x^4}} = \pi \lim_{x \rightarrow 0^+} \frac{x}{x\sqrt{2 - x^2}}$$

$$= \frac{\pi}{\sqrt{2}}$$

$LHL \Rightarrow \lim_{x \rightarrow 0^-} \frac{\cos^{-1}(1 - (1+x)^2) \sin^{-1}(1 - (1+x))}{1 \cdot (1 - (1+x)^2)}$

$$= \lim_{x \rightarrow 0^-} \frac{\cos^{-1}(-x^2 - 2x) \cdot \sin^{-1}(-x)}{-x^2 - 2x}$$

$$= \frac{\pi}{2} \lim_{x \rightarrow 0^-} \frac{-\sin^{-1} x}{-x(x+2)} = \frac{\pi}{2} \times \frac{1}{2} = \frac{\pi}{4}$$

16.

17.

18.

19.

20.

SECTION - B

Numerical Value Type Questions: This section contains 10 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

21. Let $S = \{1, 2, 3, \dots, 20\}$

$$R_1 = \{(a, b) : a \text{ divide } b\}$$

$$R_2 = \{(a, b) : a \text{ is integral multiple of } b\} \text{ } a, b \in S$$

$$n(R_1 - R_2) = ?$$

Answer (46)

Sol. $R_1 = \{(1, 1), (1, 2), (1, 3), \dots, (1, 20), (2, 2), (2, 4), \dots, (2, 20), (3, 3), (3, 6), \dots, (3, 18), (4, 4), (4, 8), \dots, (4, 20), (5, 5), (5, 10), (5, 15), (5, 20), (6, 6), (6, 12), (6, 18), (7, 7), (7, 14), (8, 8), (8, 16), (9, 9), (9, 18), (10, 10), (10, 20), (11, 11), (12, 12), \dots, (20, 20)\}$

$n(R_1) = 66$

$R_2 = \{a \text{ is integral multiple of } b\}$

$\text{So } n(R_1 - R_2) = 66 - 20 = 46$

as $R_1 \cap R_2 = \{(a, a) : a \in S\} = \{(1, 1), (2, 2), \dots, (20, 20)\}$

22. The number of solution of equation $x + 2y + 3z = 42$ and $x, y, z \in \mathbb{Z}$ and $x, y, z \geq 0$ is

Answer (168)

Sol. $x + 2y + 3z = 42$

0 $x + 2y = 42 \Rightarrow 22 \text{ cases}$

1 $x + 2y = 39 \Rightarrow 19 \text{ cases}$

2 $x + 2y = 36 \Rightarrow 19 \text{ cases}$

3 $x + 2y = 33 \Rightarrow 17 \text{ cases}$

4 $x + 2y = 30 \Rightarrow 16 \text{ cases}$

5 $x + 2y = 27 \Rightarrow 14 \text{ cases}$

6 $x + 2y = 24 \Rightarrow 13 \text{ cases}$

7 $x + 2y = 21 \Rightarrow 11 \text{ cases}$

8 $x + 2y = 18 \Rightarrow 10 \text{ cases}$

9 $x + 2y = 15 \Rightarrow 8 \text{ cases}$

10 $x + 2y = 12 \Rightarrow 7 \text{ cases}$

11 $x + 2y = 9 \Rightarrow 5 \text{ cases}$

12 $x + 2y = 6 \Rightarrow 4 \text{ cases}$

13 $x + 2y = 3 \Rightarrow 2 \text{ cases}$

14 $x + 2y = 0 \Rightarrow 1 \text{ cases}$

23.

24.

25.

26.

27.

28.

29.

30.