

1. A sequence of real numbers  $\{a_n\}_{n \geq 1}$  has a *peak* at  $n$  if  $a_n \geq a_k$  for all  $k \geq n$ . Consider the following statements.

- (I) No sequence of real numbers can have only finitely many peaks.
- (II) No sequence of real numbers can have infinitely many peaks.
- (III) Any sequence of real numbers having finitely many peaks must have the property that  $a_n \geq 0$  for all  $n$  greater than some  $k$ .

Then

- (A) only (I) is true
- (B) none of (I), (II) and (III) are true
- (C) both (II) and (III) are true
- (D) only (III) is true

2. Let  $\mathbb{C}$  denote the set of complex numbers and let  $\text{Im}(z)$  denote the imaginary part of  $z \in \mathbb{C}$ . Consider the set

$$S = \{s \in \mathbb{R} : \text{there exists } z \in \mathbb{C} \text{ such that } \text{Im}(z) \neq 0 \text{ and } s = z^2 + 2z - 1\}.$$

Then,

- (A)  $S \neq \mathbb{R}$ , but contains infinitely many elements
- (B)  $S$  is a non-empty finite set
- (C)  $S = \mathbb{R}$
- (D)  $S$  is the empty set

3. For a set  $S$ , let  $S^c$  denote the complement of  $S$ . Also, for two sets  $P$  and  $Q$ , let  $P \setminus Q = P \cap Q^c$ . Let  $A$ ,  $B_1$ ,  $B_2$  and  $B_3$  be four sets. Which of the following statements is NOT true?

- (A)  $(A \cup B_1 \cup B_2 \cup B_3)^c = A^c \cap B_1^c \cap B_2^c \cap B_3^c$   
(B)  $(A \setminus B_1) \setminus (B_2 \cup B_3) = A \setminus (B_1 \cup B_2 \cup B_3)$   
(C)  $A \setminus (B_1 \cup B_2 \cup B_3) = (A \setminus B_1) \cup (A \setminus B_2) \cup (A \setminus B_3)$   
(D)  $A \cap (B_1 \cup B_2 \cup B_3) = (A \cap B_1) \cup (A \cap B_2) \cup (A \cap B_3)$

4. For a complex number  $z$ , let  $\bar{z}$  be its complex conjugate. Then the equation

$$z\bar{z}^2 + z^2\bar{z} = 0$$

has

- (A) exactly three roots  
(B) exactly two roots  
(C) infinitely many roots  
(D) only real roots
5. Let  $f(x) = x^2 + (2a + 1)x + (a^2 + 2)$ . The number of values of  $a$  for which one of the roots of the equation  $f(x) = 0$  is twice the other root is

- (A) more than 2                      (B) 2                      (C) 1                      (D) 0

6. Let  $A$  be a finite set of real numbers having  $m$  ( $\geq 2$ ) elements. Define a function  $f : \mathbb{R} \rightarrow \mathbb{R}$ , given by

$$f(x) = \min\{|a - x| : a \in A\}.$$

Then,

- (A)  $f$  is continuous everywhere
  - (B)  $f$  is continuous only at finitely many points
  - (C)  $f$  is discontinuous everywhere
  - (D)  $f$  has  $m$  discontinuities
7. Let  $A$  be the set of functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  for which  $|f(x) - f(y)| \leq 2|x - y|^2$  for all  $x, y \in \mathbb{R}$  and  $f(0) = 0$ . Then, for any  $f \in A$ ,
- (A) the functions  $g(x) = P(f(x)) \in A$  for every polynomial  $P$
  - (B) the function  $g(x) = x + f(x) \in A$
  - (C) the function  $g(x) = xf(x) \in A$
  - (D) the function  $g(x) = e^{f(x)} \in A$

8. The number of values of  $a$  for which the three lines

$$2x + y - 1 = 0, \quad ax + 3y - 3 = 0, \quad 3x + 2y - 2 = 0$$

are concurrent is

- (A) more than 2                      (B) 1                      (C) 0                      (D) 2

9. For a non-constant geometric progression for which the second term is 2 and the common ratio is an integer, the 10th, 20th and 30th terms are in arithmetic progression. Then, the fourth term is

- (A)  $-2$                       (B)  $-4$                       (C)  $4$                       (D)  $2$

10.  $\lim_{x \rightarrow 1} \frac{\sqrt{x+8} - \sqrt{8x+1}}{\sqrt{5-x} - \sqrt{7x-3}}$  equals

- (A) does not exist                      (B)  $\frac{2}{3}$                       (C)  $\frac{1}{2}$                       (D)  $\frac{7}{12}$

11. The rank of the matrix

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 1 & 0 & a & 0 \\ 1 & 0 & 0 & 0 & b \end{pmatrix}.$$

- (A) depends on the values of both  $a$  and  $b$   
(B) is independent of the values of both  $a$  and  $b$   
(C) depends on the value of  $a$  but not on the value of  $b$   
(D) depends on the value of  $b$  but not on the value of  $a$

12. If the matrix  $A = \begin{pmatrix} a & 1 \\ 2 & 3 \end{pmatrix}$  has 1 as an eigenvalue, then the determinant of  $A$  is

- (A) 5                      (B) 2                      (C) 4                      (D) 3

13. Let  $a < 500$  be a positive integer. Consider a box containing balls numbered  $a, a+1, \dots, 500$ . Suppose that the ball numbered  $x$  is picked with probability

$$\frac{2xa}{(500+a)(500-a+1)} \quad \text{for } x = a, a+1, \dots, 500.$$

Then the value of  $a$  is

- (A) 251                      (B) 1                      (C) 499                      (D) 2

14. Let  $f(x) = ax + b$  for some  $a, b \in \mathbb{R}$ . Define  $f_n(x)$  inductively by setting

$$f_1(x) = f(x)$$

and

$$f_{n+1}(x) = f(f_n(x)) \quad \text{for } n > 1.$$

If  $f_7(x) = 128x + 381$ , then  $a^b$  equals

- (A)  $\frac{1}{8}$                       (B) 32                      (C)  $\frac{1}{32}$                       (D) 8

15. Let  $n = aaaaaaaaaabcd$  be a 12-digit number divisible by 45 where the digits  $a, b, c, d$  are not necessarily distinct and  $a \neq 0$ . How many such numbers are there?

- (A) 216                      (B) 207                      (C) 189                      (D) 198

16. Let  $a, b$  and  $c$  be the sides of a triangle such that  $c^2 = a^2 + b^2 - ab$ . Then which of the following is always true?

- (A)  $a \leq c$  and  $b \leq c$   
(B)  $a \leq c \leq b$  or  $b \leq c \leq a$   
(C)  $c \leq a$  and  $c \leq b$   
(D) None of the above

17. Let  $X$  be a discrete random variable and  $Y$  be a continuous random variable which is independent of  $X$ . Let  $U = X + Y$  and  $V = XY$ . Choose the correct statement from the options given below.

- (A) Both  $U$  and  $V$  are continuous random variables  
(B)  $U$  is a continuous random variable but  $V$  need not be  
(C)  $U$  is a discrete random variable but  $V$  need not be a continuous random variable  
(D)  $U$  is a discrete random variable and  $V$  is a continuous random variable

18. Suppose that the sample mean and sample standard deviation for a set of  $n$  observations  $x_1, x_2, \dots, x_n$  are  $m$  and  $s$  ( $> 0$ ), respectively. These values are updated to  $m_1$  and  $s_1$  after one more observation  $x_{n+1}$  is added to the data set.

Based on the above information, choose the correct statement from the options given below.

- (A) If  $m_1 = m$  then  $s_1 < s$
- (B) If  $m_1 = m$  then  $s_1 = s$
- (C) If  $m_1 < m$  then  $s_1 = s$
- (D) If  $m_1 < m$  then  $s_1 < s$

19. Suppose that  $X_1, X_2, \dots, X_n$  are independent and identically distributed random variables with probability density function

$$f_\lambda(x) = \begin{cases} \lambda e^{-\lambda x} & \text{if } x \geq 0, \\ 0 & \text{otherwise,} \end{cases}$$

where  $\lambda > 0$ . Let  $Y = X_1 + X_2 + \dots + X_n$ . Then the conditional distribution of  $X_n$  given  $Y = 1$  is

- (A) uniform on  $(0, 1)$
- (B) exponential with mean 1
- (C) beta with parameters  $n - 1$  and 1
- (D) beta with parameters 1 and  $n - 1$

20. Let  $X_1, X_2, \dots$  be a sequence of random variables such that  $E(X_i) = 1$ ,  $\text{Var}(X_i) = 1$  for all  $i$  and  $\text{Cov}(X_i, X_j) = \frac{1}{2}$  for all  $i \neq j$ . Let  $Z_n = \frac{1}{n} \sum_{i=1}^n X_i$ . Then,  $\lim_{n \rightarrow \infty} \text{Var}(Z_n)$  equals

- (A)  $\frac{1}{2}$                       (B)  $\frac{1}{4}$                       (C) 0                      (D) 1

21. Suppose that  $P(A|B) = 0.4$  and  $P(A^c|B^c) = 0.6$ . Then, the two equations are sufficient to find

- (A) neither  $P(A)$  nor  $P(B)$   
 (B) both  $P(A)$  and  $P(B)$   
 (C)  $P(B)$  but not  $P(A)$   
 (D)  $P(A)$  but not  $P(B)$

22. Let  $X_1, \dots, X_n$  be independent and identically distributed normal random variables with mean 0 and variance  $\sigma^2 > 0$ . Define  $T_n = \frac{\sqrt{n} \bar{X}_n}{S_n}$  where  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  and  $S_n^2 = \frac{1}{n} \sum_{i=1}^n X_i^2$ . Then, the distribution of  $T_n$  is

- (A) Student's  $t$  with  $n$  degrees of freedom  
 (B) Student's  $t$  with  $(n - 1)$  degrees of freedom  
 (C) normal with mean 0 and variance 1  
 (D) None of the above



23. Consider a matrix  $\mathbf{M} = \begin{pmatrix} X & -Y \\ Y & X \end{pmatrix}$  where  $X$  and  $Y$  are independent standard normal random variables. Then the probability that  $\mathbf{M}$  is a non-singular matrix is

- (A) 0                      (B) 1                      (C)  $\frac{1}{\sqrt{2}}$                       (D)  $\frac{1}{2}$


24. Let  $(U, V)$  be a point chosen uniformly at random from the unit circle  $\{(u, v) \in \mathbb{R}^2 : u^2 + v^2 = 1\}$ . Then  $\text{Var}(U)$  is

- (A)  $\frac{1}{3}$                       (B)  $\frac{1}{2}$                       (C) 1                      (D)  $\frac{1}{4}$

25. Suppose that we choose 2 cards simultaneously at random from a deck of 20 cards numbered  $1, 2, \dots, 20$ . What is the probability that the smaller of the two numbers divides the larger?


- (A)  $\frac{36}{190}$                       (B)  $\frac{46}{190}$                       (C)  $\frac{56}{190}$                       (D)  $\frac{66}{190}$

26. Let  $X$  be a random variable with probability density function displayed in the following graph.



Match the following random variables with their respective probability density functions

(i)  $X + 3$       (ii)  $X - 3$       (iii)  $2X$       (iv)  $\frac{X}{2}$ .



- (A) (i)–(d), (ii)–(c), (iii)–(b), (iv)–(a)  
 (B) (i)–(d), (ii)–(c), (iii)–(a), (iv)–(b)  
 (C) (i)–(c), (ii)–(d), (iii)–(a), (iv)–(b)  
 (D) (i)–(c), (ii)–(d), (iii)–(b), (iv)–(a)

27. Suppose that we want to fit the regression model

$$y = \beta_1 x + \beta_2 x^2 + \epsilon$$

to 10 pairs of observations  $(x_1, y_1), \dots, (x_{10}, y_{10})$  where  $x_i$ 's take two values, 0 and 1. Which of the following can be estimated using the method of least squares?

- (A) Both  $\beta_1$  and  $\beta_2$
- (B)  $\beta_1$  but not  $\beta_2$
- (C)  $\beta_2$  but not  $\beta_1$
- (D)  $\beta_1 + \beta_2$

28. Consider a bivariate sample  $(X_1, Y_1), \dots, (X_9, Y_9)$  where  $X_i = i$  for  $i = 1, 2, \dots, 9$ . The least squares regression line for this dataset is obtained as  $y = 3 + 2x$ . Later it turns out that  $Y_5$  was recorded wrongly. When the revised regression line is obtained which of the following are possible?

- (A) Intercept can change but slope cannot
- (B) Slope can change but intercept cannot
- (C) Both intercept and slope can change
- (D) Neither intercept nor slope can change

29. Assume  $X_1, \dots, X_n$  are independent and identically distributed  $N(\mu, 1)$  random variables with  $\mu \in \mathbb{R}$ . We want to test  $H_0 : \mu = 0$  versus  $H_1 : \mu \neq 0$ . Consider the following two one-sided testing problems

$$H_{0,A} : \mu = 0 \quad \text{versus} \quad H_{1,A} : \mu > 0$$

and  $H_{0,B} : \mu = 0 \quad \text{versus} \quad H_{1,B} : \mu < 0$ .

Let  $\phi_{A,\eta}(\mathbf{x})$  and  $\phi_{B,\eta}(\mathbf{x})$  denote the most powerful tests of size  $\eta \in (0, 1)$  for  $H_{0,A}$  and  $H_{0,B}$ , respectively. Then, for testing  $H_0$  versus  $H_1$ ,

- (A)  $\phi(\mathbf{x}) = \phi_{A,\eta}(\mathbf{x}) + \phi_{B,\eta}(\mathbf{x})$  is a test of size  $\eta$
- (B)  $\phi(\mathbf{x}) = \phi_{A,\eta}(\mathbf{x}) \phi_{B,\eta}(\mathbf{x})$  is a test of size  $2\eta$
- (C)  $\phi(\mathbf{x}) = \frac{1}{2} \max\{\phi_{A,\eta}(\mathbf{x}), \phi_{B,\eta}(\mathbf{x})\}$  is a test of size  $\frac{\eta}{2}$
- (D)  $\phi(\mathbf{x}) = \phi_{A,\eta}(\mathbf{x}) + \phi_{B,\eta}(\mathbf{x})$  is a test of size  $2\eta$

30. Let  $\phi$  denote the probability density function of the standard normal distribution. Let  $f_\theta$ , for  $\theta \in \{0, 1\}$ , be defined as

$$f_\theta(x) = \begin{cases} \phi(x) & \text{if } \theta = 0, \\ \frac{1}{2}\phi\left(\frac{x-1}{2}\right) & \text{if } \theta = 1. \end{cases}$$

Assume that  $X_1, \dots, X_n$  are independent and identically distributed from the density  $f_\theta(x)$ . Which of the following is a sufficient statistic for  $\theta$ ?

(A)  $\sum_{i=1}^n X_i$

(B)  $\left(\sum_{i=1}^n X_i, \sum_{i=1}^n X_i^2\right)$

(C)  $\sum_{i=1}^n X_i^2$

(D)  $\left(\sum_{i=1}^n X_i, \sum_{i=1}^n \mathbf{1}(|X_i| \geq 2)\right)$