

## FINAL JEE–MAIN EXAMINATION – APRIL, 2024

(Held On Monday 08<sup>th</sup> April, 2024)

TIME : 9 : 00 AM to 12 : 00 NOON

### SECTION-A

31. Three bodies A, B and C have equal kinetic energies and their masses are 400 g, 1.2 kg and 1.6 kg respectively. The ratio of their linear momenta is :

(1)  $1 : \sqrt{3} : 2$                       (2)  $1 : \sqrt{3} : \sqrt{2}$

(3)  $\sqrt{2} : \sqrt{3} : 1$                       (4)  $\sqrt{3} : \sqrt{2} : 1$

**Ans. (1)**

**Sol.**  $KE = \frac{P^2}{2m}$

$P \propto \sqrt{m}$

Hence,  $P_A : P_B : P_C$

$= \sqrt{400} : \sqrt{1200} : \sqrt{1600} = 1 : \sqrt{3} : 2$

32. Average force exerted on a non-reflecting surface at normal incidence is  $2.4 \times 10^{-4}$  N. If  $360 \text{ W/cm}^2$  is the light energy flux during span of 1 hour 30 minutes. Then the area of the surface is:

(1)  $0.2 \text{ m}^2$                       (2)  $0.02 \text{ m}^2$

(3)  $20 \text{ m}^2$                       (4)  $0.1 \text{ m}^2$

**Ans. (2)**

**Sol.** Pressure =  $\frac{C}{A}$

$\Rightarrow \frac{360}{10^{-4} \times 3 \times 10^8} = \frac{2.4 \times 10^{-4}}{A}$

$\Rightarrow A = 2 \times 10^{-2} \text{ m}^2 = 0.02 \text{ m}^2$

33. A proton and an electron are associated with same de-Broglie wavelength. The ratio of their kinetic energies is:

(Assume  $h = 6.63 \times 10^{-34} \text{ J s}$ ,  $m_e = 9.0 \times 10^{-31} \text{ kg}$  and  $m_p = 1836 \text{ times } m_e$ )

(1)  $1 : 1836$                       (2)  $1 : \frac{1}{1836}$

(3)  $1 : \frac{1}{\sqrt{1836}}$                       (4)  $1 : \sqrt{1836}$

**Ans. (1)**

**Sol.**  $\lambda$  is same for both

$P = \frac{h}{\lambda}$  same for both

$P = \sqrt{2mK}$

Hence,

$K \propto \frac{1}{m}$

$\Rightarrow \frac{KE_p}{KE_e} = \frac{m_e}{m_p} = \frac{1}{1836}$

34. A mixture of one mole of monoatomic gas and one mole of a diatomic gas (rigid) are kept at room temperature ( $27^\circ\text{C}$ ). The ratio of specific heat of gases at constant volume respectively is:

(1)  $\frac{7}{5}$                       (2)  $\frac{3}{2}$

(3)  $\frac{3}{5}$                       (4)  $\frac{5}{3}$

**Ans. (3)**

**Sol.**  $\frac{(C_v)_{\text{mono}}}{(C_v)_{\text{dia}}} = \frac{\frac{3}{2}R}{\frac{5}{2}R} = \frac{3}{5}$

35. In an expression  $a \times 10^b$  :

(1) a is order of magnitude for  $b \leq 5$

(2) b is order of magnitude for  $a \leq 5$

(3) b is order of magnitude for  $5 < a \leq 10$

(4) b is order of magnitude for  $a \geq 5$

**Ans. (2)**

**Sol.**  $a \times 10^b$

if  $a \leq 5$  order is b

$a > 5$  order is  $b + 1$




Sol.  $\frac{\Delta Y}{Y} = \frac{\Delta m}{m} + \frac{\Delta \ell}{\ell}$   
 $= \frac{5}{500} + \frac{0.02}{2} = 0.01 + 0.01$

$\frac{\Delta Y}{Y} = 0.02 \Rightarrow \% \frac{\Delta Y}{Y} = 2\%$

41. Two different adiabatic paths for the same gas intersect two isothermal curves as shown in P-V diagram. The relation between the ratio  $\frac{V_a}{V_d}$  and the

ratio  $\frac{V_b}{V_c}$  is:



- (1)  $\frac{V_a}{V_d} = \left(\frac{V_b}{V_c}\right)^{-1}$       (2)  $\frac{V_a}{V_d} \neq \frac{V_b}{V_c}$   
 (3)  $\frac{V_a}{V_d} = \frac{V_b}{V_c}$       (4)  $\frac{V_a}{V_d} = \left(\frac{V_b}{V_c}\right)^2$

Ans. (3)

Sol. For adiabatic process

$TV^{\gamma-1} = \text{constant}$

$T_a \cdot V_a^{\gamma-1} = T_d \cdot V_d^{\gamma-1}$

$\left(\frac{V_a}{V_d}\right)^{\gamma-1} = \frac{T_d}{T_a}$

$\left(\frac{V_b}{V_c}\right)^{\gamma-1} = \frac{T_c}{T_b}$

$\frac{V_a}{V_d} = \frac{V_b}{V_c} \quad \left( \begin{array}{l} \because T_d = T_c \\ T_a = T_b \end{array} \right)$

42. Two planets A and B having masses  $m_1$  and  $m_2$  move around the sun in circular orbits of  $r_1$  and  $r_2$  radii respectively. If angular momentum of A is  $L$  and that of B is  $3L$ , the ratio of time period  $\left(\frac{T_A}{T_B}\right)$  is:

- (1)  $\left(\frac{r_2}{r_1}\right)^{\frac{3}{2}}$       (2)  $\left(\frac{r_1}{r_2}\right)^3$   
 (3)  $\frac{1}{27} \left(\frac{m_2}{m_1}\right)^3$       (4)  $27 \left(\frac{m_1}{m_2}\right)^3$

Ans. (3)

Sol.  $\frac{\pi r_1^2}{T_A} = \frac{L}{2m_1}$  ..... (1)

$\frac{\pi r_2^2}{T_B} = \frac{3L}{2m_2}$  ..... (2)

$\Rightarrow \frac{T_A}{T_B} = 3 \cdot \frac{m_1}{m_2} \cdot \left(\frac{r_1}{r_2}\right)^2$

$\left(\frac{T_A}{T_B}\right)^2 = \left(\frac{r_1}{r_2}\right)^3 \Rightarrow \left(\frac{r_1}{r_2}\right)^2 = \left(\frac{T_A}{T_B}\right)^{\frac{4}{3}}$

$\Rightarrow \frac{1}{27} \cdot \left(\frac{m_2}{m_1}\right)^3 = \left(\frac{T_A}{T_B}\right)$

43. A LCR circuit is at resonance for a capacitor  $C$ , inductance  $L$  and resistance  $R$ . Now the value of resistance is halved keeping all other parameters same. The current amplitude at resonance will be now:

- (1) Zero      (2) double  
 (3) same      (4) halved

Ans. (2)

Sol. In resonance  $Z = R$


$I = \frac{V}{R}$

$R \rightarrow \text{halved}$

$\Rightarrow I \rightarrow 2I$

$I$  becomes doubled.

44. The output Y of following circuit for given inputs is :



- (1)  $A \cdot B(A + B)$                       (2)  $A \cdot B$   
 (3) 0    (4)  $\bar{A} \cdot B$

Ans. (3)

Sol. By truth table

A	B	Y
0	0	0
0	1	0
1	0	0
1	1	0

45. Two charged conducting spheres of radii a and b are connected to each other by a conducting wire. The ratio of charges of the two spheres respectively is:

- (1)  $\sqrt{ab}$                                       (2) ab  
 (3)  $\frac{a}{b}$     (4)  $\frac{b}{a}$

Ans. (3)

Sol. Potential at surface will be same

$$\frac{Kq_1}{a} = \frac{Kq_2}{b}$$

$$\frac{q_1}{q_2} = \frac{a}{b}$$

46. Correct Bernoulli's equation is (symbols have their usual meaning) :

- (1)  $P + mgh + \frac{1}{2}mv^2 = \text{constant}$   
 (2)  $P + \rho gh + \frac{1}{2}\rho v^2 = \text{constant}$   
 (3)  $P + \rho gh + \rho v^2 = \text{constant}$   
 (4)  $P + \frac{1}{2}\rho gh + \frac{1}{2}\rho v^2 = \text{constant}$

Ans. (2)

Sol.  $P + \rho gh + \frac{1}{2}\rho v^2 = \text{constant}$

47. A player caught a cricket ball of mass 150 g moving at a speed of 20 m/s. If the catching process is completed in 0.1 s, the magnitude of force exerted by the ball on the hand of the player is:

- (1) 150 N    (2) 3 N  
 (3) 30 N    (4) 300 N

Ans. (3)

Sol.  $F = \frac{\Delta P}{\Delta t} = \frac{mv - 0}{0.1}$

$$= \frac{150 \times 10^{-3} \times 20}{0.1} = 30 \text{ N}$$

48. A stationary particle breaks into two parts of masses  $m_A$  and  $m_B$  which move with velocities  $v_A$  and  $v_B$  respectively. The ratio of their kinetic energies ( $K_B : K_A$ ) is :

- (1)  $v_B : v_A$                                       (2)  $m_B : m_A$   
 (3)  $m_B v_B : m_A v_A$                       (4) 1 : 1

Ans. (1)

Sol. Initial momentum is zero.

Hence  $|P_A| = |P_B|$

$\Rightarrow m_A v_B = m_B v_A$

$$\frac{(KE)_A}{(KE)_B} = \frac{\frac{1}{2}m_A v_A^2}{\frac{1}{2}m_B v_B^2} = \frac{v_A}{v_B}$$

$$\frac{(KE)_B}{(KE)_A} = \frac{v_B}{v_A}$$

49. Critical angle of incidence for a pair of optical media is  $45^\circ$ . The refractive indices of first and second media are in the ratio:

- (1)  $\sqrt{2} : 1$                                       (2) 1 : 2  
 (3)  $1 : \sqrt{2}$                                       (4) 2 : 1

Ans. (1)

Sol.  $\sin\theta_c = \frac{\mu_R}{\mu_d} = \frac{\mu_2}{\mu_1}$

$\sin 45^\circ = \frac{\mu_2}{\mu_1}$

$\Rightarrow \frac{1}{\sqrt{2}} = \frac{\mu_2}{\mu_1}$

$\Rightarrow \frac{\mu_1}{\mu_2} = \frac{\sqrt{2}}{1}$

50. The diameter of a sphere is measured using a vernier caliper whose 9 divisions of main scale are equal to 10 divisions of vernier scale. The shortest division on the main scale is equal to 1 mm. The main scale reading is 2 cm and second division of vernier scale coincides with a division on main scale. If mass of the sphere is 8.635 g, the density of the sphere is:

- (1) 2.5 g/cm<sup>3</sup>                      (2) 1.7 g/cm<sup>3</sup>
- (3) 2.2 g/cm<sup>3</sup>                      (4) 2.0 g/cm<sup>3</sup>

Ans. (4)

Sol. Given 9MSD = 10VSD

mass = 8.635 g

LC = 1 MSD - 1 VSD

LC = 1 MSD -  $\frac{9}{10}$  MSD

LC =  $\frac{1}{10}$  MSD

LC = 0.01 cm


Reading of diameter = MSR + LC × VSR  
 = 2 cm + (0.01) × (2)  
 = 2.02 cm

Volume of sphere =  $\frac{4}{3}\pi\left(\frac{d}{2}\right)^3 = \frac{4}{3}\pi\left(\frac{2.02}{2}\right)^3$   
 = 4.32 cm<sup>3</sup>

Density =  $\frac{\text{mass}}{\text{volume}} = \frac{8.635}{4.32} = 1.998 \sim 2.00\text{g}$



SECTION-B

51. A uniform thin metal plate of mass 10 kg with dimensions is shown. The ratio of x and y coordinates of center of mass of plate in  $\frac{n}{9}$ . The value of n is \_\_\_\_\_.



Ans. (15)

Sol.  $m_1 = \sigma \times 5 = 10\text{Kg}$



$\Rightarrow m_1 x_1 + m_2 x_2 = m_3 x_3$

$10x_1 + 2(1.5) = 12(1.5) \Rightarrow x_1 = 1.5\text{ cm}$

$\Rightarrow m_1 y_1 + m_2 y_2 = m_3 y_3$

$10y_1 + 2(1.5) = 12 \times 1 \Rightarrow y_1 = 0.9\text{ cm}$

$\frac{x_1}{y_1} = \frac{1.5}{0.9} = \frac{15}{9}$

n = 15

52. An electron with kinetic energy 5 eV enters a region of uniform magnetic field of 3 μT perpendicular to its direction. An electric field E is applied perpendicular to the direction of velocity and magnetic field. The value of E, so that electron moves along the same path, is \_\_\_\_\_ NC<sup>-1</sup>.  
(Given, mass of electron = 9 × 10<sup>-31</sup> kg, electric charge = 1.6 × 10<sup>-19</sup>C)

Ans. (4)

Sol. For the given condition of moving undeflected, net force should be zero.

$$qE = qvB$$


$$E = vB$$

$$= \sqrt{\frac{2 \times KE}{m}} \times B$$

$$= \sqrt{\frac{2 \times 5 \times 1.6 \times 10^{-19}}{9 \times 10^{-31}}} \times 3 \times 10^{-6}$$

$$= 4 \text{ N/C}$$

53. A square loop PQRS having 10 turns, area 3.6 × 10<sup>-3</sup> m<sup>2</sup> and resistance 100 Ω is slowly and uniformly being pulled out of a uniform magnetic field of magnitude B = 0.5 T as shown. Work done in pulling the loop out of the field in 1.0 s is \_\_\_\_\_ × 10<sup>-6</sup> J.



Ans. (3)

Sol.  $\epsilon = NB\ell v$

$$i = \frac{\epsilon}{R} = \frac{NB\ell v}{R}$$

$$F = N(i\ell B) = \frac{N^2 B^2 \ell^2 v}{R}$$

$$W = F \times \ell = \frac{N^2 B^2 \ell^3}{R} \left( \frac{\ell}{t} \right)$$

$$A = \ell^2$$

$$W = \frac{(10 \times 10)(0.5)^2 \times (3.6 \times 10^{-3})^2}{100 \times 1}$$

$$W = 3.24 \times 10^{-6} \text{ J}$$

54. Resistance of a wire at 0 °C, 100 °C and t °C is found to be 10 Ω, 10.2 Ω and 10.95 Ω respectively. The temperature t in Kelvin scale is \_\_\_\_\_.

Ans. (748)

Sol.  $R = R_0(1 + \alpha\Delta T)$

$$\frac{\Delta R}{R_0} = \alpha\Delta T$$

Case-I

$$0^\circ\text{C} \rightarrow 100^\circ\text{C}$$

$$\frac{10.2 - 10}{10} = \alpha(100 - 0) \quad \dots (1)$$

Case-II

$$0^\circ\text{C} \rightarrow t^\circ\text{C}$$

$$\frac{10.95 - 10}{10} = \alpha(t - 0) \quad \dots (2)$$

$$\Rightarrow \frac{t}{100} = \frac{0.95}{0.2} = 475^\circ\text{C}$$

$$t = 475 + 273 = 748 \text{ K}$$

55. An electric field,  $\vec{E} = \frac{2\hat{i} + 6\hat{j} + 8\hat{k}}{\sqrt{6}}$  passes through the surface of 4 m<sup>2</sup> area having unit vector  $\hat{n} = \left( \frac{2\hat{i} + \hat{j} + \hat{k}}{\sqrt{6}} \right)$ . The electric flux for that surface is \_\_\_\_\_ V m.

Ans. (12)

Sol.  $\phi = \vec{E} \cdot \vec{A}$

$$= \left( \frac{2\hat{i} + 6\hat{j} + 8\hat{k}}{\sqrt{6}} \right) \cdot 4 \left( \frac{2\hat{i} + \hat{j} + \hat{k}}{\sqrt{6}} \right)$$

$$= \frac{4}{6} \times (4 + 6 + 8) = 12 \text{ Vm}$$

56. A liquid column of height 0.04 cm balances excess pressure of soap bubble of certain radius. If density of liquid is  $8 \times 10^3 \text{ kg m}^{-3}$  and surface tension of soap solution is  $0.28 \text{ Nm}^{-1}$ , then diameter of the soap bubble is \_\_\_\_\_ cm.  
(if  $g = 10 \text{ ms}^{-2}$ )

Ans. (7)

Sol.  $\rho gh = \frac{4S}{R}$

$$\Rightarrow R = \frac{4 \times 0.28}{8 \times 10^3 \times 10 \times 4 \times 10^{-4}}$$

$$\Rightarrow \frac{0.28}{8} \text{ m} = \frac{28}{8} \text{ cm}$$

$$\Rightarrow R = 3.5 \text{ cm}$$

$$\text{Diameter} = 7 \text{ cm}$$

57. A closed and an open organ pipe have same lengths. If the ratio of frequencies of their seventh overtones is  $\left(\frac{a-1}{a}\right)$  then the value of a is \_\_\_\_\_.

Ans. (16)

Sol. For closed organ pipe

$$f_c = (2n+1) \frac{v}{4l} = \frac{15v}{4l}$$


For open organ pipe

$$f_o = (n+1) \frac{v}{2l} = \frac{8v}{2l}$$

$$\frac{f_c}{f_o} = \frac{15}{16} = \frac{a-1}{a}$$


$$\Rightarrow a = 16$$

58. Three vectors  $\vec{OP}$ ,  $\vec{OQ}$  and  $\vec{OR}$  each of magnitude A are acting as shown in figure. The resultant of the three vectors is  $A\sqrt{x}$ . The value of x is \_\_\_\_\_.



Ans. (3)

Sol.



$$\vec{R} = \left(A + \frac{A}{\sqrt{2}}\right)\hat{i} + \left(A - \frac{A}{\sqrt{2}}\right)\hat{j}$$

$$|\vec{R}| = \sqrt{\left(A + \frac{A}{\sqrt{2}}\right)^2 + \left(A - \frac{A}{\sqrt{2}}\right)^2} = \sqrt{3}A$$

59. A parallel beam of monochromatic light of wavelength 600 nm passes through single slit of 0.4 mm width. Angular divergence corresponding to second order minima would be \_\_\_\_\_  $\times 10^{-3}$  rad.

Ans. (6)

Sol.  $\sin \theta \approx \theta = \frac{2\lambda}{b}$

$$= \frac{2 \times 600 \times 10^{-9}}{4 \times 10^{-4}} = 3 \times 10^{-3} \text{ rad}$$

$$\text{Total divergence} = (3 + 3) \times 10^{-3} = 6 \times 10^{-3} \text{ rad}$$

60. In an alpha particle scattering experiment distance of closest approach for the  $\alpha$  particle is  $4.5 \times 10^{-14} \text{ m}$ . If target nucleus has atomic number 80, then maximum velocity of  $\alpha$ -particle is \_\_\_\_\_  $\times 10^5 \text{ m/s}$  approximately.

$$\left(\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ SI unit, mass of } \alpha \text{ particle} = \right.$$

$$\left. 6.72 \times 10^{-27} \text{ kg}\right)$$

Ans. (156)

Sol.  $v = \sqrt{\frac{4KZe^2}{mr_{\min}}}$

$$= \sqrt{\frac{4 \times 9 \times 10^9 \times 80}{6.72 \times 10^{-27} \times 4.5 \times 10^{-14}}} \times 1.6 \times 10^{-19}$$

$$= 9.759 \times 10^{25} \times 1.6 \times 10^{-19}$$

$$= 156 \times 10^5 \text{ m/s}$$