

FINAL JEE-MAIN EXAMINATION – JANUARY, 2023

(Held On Wednesday 1st February, 2023)

TIME : 9 : 00 AM to 12 : 00

SECTION-A

61. $\lim_{n \rightarrow \infty} \left(\frac{1}{1+n} + \frac{1}{2+n} + \frac{1}{3+n} + \dots + \frac{1}{2n} \right)$ is equal to :-
 (1) 0 (2) $\log_e 2$
 (3) $\log_e \left(\frac{3}{2} \right)$ (4) $\log_e \left(\frac{2}{3} \right)$

Official Ans. by NTA (2)

Ans. (2)

Sol. $\lim_{n \rightarrow \infty} \left(\frac{1}{1+n} + \dots + \frac{1}{n+n} \right) = \lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{1}{n+r}$
 $= \lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{1}{n} \left(\frac{1}{1+\frac{r}{n}} \right)$
 $= \int_0^1 \frac{1}{1+x} dx$ $\int_0^1 \frac{1}{1+x} dx = \ln 2$

62. The negation of the expression $q \vee ((\sim q) \wedge p)$ is equivalent to
 (1) $(\sim p) \wedge (\sim q)$ (2) $p \wedge (\sim q)$
 (3) $(\sim p) \vee (\sim q)$ (4) $(\sim p) \vee q$

Official Ans. by NTA (1)

Ans. (1)

Sol. $\sim (q \vee ((\sim q) \wedge p))$
 $= \sim q \wedge \sim ((\sim q) \wedge p)$
 $= \sim q \wedge (q \vee \sim p)$
 $= (\sim q \wedge q) \vee (\sim q \wedge \sim p)$
 $= (\sim q \wedge \sim p)$

63. In a binomial distribution $B(n, p)$, the sum and product of the mean & variance are 5 and 6 respectively, then find $6(n + p - q)$ is equal to :-
 (1) 51
 (2) 52
 (3) 53
 (4) 50

Official Ans. by NTA (2)

Ans. (2)

Sol. $np + npq = 5, np \cdot npq = 6$
 $np(1+q) = 5, n^2p^2q = 6$
 $n^2p^2(1+q)^2 = 25, n^2p^2q = 6$
 $\frac{6}{q}(1+q)^2 = 25$
 $6q^2 + 12q + 6 = 25q$
 $6q^2 - 13q + 6 = 0$
 $6q^2 - 9q - 4q + 6 = 0$
 $(3q-2)(2q-3) = 0$

$q = \frac{2}{3}, \frac{3}{2}, q = \frac{2}{3}$ is accepted

$p = \frac{1}{3} \Rightarrow n \cdot \frac{1}{3} + n \cdot \frac{1}{3} \cdot \frac{2}{3} = 5$

$\frac{3n+2n}{9} = 5$

$n = 9$

So $6(n+p-q) = 6 \left(9 + \frac{1}{3} - \frac{2}{3} \right) = 52$

64. The sum to 10 terms of the series

$\frac{1}{1+1^2+1^4} + \frac{2}{1+2^2} + \frac{3}{1+3^2} + \dots$ is:-

(1) $\frac{59}{111}$ (2) $\frac{55}{111}$

(3) $\frac{56}{111}$ (4) $\frac{58}{111}$

Official Ans. by NTA (2)

Ans. (2)

Sol. $T_r = \frac{(r^2+r+1) - (r^2-r+1)}{2(r^4+r^2+1)}$

$\Rightarrow T_r = \frac{1}{2} \left[\frac{1}{r^2-r+1} - \frac{1}{r^2+r+1} \right]$

$T_1 = \frac{1}{2} \left[\frac{1}{1} - \frac{1}{3} \right]$

$T_2 = \frac{1}{2} \left[\frac{1}{3} - \frac{1}{7} \right]$

$$T_3 = \frac{1}{2} \left[\frac{1}{7} - \frac{1}{13} \right]$$

$$\vdots$$

$$T_{10} = \frac{1}{2} \left[\frac{1}{91} - \frac{1}{111} \right]$$

$$\Rightarrow \sum_{r=1}^{10} T_r = \frac{1}{2} \left[1 - \frac{1}{111} \right] = \frac{55}{111}$$

65. The value of

$$\frac{1}{1!50!} + \frac{1}{3!48!} + \frac{1}{5!46!} + \dots + \frac{1}{49!2!} + \frac{1}{51!1!}$$
 is

- (1) $\frac{2^{50}}{50!}$ (2) $\frac{2^{50}}{51!}$
 (3) $\frac{2^{51}}{51!}$ (4) $\frac{2^{51}}{50!}$

Official Ans. by NTA (2)

Ans. (2)

Sol.
$$\sum_{r=1}^{26} \frac{1}{(2r-1)!(51-(2r-1))!} = \sum_{r=1}^{26} {}^{51}C_{(2r-1)} \frac{1}{51!}$$

$$= \frac{1}{51!} \{ {}^{51}C_1 + {}^{51}C_3 + \dots + {}^{51}C_{51} \} = \frac{1}{51!} (2^{50})$$


66. If the orthocentre of the triangle, whose vertices are (1, 2), (2, 3) and (3, 1) is (α, β) , then the quadratic equation whose roots are $\alpha + 4\beta$ and $4\alpha + \beta$, is

- (1) $x^2 - 19x + 90 = 0$
 (2) $x^2 - 18x + 80 = 0$
 (3) $x^2 - 22x + 120 = 0$
 (4) $x^2 - 20x + 99 = 0$

Official Ans. by NTA (4)

Ans. (4)

Sol.



Here $m_{BH} \times m_{AC} = -1$

$$\left(\frac{\beta - 3}{\alpha - 2} \right) \left(\frac{1}{-2} \right) = -1$$

$$\beta - 3 = 2\alpha - 4$$

$$\beta = 2\alpha - 1$$

$$m_{AH} \times m_{BC} = -1$$

$$\Rightarrow \left(\frac{\beta - 2}{\alpha - 1} \right) (-2) = -1$$

$$\Rightarrow 2\beta - 4 = \alpha - 1$$

$$\Rightarrow 2(2\alpha - 1) = \alpha + 3$$

$$\Rightarrow 3\alpha = 5$$

$$\alpha = \frac{5}{3}, \beta = \frac{7}{3} \Rightarrow H\left(\frac{5}{3}, \frac{7}{3}\right)$$

$$\alpha + 4\beta = \frac{5}{3} + \frac{28}{3} = \frac{33}{3} = 11$$

$$\beta + 4\alpha = \frac{7}{3} + \frac{20}{3} = \frac{27}{3} = 9$$

$$x^2 - 20x + 99 = 0$$


67. For a triangle ABC, the value of $\cos 2A + \cos 2B + \cos 2C$ is least. If its inradius is 3 and incentre is M, then which of the following is NOT correct?

- (1) Perimeter of ΔABC is $18\sqrt{3}$
 (2) $\sin 2A + \sin 2B + \sin 2C = \sin A + \sin B + \sin C$
 (3) $\overrightarrow{MA} \cdot \overrightarrow{MB} = -18$
 (4) area of ΔABC is $\frac{27\sqrt{3}}{2}$

Official Ans. by NTA (4)

Ans. (4)

Sol.



If $\cos 2A + \cos 2B + \cos 2C$ is minimum then $A = B = C = 60^\circ$

So ΔABC is equilateral

Now in-radius $r = 3$

So in ΔMBD we have

$$\tan 30^\circ = \frac{MD}{BD} = \frac{r}{a/2} = \frac{6}{a}$$

$$1/\sqrt{3} = \frac{1}{a} = a = 6\sqrt{3}$$

$$\text{Perimeter of } \Delta ABC = 18\sqrt{3}$$

$$\text{Area of } \Delta ABC = \frac{\sqrt{3}}{4} a^2 = 27\sqrt{3}$$

68. The combined equation of the two lines $ax + by + c = 0$ and $a'x + b'y + c' = 0$ can be written as $(ax + by + c)(a'x + b'y + c') = 0$

The equation of the angle bisectors of the lines represented by the equation $2x^2 + xy - 3y^2 = 0$ is

(1) $3x^2 + 5xy + 2y^2 = 0$

(2) $x^2 - y^2 + 10xy = 0$

(3) $3x^2 + xy - 2y^2 = 0$

(4) $x^2 - y^2 - 10xy = 0$

Official Ans. by NTA (4)

□ Ans. (4)

Sol.

Equation of the pair of angle bisector for the homogenous equation $ax^2 + 2hxy + by^2 = 0$ is given as

$$\frac{x}{a-b} = \frac{y}{h}$$

Here $a = 2, h = 1/2$ & $b = -3$

Equation will become

$$\frac{x^2 - y^2}{2 - (-3)} = \frac{xy}{1/2}$$

$$x^2 - y^2 = 10xy$$

$$x^2 - y^2 - 10xy = 0$$

69. The shortest distance between the lines

$$\frac{x-5}{1} = \frac{y-2}{2} = \frac{z-4}{-3} \text{ and } \frac{x+3}{1} = \frac{y+5}{4} = \frac{z-1}{-5} \text{ is}$$

(1) $7\sqrt{3}$

(2) $5\sqrt{3}$

(3) $6\sqrt{3}$

(4) $4\sqrt{3}$

Official Ans. by NTA (3)

Ans. (3)

Sol.

Shortest distance between two lines

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{a_2} = \frac{z-z_1}{a_3} \text{ \&}$$

$$\frac{x-x_2}{b_1} = \frac{y-y_2}{b_2} = \frac{z-z_2}{b_3} \text{ is given as}$$

$$\begin{vmatrix} x_1 - x_2 & y_1 - y_2 & z_1 - z_2 \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

$$\sqrt{(a_1 b_3 - a_3 b_2)^2 + (a_1 b_3 - a_3 b_1)^2 + (a_1 b_2 - a_2 b_1)^2}$$

$$\begin{vmatrix} 5 - (-3) & 2 - (-5) & 4 - 1 \\ 1 & 2 & -3 \\ 1 & 4 & -5 \end{vmatrix}$$

$$\sqrt{(-10+12)^2 + (-5+3)^2 + (4-2)^2}$$

$$\begin{vmatrix} 8 & 7 & 3 \\ 1 & 2 & -3 \\ 1 & 4 & -5 \end{vmatrix}$$

$$\sqrt{(2)^2 + (2)^2 + (2)^2}$$

$$= \frac{|8(-10+12) - 7(-5+3) + 3(4-2)|}{\sqrt{4+4+4}}$$

$$= \frac{16+14+6}{\sqrt{12}} = \frac{36}{\sqrt{12}} = \frac{36}{2\sqrt{3}}$$

$$= \frac{18}{\sqrt{3}} = 6\sqrt{3}$$

70. Let S denote the set of all real values of λ such that the system of equations

$$\lambda x + y + z = 1$$

$$x + \lambda y + z = 1$$

$$x + y + \lambda z = 1$$

is inconsistent, then $\sum_{\lambda \in S} (|\lambda|^2 + |\lambda|)$ is equal to

(1) 2

(2) 12

(3) 4

(4) 6

Official Ans. by NTA (4)

Ans. (4)

Sol. $\begin{vmatrix} \lambda & 1 & 1 \\ 1 & \lambda & 1 \\ 1 & 1 & \lambda \end{vmatrix} = 0$

$$(\lambda + 2) \begin{vmatrix} 1 & 1 & 1 \\ 1 & \lambda & 1 \\ 1 & 1 & \lambda \end{vmatrix} = 0$$

$$(\lambda + 2)[1(\lambda^2 - 1) - 1(\lambda - 1) + (1 - \lambda)] = 0$$

$$(\lambda + 2)(\lambda^2 - 2\lambda + 1) = 0$$

$$(\lambda + 2)(\lambda - 1)^2 = 0 \Rightarrow \lambda = -2, \lambda = 1$$

at $\lambda = 1$ system has infinite solution, for inconsistent $\lambda = -2$

$$\text{so } \sum (|-2|^2 + |-2|) = 6$$

71. Let

$$S = \left\{ x : x \in \mathbb{R} \text{ and } (\sqrt{3} + \sqrt{2})^{x^2-4} + (\sqrt{3} - \sqrt{2})^{x^2-4} = 10 \right\}.$$

Then $n(S)$ is equal to

(1) 2 (2) 4

(3) 6 (4) 0

Official Ans. by NTA (2)

Ans. (2)

Sol. Let $(\sqrt{3} + \sqrt{2})^{x^2-4} = t$

$$t + \frac{1}{t} = 10$$

$$\Rightarrow t = 5 + 2\sqrt{6}, 5 - 2\sqrt{6}$$

$$\Rightarrow \sqrt{x^2-4} = \sqrt{5+2\sqrt{6}}, \sqrt{5-2\sqrt{6}}$$

$$\Rightarrow x^2 - 4 = 2, -2 \text{ or } x^2 = 6, 2$$

$$\Rightarrow x = \pm\sqrt{2}, \pm\sqrt{6}$$

72. Let S be the set of all solutions of the equation

$$\cos^{-1}(2x) - 2\cos^{-1}(\sqrt{1-x^2}) = \pi, x \in \left[-\frac{1}{2}, \frac{1}{2}\right].$$

Then $\sum_{x \in S} 2\sin^{-1}(x^2 - 1)$ is equal to

(1) 0 (2) $\frac{-2\pi}{3}$

(3) $\pi - \sin^{-1}\left(\frac{\sqrt{3}}{4}\right)$ (4) $\pi - 2\sin^{-1}\left(\frac{\sqrt{3}}{4}\right)$

Official Ans. by NTA (2)

Ans. (2)

Sol. $\cos^{-1}(2x) - 2\cos^{-1}\sqrt{1-x^2} = \pi$

$$\cos^{-1}(2x) - \cos^{-1}(2(1-x^2) - 1) = \pi$$

$$\cos^{-1}(2x) - \cos^{-1}(1-2x^2) = \pi$$

$$-\cos^{-1}(1-2x^2) = \pi - \cos^{-1}(2x)$$

Taking cos both sides we get

$$\cos(-\cos^{-1}(1-2x^2)) = \cos(\pi - \cos^{-1}(2x))$$

$$1 - 2x^2 = -2x$$

$$2x^2 - 2x - 1 = 0$$

On solving, $x = \frac{1-\sqrt{3}}{2}, \frac{1+\sqrt{3}}{2}$

As $x = [-1/2, 1/2]$, $x = \frac{1+\sqrt{3}}{2}$ = rejected

So $x = \frac{1-\sqrt{3}}{2} \Rightarrow x^2 - 1 = -\sqrt{3}/2$

$$= 2\sin^{-1}(x^2 - 1) = 2\sin^{-1}\left(\frac{-\sqrt{3}}{2}\right) = \frac{-2\pi}{3}$$

73. If the center and radius of the circle $\left|\frac{z-2}{z-3}\right| = 2$ are

respectively (α, β) and γ , then $3(\alpha + \beta + \gamma)$ is equal to

(1) 11

(2) 9

(3) 10

(4) 12

Official Ans. by NTA (4)

Ans. (4)

Sol.

$$\sqrt{(x-2)^2 + y^2} = 2\sqrt{(x-3)^2 + y^2}$$

$$= x^2 + y^2 - 4x + 4 = 4x^2 + 4y^2 - 24x + 36$$

$$= 3x^2 + 3y^2 - 20x + 32 = 0$$

$$= x^2 + y^2 - \frac{20}{3}x + \frac{32}{3} = 0$$

$$= (\alpha, \beta) = \left(\frac{10}{3}, 0\right)$$

$$\gamma = \sqrt{\frac{100}{9} - \frac{32}{3}} = \sqrt{\frac{4}{9}} = \frac{2}{3}$$

$$3(\alpha, \beta, \gamma) = 3\left(\frac{10}{3} + \frac{2}{3}\right)$$

$$= 12$$

74. If $y = y(x)$ is the solution curve of the differential equation $\frac{dy}{dx} + y \tan x = x \sec x$, $0 \leq x \leq \frac{\pi}{3}$,

$y(0) = 1$, then $y\left(\frac{\pi}{6}\right)$ is equal to

(1) $\frac{\pi}{12} - \frac{\sqrt{3}}{2} \log_e \left(\frac{2}{e\sqrt{3}} \right)$

(2) $\frac{\pi}{12} + \frac{\sqrt{3}}{2} \log_e \left(\frac{2\sqrt{3}}{e} \right)$

(3) $\frac{\pi}{12} - \frac{\sqrt{3}}{2} \log_e \left(\frac{2\sqrt{3}}{e} \right)$

(4) $\frac{\pi}{12} + \frac{\sqrt{3}}{2} \log_e \left(\frac{2}{e\sqrt{3}} \right)$

Official Ans. by NTA (1)

Ans. (1)

Sol. Here I.F. = $\sec x$

Then solution of D.E :

$$y(\sec x) = x \tan x - \ln(\sec x) + c$$

Given $y(0) = 1 \Rightarrow c = 1$

$$\therefore y(\sec x) = x \tan x - \ln(\sec x) + 1$$

At $x = \frac{\pi}{6}$, $y = \frac{\pi}{12} + \frac{\sqrt{3}}{2} \ln \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{2}$

75. Let R be a relation on \mathbb{R} , given by $R = \{(a, b) : 3a - 3b + \sqrt{7} \text{ is an irrational number}\}$. Then R is

- (1) Reflexive but neither symmetric nor transitive
- (2) Reflexive and transitive but not symmetric
- (3) Reflexive and symmetric but not transitive
- (4) An equivalence relation

Official Ans. by NTA (1)

Ans. (1)

Sol. Check for reflexivity:

As $3(a - a) + \sqrt{7} = \sqrt{7}$ which belongs to relation so relation is reflexive

Check for symmetric:

Take $a = \frac{\sqrt{7}}{3}, b = 0$

Now $(a, b) \in R$ but $(b, a) \notin R$

As $3(b - a) + \sqrt{7} = 0$ which is rational so relation is not symmetric.

Check for Transitivity:

Take (a, b) as $\left(\frac{\sqrt{7}}{3}, 1\right)$

& (b, c) as $\left(1, \frac{2\sqrt{7}}{3}\right)$

So now $(a, b) \in R$ & $(b, c) \in R$ but $(a, c) \notin R$ which means relation is not transitive

76. Let the image of the point $P(2, -1, 3)$ in the plane $x + 2y - z = 0$ be Q . Then the distance of the plane $3x + 2y + z + 29 = 0$ from the point Q is

(1) $\frac{22\sqrt{2}}{7}$

(2) $\frac{24\sqrt{2}}{7}$


(3) $2\sqrt{14}$

(4) $3\sqrt{14}$

Official Ans. by NTA (4)

Ans. (4)

Sol.



eq. of line $PM \frac{x-2}{1} = \frac{y+1}{2} = \frac{z-3}{-1} = \lambda$

any point on line = $(\lambda + 2, 2\lambda - 1, -\lambda + 3)$

for point 'm' $(\lambda + 2) + 2(2\lambda - 1) - (3 - \lambda) = 0$

$$\lambda = \frac{1}{2}$$

Point m $\left(\frac{1}{2} + 2, 2 \times \frac{1}{2} - 1, -\frac{1}{2} + 3\right)$

= $\left(\frac{5}{2}, 0, \frac{5}{2}\right)$

For Image $Q (\alpha, \beta, \gamma)$

$$\frac{\alpha+2}{2} = \frac{5}{2}, \frac{\beta-1}{2} = 0,$$

$$\frac{\gamma+3}{2} = \frac{5}{2}$$

$$Q : (3, 1, 2)$$

$$d = \frac{|3(3) + 2(1) + 2 + 29|}{\sqrt{3^2 + 2^2 + 1^2}}$$

$$d = \frac{42}{\sqrt{14}} = 3\sqrt{14}$$

77. Let $f(x) = \begin{vmatrix} 1 + \sin^2 x & \cos^2 x & \sin 2x \\ \sin^2 x & 1 + \cos^2 x & \sin 2x \\ \sin^2 x & \cos^2 x & 1 + \sin 2x \end{vmatrix}$,

$$x \in \left[\frac{\pi}{6}, \frac{\pi}{3} \right].$$
 If α & β respectively are the maximum

and the minimum values of f , then

$$(1) \beta^2 - 2\sqrt{\alpha} = \frac{19}{4}$$

$$(2) \beta^2 + 2\sqrt{\alpha} = \frac{19}{4}$$

$$(3) \alpha^2 - \beta^2 = 4\sqrt{3}$$

$$(4) \alpha^2 + \beta^2 = \frac{9}{2}$$

Official Ans. by NTA (1)

Ans. (1)

Sol.

$$C_1 \rightarrow C_1 + C_2 + C_3$$

$$f(x) = \begin{vmatrix} 2 + \cos^2 x & \sin 2x & \sin 2x \\ 2 + \sin 2x & 1 + \cos^2 x & \sin 2x \\ 2 + \sin 2x & \cos^2 x & 1 + \sin 2x \end{vmatrix}$$

$$f(x) = (2 + \sin 2x) \begin{vmatrix} 1 & \cos^2 x & \sin 2x \\ 1 & 1 + \cos^2 x & \sin 2x \\ 1 & \cos^2 x & 1 + \sin 2x \end{vmatrix}$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_1$$

$$f(x) = 2 + \sin 2x \begin{vmatrix} 1 & \cos^2 x & \sin 2x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$= (2 + \sin 2x)(1) = 2 + \sin 2x$$

$$= \sin 2x \in \left[\frac{\sqrt{3}}{2}, 1 \right]$$

$$\text{Hence } 2 + \sin 2x \in \left[2 + \frac{\sqrt{3}}{2}, 3 \right]$$

78. Let $f(x) = 2x + \tan^{-1}x$ and $g(x) = \log_e(\sqrt{1+x^2} + x)$, $x \in [0, 3]$. Then

(1) There exists $x \in [0, 3]$ such that $f'(x) < g'(x)$

(2) $\max f(x) > \max g(x)$

(3) There exist $0 < x_1 < x_2 < 3$ such that $f(x) < g(x)$, $\forall x \in (x_1, x_2)$

(4) $\min f'(x) = 1 + \max g'(x)$

Official Ans. by NTA (2)

Ans. (2)

Sol.

$$f(x) = 2x + \tan^{-1}x \text{ and } g(x) = \ln(\sqrt{1+x^2} + x)$$

and $x \in [0, 3]$

$$g'(x) = \frac{1}{\sqrt{1+x^2}}$$

Now, $0 \leq x \leq 3$

$$0 \leq x^2 \leq 9$$

$$1 \leq 1+x^2 \leq 10$$

$$\text{So, } 2 + \frac{1}{10} \leq f'(x) \leq 3$$

$$\frac{21}{10} \leq f'(x) \leq 3 \text{ and } \frac{1}{\sqrt{10}} \leq g'(x) \leq 1$$

option (4) is incorrect

From above, $g'(x) < f'(x) \forall x \in [0, 3]$

Option (1) is incorrect.

$f'(x)$ & $g'(x)$ both positive so $f(x)$ & $g(x)$ both are increasing

So, $\max(f(x))$ at $x=3$ is $6 + \tan^{-1}3$

$\max(g(x))$ at $x=3$ is $\ln(3 + \sqrt{10})$

And $6 + \tan^{-1}3 > \ln(3 + \sqrt{10})$

Option (2) is correct

79. The mean and variance of 5 observations are 5 and 8 respectively. If 3 observations are 1, 3, 5, then the sum of cubes of the remaining two observations is

- (1) 1072 (2) 1792
(3) 1216 (4) 1456

Official Ans. by NTA (1)

Ans. (1)

Sol. $\frac{1+3+5+a+b}{5} = 5$

$a + b = 16 \dots\dots(1)$

$\sigma^2 = \frac{\sum x_i^2}{5} - \left(\frac{\sum x}{5}\right)^2$

$8 = \frac{1^2 + 3^2 + 5^2 + a^2 + b^2}{5} - 25$

$a^2 + b^2 = 130 \dots\dots(2)$

by (1), (2)

$a = 7, b = 9$

or $a = 9, b = 7$

80. The area enclosed by the closed curve C given by the differential equation $\frac{dy}{dx} + \frac{x+a}{y-2} = 0, y(1) = 0$

is 4π .

Let P and Q be the points of intersection of the curve C and the y-axis. If normals at P and Q on the curve C intersect x-axis at points R and S respectively, then the length of the line segment RS is

- (1) $2\sqrt{3}$ (2) $\frac{2\sqrt{3}}{3}$
(3) 2 (4) $\frac{4\sqrt{3}}{3}$

Official Ans. by NTA (4)

Ans. (4)

Sol. $\frac{dy}{dx} + \frac{x+a}{y-2} = 0$

$\frac{dy}{dx} = \frac{x+a}{2-y}$

$(2-y) dy = (x+a) dx$

$2y \frac{-y}{2} = \frac{x^2}{2} + ax + c$

$a + c = -\frac{1}{2}$ as $y(1) = 0$

$X^2 + y^2 + 2ax - 4y - 1 - 2a = 0$

$\pi r^2 = 4\pi$

$r^2 = 4$

$4 = \sqrt{a^2 + 4 + 1 + 2a}$

$(a + 1)^2 = 0$

$P, Q = (0, 2 \pm \sqrt{3})$

Equation of normal at P, Q are $y - 2 = \sqrt{3}(x - 1)$

$y - 2 = -\sqrt{3}(x - 1)$

$R = \left(1 - \frac{2}{\sqrt{3}}, 0\right)$

$S = \left(1 + \frac{2}{\sqrt{3}}, 0\right)$

$RS = \frac{4}{\sqrt{3}} = 4 \frac{\sqrt{3}}{3}$

SECTION-B

81. Let $a_1 = 8, a_2, a_3, \dots, a_n$ be an A.P. If the sum of its first four terms is 50 and the sum of its last four terms is 170, then the product of its middle two terms is _____.

Official Ans. by NTA (754)

Ans. (754)

Sol. $a_1 + a_2 + a_3 + a_4 = 50$

$\Rightarrow 32 + 6d = 50$

$\Rightarrow d = 3$

and, $a_{n-3} + a_{n-2} + a_{n-1} + a_n = 170$

$\Rightarrow 32 + (4n - 10) \cdot 3 = 170$

$\Rightarrow n = 14$

$a_7 = 26, a_8 = 29$

$\Rightarrow a_7 \cdot a_8 = 754$

82. $A(2, 6, 2), B(-4, 0, \lambda), C(2, 3, -1)$ and $D(4, 5, 0), |\lambda| \leq 5$ are the vertices of a quadrilateral ABCD. If its area is 18 square units, then $5 - 6\lambda$ is equal to _____.

Official Ans. by NTA (11)

Ans. (11)

Sol. A(2, 6, 2) B(-4, 0, λ), C(2, 3, -1) D(4, 5, 0)

$$\text{Area} = \frac{1}{2} |\overrightarrow{BD} \times \overrightarrow{AC}| = 18$$

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 8 & 5 & -\lambda \\ 3 & -3 & -3 \end{vmatrix}$$

$$= (3\lambda + 15)\hat{i} - j(-24) + k(-24)$$

$$\overrightarrow{AC} \times \overrightarrow{BD} = (3\lambda + 15)\hat{i} + 24j - 24k$$

$$= \sqrt{(3\lambda + 15)^2 + (24)^2 + (24)^2} = 36$$

$$= \lambda^2 + 10\lambda + 9 = 0$$

$$= \lambda = -1, -9$$

$$|\lambda| \leq 5 \Rightarrow \lambda = -1$$

$$5 - 6\lambda = 5 - 6(-1) = 11$$

83. The number of 3-digit numbers, that are divisible by either 2 or 3 but not divisible by 7 is _____.

Official Ans. by NTA (514)

Ans. (514)

Sol. Divisible by 2 → 450

Divisible by 3 → 300

Divisible by 7 → 128

Divisible by 2 & 7 → 64

Divisible by 3 & 7 → 43

Divisible by 2 & 3 → 150

Divisible by 2, 3 & 7 → 21

$$\therefore \text{Total numbers} = 450 + 300 - 150 - 64 - 43 + 21 = 514$$

84. The remainder when $19^{200} + 23^{200}$ is divided by 49, is _____.

Official Ans. by NTA (29)

Ans. (29)

Sol. $(21 + 2)^{200} + (21 - 2)^{200}$

$$2[{}^{100}C_0 21^{200} + 200C_2 21^{198} \cdot 2^2 + \dots + {}^{200}C_{198} 21^2 \cdot 2^{198} + 2^{200}]$$

$$\Rightarrow 2[49 I_1 + 2^{200}] = 49I_1 + 2^{201}$$

$$\text{Now, } 2^{201} = (8)^{67} = (1 + 7)^{67} = 49I_2 + {}^{67}C_0 {}^{67}C_1 \cdot 7 = 49I_2 + 470 = 49I_2 + 49 \times 9 + 29$$

\therefore Remainder is 29

85. If

$$\int_0^1 (x^{21} + x^{14} + x^7)(2x^{14} + 3x^7 + 6)^{1/7} dx = \frac{1}{l} (11)^{m/n}$$

where $l, m, n \in \mathbb{N}$, m and n are coprime then $l + m + n$ is equal to _____.

Official Ans. by NTA (63)

Ans. (63)

Sol. $\int (x^{20} + x^{13} + x^6)(2x^{21} + 3x^{14} + 6x^7)^{1/7} dx$

$$2x^{21} + 3x^{14} + 6x^7 = t$$

$$42(x^{20} + x^{13} + x^6) dx = dt$$

$$\frac{1}{42} \int_0^{11} t^{1/7} dt = \left(\frac{t^{8/7}}{8/7} \times \frac{1}{42} \right)_0^{11}$$

$$= \frac{1}{48} \left(t^{8/7} \right)_0^{11} = \frac{1}{48} (11)^{8/7}$$

$$l = 48, m = 8, n = 7$$

$$l + m + n = 63$$

86. If $f(x) = x^2 + g'(1)x + g''(2)$ and

$$g(x) = f(1)x^2 + xf'(x) + f''(x),$$

then the value of $f(4) - g(4)$ is equal to _____.

Official Ans. by NTA (14)

Ans. (14)

Sol. $f(x) = x^2 + g'(1)x + g''(2)$

$$f'(x) = 2x + g'(1)$$

$$f''(x) = 2$$

$$g(x) = f(1)x^2 + x[2x + g'(1)] + 2$$

$$g'(x) = 2f(1)x + 4x + g'(1)$$

$$g''(x) = 2f(1) + 4$$

$$g''(x) = 0$$

$$2f(1) + 4 = 0$$

$$f(1) = -2$$

$$-2 = 1 + g'(1) = g'(1) = -3$$

So, $f'(x) = 2x - 3$

$$f(x) = x^2 - 3x + c$$

$$c = 0$$

$$f(x) = x^2 - 3x$$

$$g(x) = -3x + 2$$

$$f(4) - g(4) = 14$$

87. Let $\vec{v} = \alpha\hat{i} + 2j - 3k$, $\vec{w} = 2\alpha\hat{i} + j - k$, and \vec{u} vector such that $|\vec{u}| = \alpha > 0$. If the minimum value of the scalar triple product $[\vec{u}\vec{v}\vec{w}]$ is $-\alpha\sqrt{3401}$, and $|\vec{u}\cdot\hat{i}|^2 = \frac{m}{n}$ where m and n are coprime natural numbers, then m + n is equal to _____.

Official Ans. by NTA (3501)

Ans. (3501)

Sol. $[\vec{u}\vec{v}\vec{w}] = \vec{u} \cdot (\vec{v} \times \vec{w})$

$$\min. (|\vec{u}| |\vec{v} \times \vec{w}| \cos \theta) = -\alpha\sqrt{3401}$$

$$\Rightarrow \cos \theta = -1$$

$$|\vec{u}| = \alpha \text{ (Given)}$$

$$|\vec{v} \times \vec{w}| = \sqrt{3401}$$

$$\vec{v} \times \vec{w} = \begin{vmatrix} \hat{i} & j & k \\ \alpha & 2 & -3 \\ 2\alpha & 1 & -1 \end{vmatrix}$$

$$\vec{v} \times \vec{w} = \hat{i} - 5\alpha j - 3\alpha k$$

$$|\vec{v} \times \vec{w}| = \sqrt{1 + 25\alpha^2 + 9\alpha^2} = \sqrt{3401}$$

$$34\alpha^2 = 3400$$

$$\alpha^2 = 100$$

$$\alpha = 10 \quad (\text{as } \alpha > 0)$$

So $u = \lambda(i \quad k)$

$$\vec{u} = \sqrt{\lambda^2 + 25\alpha^2\lambda^2 + 9\alpha^2\lambda^2}$$

$$\alpha^2 = \lambda^2(1 + 25\alpha^2 + 9\alpha^2)$$

$$100 = \lambda^2(1 + 34 \times 100)$$

$$\lambda^2 = \frac{100}{3401} = \frac{m}{n}$$

88. The number of words, with or without meaning, that can be formed using all the letters of the word ASSASSINATION so that the vowels occur together, is _____.

Official Ans. by NTA (50400)

Ans. (50400)

ol. Vowels : A,A,A,I,I,O

Consonants : S,S,S,S,N,N,T

Total number of ways in which vowels come together

$$= \frac{|8}{|4|2} \times \frac{|6}{|3|2} = 50400$$

89. Let A be the area bounded by the curve $y = x|x - 3|$, the x-axis and the ordinates $x = -1$ and $x = 2$. Then 12A is equal to _____.

Official Ans. by NTA (62)

Ans. (62)

Sol. $A = \int_{-1}^0 (x^2 - 3x) dx + \int_0^2 (3x - x^2) dx$

$$\Rightarrow A = \frac{x^3}{3} - \frac{3x^2}{2} \Big|_{-1}^0 + \frac{3x^2}{2} - \frac{x^3}{3} \Big|_0^2$$

$$\Rightarrow A = \frac{11}{6} + \frac{10}{3} = \frac{31}{6}$$

$$\therefore 12A = 62$$

90. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function such that $f'(x) + f(x) = \int_0^2 f(t) dt$. If $f(0) = e^{-2}$, then $2f(0) - f(2)$ is equal to _____.

Official Ans. by NTA (1)

Ans. (1)

Sol. $\frac{dy}{dx} + y = k$

$$y \cdot e^x = k \cdot e^x + c$$

$$f(0) = e^{-2}$$

$$\Rightarrow c = e^{-2} - k$$

$$\therefore y = k + (e^{-2} - k)e^{-x}$$

$$\text{now } k = \int_0^2 (k + (e^{-2} - k)e^{-x}) dx$$

$$\Rightarrow k = e^{-2} - 1$$

$$\therefore y = (e^{-2} - 1) + e^{-x}$$

$$f(2) = 2e^{-2} - 1, f(0) = e^{-2}$$

$$2f(0) - f(2) = 1$$