

FINAL JEE-MAIN EXAMINATION – JANUARY, 2023

(Held On Wednesday 1st February, 2023)


TIME : 9 : 00 AM to 12 : 00 NOON

SECTION-A


31. Which of the following represents the lattice structure of $A_{0.95}O$ containing A^{2+} , A^{3+} and O^{2-} ions?

○ A^{2+} ○ A^{3+} ○ O^{2-}


A.



B.



C.



(1) B and C only

(2) B only

(3) A and B only

(4) A only






Official Ans. by NTA (4)

Ans. (4)

Sol. Applying electrical neutrality principle in metal deficiency defect.

3 A^{2+} are replaced by 2 A^{3+} , thus one vacant site per pair of A^{3+} is created

32. The correct representation in six membered pyranose form for the following sugar [X] is



Official Ans. by NTA (2)

Ans. (2)

Sol. By Haworth structure of mannose.


33. Highest oxidation state of Mn is exhibited in Mn_2O_7 . The correct statements about Mn_2O_7 are
 (A) Mn is tetrahedrally surrounded by oxygen atoms
 (B) Mn is octahedrally surrounded by oxygen atoms
 (C) Contains Mn-O-Mn bridge
 (D) Contains Mn-Mn bond.

Choose the correct answer from the options given below


- (1) A and C only (2) A and D only
 (3) B and D only (4) B and C only

Official Ans. by NTA (1)

Ans. (1)



34. Decreasing order of dehydration of the following alcohols is



- (1) $a > d > b > c$ (2) $b > d > c > a$
 (3) $b > a > d > c$ (4) $d > b > c > a$

Official Ans. by NTA (2)

Ans. (2)

- Sol. Dehydration of alcohol is directly proportional to the stability of carbocation.

35. Given below are two statements: One is labelled as **Assertion A** and the other is labelled as **Reason R**.

Assertion A: Amongst He, Ne, Ar and Kr;

1 g of activated charcoal adsorbs more of Kr.

Reason R : The critical volume V_c ($cm^3 mol^{-1}$) and critical pressure P_c (atm) is highest for Krypton but the compressibility factor at critical point Z_c is lowest for Krypton.

In the light of the above statements, choose the **correct** answer from the options given below.

- (1) A is true but R is false
 (2) A is false but R is true
 (3) Both A and R are true but R is NOT the correct explanation of A
 (4) Both A and R are true and R is the correct explanation A


Official Ans. by NTA (1)

Ans. (1)





- Sol. Adsorption \propto vanderwaal attraction forces

$$Z_c = \frac{3}{8} \text{ for all real gases}$$

36. In the following reaction, 'A' is



'A' Major product.




Official Ans. by NTA (2)

Ans. (2)

- Sol. Initially lone pair electron of $-NH_2$ attack on electrophilic carbon, after then lone pair electron of

oxygen attacks leading to formation of cyclic compound.



37. Match List I with List II

List-I	List-II
(A) Tranquilizers	(I) Anti blood clotting
(B) Aspirin	(II) Salvarsan
(C) Antibiotic	(III) Antidepressant drugs
(D) Antiseptic	(IV) Soframicine

Choose the correct answer from the options given below:

- (1) (A) – IV, (B) – II, (C) – I, (D) – III
- (2) (A) – II, (B) – I, (C) – III, (D) – IV
- (3) (A) – III, (B) – I, (C) – II, (D) – IV
- (4) (A) – II, (B) – IV, (C) – I, (D) – III

Official Ans. by NTA (3)

Ans. (3)

Sol. NCERT (Chemistry in every day life)

38. Given below are two statements:

Statement I: Chlorine can easily combine with oxygen to form oxides: and the product has a tendency to explode.

Statement II: Chemical reactivity of an element can be determined by its reaction with oxygen and halogens.

In the light of the above statements, choose the **correct** answer from the options given below.


- (1) Both the statements I and II are true
- (2) Statement I is true but Statement II is false
- (3) Statement I is false but Statement II is true
- (4) Both the Statements I and II are false

Official Ans. by NTA (1)

Ans. (1)

Sol. Chlorine oxides, Cl_2O , ClO_2 , Cl_2O_6 and Cl_2O_7 are highly reactive oxidising agents and tend to explode.

39. Resonance in carbonate ion (CO_3^{2-}) is



Which of the following is true?





- (1) It is possible to identify each structure individually by some physical or chemical method.
- (2) All these structures are in dynamic equilibrium with each other.
- (3) Each structure exists for equal amount of time.
- (4) CO_3^{2-} has a single structure i.e., resonance hybrid of the above three structures.

Official Ans. by NTA (4)

Ans. (4)

Sol. Resonating structure are hypothetical and resonance hybrid is real structure which is weighted average of all the resonating structures.

40. Identify the incorrect option from the following:



Official Ans. by NTA (2)

Ans. (2)

Sol. In alcoholic KOH, elimination reaction takes place.

41. A solution of FeCl₃ when treated with K₄[Fe(CN)₆] gives a prussian blue precipitate due to the formation of

- (1) K[Fe₂(CN)₆]
- (2) Fe[Fe(CN)₆]
- (3) Fe₃[Fe(CN)₆]₂
- (4) Fe₄[Fe(CN)₆]₃

Official Ans. by NTA (4)

Ans. (4)

Sol. Formation of Prussian blue complex takes place.

42. Which of the following are the example of double salt?

- (A) FeSO₄·(NH₄)₂SO₄·6H₂O
- (B) CuSO₄·4NH₃·H₂O
- (C) K₂SO₄·Al₂(SO₄)₃·24H₂O
- (D) Fe(CN)₂·4KCN

Choose the correct answer.

- (1) A and C only
- (2) A and B only

(3) A, B and D only

(4) B and D only

Official Ans. by NTA (1)

Ans. (1)

Sol. Double salt contain's two or more types of salts. CuSO₄·4NH₃·H₂O and Fe(CN)₂·4KCN are complex compounds.

43. Which of the following complex will show largest splitting of d-orbitals?

- (1) [Fe(C₂O₄)₃]³⁻
- (2) [FeF₆]³⁻
- (3) [Fe(CN)₆]³⁻
- (4) [Fe(NH₃)₆]³⁺

Official Ans. by NTA (3)

Ans. (3)

Sol. $\bar{C}N$ is a strong field ligand so maximum splitting in d orbitals take place.

44. How can photochemical smog be controlled?

- (1) By using tall chimneys
- (2) By complete combustion of fuel
- (3) By using catalytic converters in the automobiles/industry
- (4) By using catalyst

Official Ans. by NTA (3)

Ans. (3)

Sol. NCERT (Environmental chemistry)

45. Match List I with List II

- | | |
|------------------------|---|
| (A) Slaked lime | (I) NaOH |
| (B) Dead burnt plaster | (II) Ca(OH) ₂ |
| (C) Caustic soda | (III) Na ₂ CO ₃ ·10H ₂ O |
| (D) Washing soda | (IV) CaSO ₄ |

Choose the correct answer form the options given below:

- (1) (A) – I, (B) – IV, (C) – II, (D) – III
- (2) (A) – III, (B) – IV, (C) – II, (D) – I
- (3) (A) – II, (B) – IV, (C) – I, (D) – III
- (4) (A) – III, (B) – II, (C) – IV, (D) – I

Official Ans. by NTA (3)
Ans. (3)
Sol. From S-block NCERT

46. Choose the correct statement(s):

- A. Beryllium oxide is purely acidic in nature.
 B. Beryllium carbonate is kept in the atmosphere of CO₂.
 C. Beryllium sulphate is readily soluble in water.
 D. Beryllium shows anomalous behavior.

Choose the correct answer from the options given below:

- (1) A, B and C only
 (2) B, C and D only
 (3) A and B only
 (4) A only

Official Ans. by NTA (2)
 Ans. (2)
Sol. A. Beryllium oxide is amphoteric in nature.

 B. Beryllium carbonate is kept in the atmosphere of CO₂ because it is thermally less stable.

C. Beryllium sulphate is readily soluble in water due to high degree of hydration.

 D. Beryllium shows anomalous behaviour due to small size, high ionization energy and high value of ϕ (polarising power).

47. Given below are two statements: one is labelled as

Assertion A and the other is labelled as **Reason R**

Assertion A: In an Ellingham diagram, the oxidation of carbon to carbon monoxide shows a negative slope with respect to temperature.

Reason R: CO tends to get decomposed at higher temperature.


In the light of the above statements, choose the correct answer from the options given below

- (1) Both A and R are correct and R is the correct explanation of A
 (2) A is not correct but R is correct
 (3) Both A and R are correct but R is NOT the correct explanation of A

(4) A is correct but R is not correct

Official Ans. by NTA (4)
Ans. (4)
Sol. $2C(s) + O_2(g) \rightarrow 2CO(g)$
 $\Delta_r S^\circ$ is +ve, $\Delta_r G^\circ = \Delta_r H^\circ - T\Delta_r S^\circ$; thus slope is negative

 As temperature increases $\Delta_r G^\circ$ becomes more negative thus it has lower tendency to get decomposed.

48. But-2-yne is reacted separately with one mole of Hydrogen as shown below:


Identify the incorrect statements from the options given below:

- A. A is more soluble than B.
 B. The boiling point & melting point of A are higher and lower than B respectively.
 C. A is more polar than B because dipole moment of A is zero.
 D. Br₂ adds easily to B than A.

- (1) B and C only
 (2) B, C and D only
 (3) A, C and D only
 (4) A and B only

Official Ans. by NTA (2)
 Ans. (Bonus)
Sol. Incorrect statements are C and D only, correct choice is not available.

49. Given below are two statements: one is labelled as **Assertion A** and the other is labelled as **Reason R**
Assertion A: Hydrogen is an environment friendly fuel.

Reason R: Atomic number of hydrogen is 1 and it is a very light element.

In the light of the above statements, choose the correct answer from the options given below

- (1) A is true but R is false
 (2) Both A and R are true but R is NOT the correct explanation of A
 (3) A is false but R is true
 (4) Both A and R are true and R is the correct explanation of A

Official Ans. by NTA (2)

Ans. (2)

Sol. No pollution occurs by combustion of hydrogen and very low density of hydrogen.

50. Match List I and List II

List I	List II
Test	Functional group / Class of Compound
(A) Molisch's Test	(I) Peptide
(B) Biuret Test	(II) Carbohydrate
(C) Carbylamine Test	(III) Primary amine
(D) Schiff's Test	(IV) Aldehyde

Choose the correct answer from the options given below:

- (1) (A) – I, (B) – II, (C) – III, (D) – IV
 (2) (A) – III, (B) – IV, (C) – I, (D) – II
 (3) (A) – II, (B) – I, (C) – III, (D) – IV
 (4) (A) – III, (B) – IV, (C) – II, (D) – I

Official Ans. by NTA (3)

Ans. (3)

Sol.

List I	List II
Test	Functional group / Class of Compound
(A) Molisch's Test	(II) Carbohydrate
(B) Biuret Test	(I) Peptide
(C) Carbylamine Test	(III) Primary amine
(D) Schiff's Test	(IV) Aldehyde

SECTION-B

51. The density of 3 M solution of NaCl is 1.0 g mL⁻¹. Molality of the solution is _____ × 10⁻² m. (Nearest integer).

Given: Molar mass of Na and Cl is 23 and 35.5 g mol⁻¹ respectively.

Official Ans. by NTA (364)

Ans. (364)

$$\begin{aligned} \text{Sol. } m &= \frac{1000 \times M}{1000 \times d - M \times M.W \text{ of solute}} \\ &= \frac{1000 \times 3}{1000 \times 1 - (3 \times 58.5)} = 3.64 \\ &= 364 \times 10^{-2} \end{aligned}$$

52. Electrons in a cathode ray tube have been emitted with a velocity of 1000 ms⁻¹. The number of following statements which is/are true about the emitted radiation is _____.

Given : $h = 6 \times 10^{-34}$ Js, $m_e = 9 \times 10^{-31}$ kg.

- (A) The deBroglie wavelength of the electron emitted is 666.67nm.
 (B) The characteristic of electrons emitted depend upon the material of the electrodes of the cathode ray tube.
 (C) The cathode rays start from cathode and move towards anode.
 (D) The nature of the emitted electrons depends on the nature of the gas present in cathode ray tube.

Official Ans. by NTA (2)

Ans. (2)

Sol. (A) $V_e = 1000$ m/s ; $h = 6 \times 10^{-34}$ Js ;

$$m_e = 9 \times 10^{-31} \text{ kg}$$

$$\lambda = \frac{h}{mv} = \frac{6 \times 10^{-34}}{9 \times 10^{-31} \times 1000} = 666.67 \times 10^{-9} \text{ m}$$

$$= 666.67 \text{ nm}$$

(B) The characteristic of electrons emitted is independent of the material of the electrodes of the cathode ray tube.

(C) The cathode rays start from cathode and move towards anode.

(D) The nature of the emitted electrons is independent on the nature of the gas present in cathode ray tube.

53. Sum of oxidation states of bromine in bromic acid and perbromic acid is _____ .

Official Ans. by NTA (12)

Ans. (12)

Sol. HBrO_3 (Bromic acid)

Ox. State of Br = +5

HBrO_4 (per bromic acid)

Ox. State of Br = +7

Sum of Ox. State = 12

54. At what pH, given half cell MnO_4^- (0.1 M) | Mn^{2+} (0.001 M) will have electrode potential of 1.282 V? _____ (Nearest Integer)

$$\text{Given } E_{\text{MnO}_4^-/\text{Mn}^{2+}}^{\circ} = 1.54 \text{ V, } \frac{2.303RT}{F} = 0.059 \text{ V}$$

Official Ans. by NTA (3)

Ans. (3)

Sol. $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$

$$E = E^{\circ} - \frac{0.059}{5} \log \frac{[\text{Mn}^{2+}]}{[\text{MnO}_4^-][\text{H}^+]^8}$$

$$1.282 = 1.54 - \frac{0.059}{5} \log \frac{10^{-3}}{10^{-1} \times [\text{H}^+]^8}$$

$$\frac{0.2}{0.059} = \log \frac{10^{-2}}{[\text{H}^+]^8}$$

$$\Rightarrow 21.86 = -2 + 8\text{pH}$$

$$\therefore \text{pH} = 2.98$$

$$\approx 3$$

55. Number of isomeric compounds with molecular formula $\text{C}_9\text{H}_{10}\text{O}$ which (i) do not dissolve in NaOH (ii) do not dissolve in HCl. (iii) do not give orange precipitate with 2, 4 - DNP (iv) on hydrogenation give identical compound with molecular formula $\text{C}_9\text{H}_{12}\text{O}$ is _____ .

Official Ans. by NTA (2)

Ans. (2)

Sol. As per the language of given question, the best possible isomeric structure is $\text{Ph}-\text{CH}=\text{CH}-\text{O}-\text{CH}_3$ (cis and trans). So, the answer is 2.

56. (i) $\text{X}(\text{g}) \rightleftharpoons \text{Y}(\text{g}) + \text{Z}(\text{g})$ $K_{p1} = 3$

$$(ii) \text{A}(\text{g}) \rightleftharpoons 2\text{B}(\text{g}) \quad K_{p2} = 1$$

If the degree of dissociation and initial concentration of both the reactants $\text{X}(\text{g})$ and $\text{A}(\text{g})$ are equal, then the ratio of the total pressure at equilibrium $\left(\frac{P_1}{P_2}\right)$ is equal to $x : 1$. The value of x is _____ (Nearest integer)

Official Ans. by NTA (12)

Ans. (12)

Sol. $\text{x}(\text{g}) \rightleftharpoons \text{y}(\text{g}) + \text{z}(\text{g}) \quad k_{p1} = 3$

Initial moles n - -

at equilibrium n - α n α n α n

$$k_{p1} = \frac{\left(\frac{\alpha}{1+\alpha} \times p_1\right)^2}{\frac{1-\alpha}{1+\alpha} p_1}$$

$$3 = \frac{\alpha^2 \times p_1}{1-\alpha^2}$$

$\text{A}(\text{g}) \rightleftharpoons 2\text{B}(\text{g}) \quad k_{p2} = 1$

Initial mole n -

at equilibrium x - α n 2 α n $p_{\text{total}} = p_2$

$$k_{p2} = \frac{\left(\frac{2\alpha}{1+\alpha} \times p_2\right)^2}{\frac{1-\alpha}{1+\alpha} \times p_2}$$

$$1 = \frac{4\alpha^2 \times p_2}{1-\alpha^2}$$


$$\frac{k_{p_1}}{k_{p_2}} = \frac{p_1}{4p_2}$$

$$\frac{3}{1} = \frac{p_1}{4p_2}$$

$$\therefore p_1 : p_2 = 12 : 1$$

$$x = 12$$




57. The total number of chiral compound/s from the following is _____.



Official Ans. by NTA (2)

Ans. (2)

Sol.



58. A and B are two substances undergoing radioactive decay in a container. The half life of A is 15 min and that of B is 5 min. If the initial concentration of B is 4 times that of A and they both start decaying at the same time, how much time will it take for the concentration of both of them to be same? _____ min.

Official Ans. by NTA (15)

Ans. (15)

Sol. $[A]_t = [A]_0 e^{-kt}$

For A : Let $[A]_t$ be y and $[A]_0$ be x ; $k = \frac{\ln 2}{t_{1/2}} =$

$$\frac{\ln 2}{15 \text{ min}}$$

$$y = x e^{-kt}$$

$$= xe^{-\left(\frac{\ln 2}{15}\right)t}$$

For B : $[B]_t = [B]_0 e^{-kt}$

Let $[B]_t = y$; $[B]_0 = 4x$; $k = \frac{\ln 2}{t_{1/2}} = \frac{\ln 2}{5 \text{ min}}$

$$y = 4xe^{-\left(\frac{\ln 2}{5}\right)t}$$

$$\Rightarrow xe^{-\left(\frac{\ln 2}{15}\right)t} = 4xe^{-\left(\frac{\ln 2}{5}\right)t}$$


$$e^{t\left(\frac{\ln 2}{5} - \frac{\ln 2}{15}\right)} = 4$$

$$t \times \left[\frac{\ln 2}{5} - \frac{\ln 2}{15} \right] = \ln 4$$

$$t \times \ln 2 \left[\frac{1}{5} - \frac{1}{15} \right] = 2 \ln 2$$

$$t = 15 \text{ min}$$

59. At 25°C, the enthalpy of the following processes are given:






What would be the value of X for the following reaction? _____ (Nearest integer)



Official Ans. by NTA (499)

Ans. (499)



60. 25 mL of an aqueous solution of KCl was found to require 20 mL of 1 M AgNO₃ solution when titrated using K₂CrO₄ as an indicator. What is the depression in freezing point of KCl solution of the given concentration? _____ (Nearest integer).

(Given : K_f = 2.0 K kg mol⁻¹)

Assume


1) 100% ionization and

2) density of the aqueous solution as 1 g mL⁻¹

Official Ans. by NTA (3)

Ans. (3)

Sol.



At equivalence point,

mmole of KCl = mmole of AgNO₃

= 20 mmole

Volume of solution = 25 ml

Mass of solution = 25 gm

Mass of solvent

= 25 – mass of solute

= 25 – [20 × 10⁻³ × 74.5]

= 23.51 gm

$$\text{Molality of KCl} = \frac{\text{mole of KCl}}{\text{mass of solvent in kg}}$$

$$= \frac{20 \times 10^{-3}}{23.51 \times 10^{-3}} = 0.85$$

i of KCl = 2 (100% ionisation)

$$\Delta T_f = i \times K_f \times m$$

$$= 2 \times 2 \times 0.85$$

$$= 3.4$$

$$\approx 3$$