


5. The variation of molar conductivity with concentration of an electrolyte (X) in aqueous solution is shown in the given figure.



The electrolyte X is :

- (1) CH_3COOH (2) KNO_3
 (3) HCl (4) NaCl

Official Ans. by NTA (1)

Sol. Its a weak electrolyte hence : CH_3COOH

6. The one that is NOT suitable for the removal of permanent hardness of water is :

- (1) Treatment with sodium carbonate
 (2) Calgon's method
 (3) Clark's method
 (4) Ion-exchange method


Official Ans. by NTA (3)

Sol. Temporary hardness of water is removed by clark method and boiling. While permanent hardness of water is removed by treatment with sodium carbonate (Na_2CO_3), calgons method and ion-exchange method


7. The correct statement about probability density (except at infinite distance from nucleus) is :

- (1) It can be negative for 2p orbital
 (2) It can be zero for 3p orbital
 (3) It can be zero for 1s orbital
 (4) It can never be zero for 2s orbital

Official Ans. by NTA (2)




8. The increasing order of boiling points of the following compounds is :



- (1) $\text{I} < \text{IV} < \text{III} < \text{II}$
 (2) $\text{IV} < \text{I} < \text{II} < \text{III}$
 (3) $\text{I} < \text{III} < \text{IV} < \text{II}$
 (4) $\text{III} < \text{I} < \text{II} < \text{IV}$

Official Ans. by NTA (1)


Sol.




$$\text{BP} \propto \text{dipole moment } (\mu)$$

Alter


Increasing order of boiling point is :




\Rightarrow Shows hydrogen bonding from $-\text{O}-\text{H}$ group only



\Rightarrow Shows strongest hydrogen bonding from both sides of $-\text{OH}$ group as well as $-\text{NO}_2$ group.



⇒ Shows stronger hydrogen from both side of -OH group as well as -NH₂ group.



⇒ Shows stronger hydrogen bonding from one side -OH-group and another side of -OCH₃ group shows only dipole-dipole interaction.


⇒ Hence correct order of boiling point is:

(I) < (IV) < (III) < (II)






9. The compound that has the largest H-M-H bond angle (M=N, O, S, C), is :

- (1) H₂O (2) CH₄
 (3) NH₃ (4) H₂S

Official Ans. by NTA (2)




10. Among the following compounds, geometrical isomerism is exhibited by :





11. Which one of the following polymers is not obtained by condensation polymerisation?


- (1) Buna - N (2) Bakelite
 (3) Nylon 6 (4) Nylon 6, 6

Official Ans. by NTA (1)



12. The final major product of the following reaction is :






13. Hydrogen peroxide, in the pure state, is :

- (1) non-planar and almost colorless
- (2) linear and almost colorless
- (3) planar and blue in color
- (4) linear and blue in color

Official Ans. by NTA (1)



hydrogen peroxide, in the pure state, is non-planar and almost colourless (very pale blue) liquid.

14. Boron and silicon of very high purity can be obtained through :

- (1) vapour phase refining
- (2) electrolytic refining
- (3) liquation
- (4) zone refining

Official Ans. by NTA (4)

Sol. "Boron" and "Silicon" of very high purity can be obtained through :-
zone refining method only.

While other methods are used for other metals/elements i.e.


- (i) Vapour phase refining
- (ii) electrolytic refining

15. Lattice enthalpy and enthalpy of solution of NaCl are 788 kJ mol⁻¹ and 4 kJ mol⁻¹, respectively.

The hydration enthalpy of NaCl is :

- (1) -780 kJ mol⁻¹
- (2) -784 kJ mol⁻¹
- (3) 780 kJ mol⁻¹
- (4) 784 kJ mol⁻¹

Official Ans. by NTA (2)




$$4 = 788 + \Delta H$$

$$\Delta H = -784 \text{ kJ}$$

16. Reaction of ammonia with excess Cl₂ gives :

- (1) NH₄Cl and N₂
- (2) NCl₃ and NH₄Cl
- (3) NH₄Cl and HCl
- (4) NCl₃ and HCl

Official Ans. by NTA (4)



17. The correct order of the ionic radii of O²⁻, N³⁻, F⁻, Mg²⁺, Na⁺ and Al³⁺ is :

- (1) Al³⁺ < Na⁺ < Mg²⁺ < O²⁻ < F⁻ < N³⁻
- (2) N³⁻ < O²⁻ < F⁻ < Na⁺ < Mg²⁺ < Al³⁺
- (3) Al³⁺ < Mg²⁺ < Na⁺ < F⁻ < O²⁻ < N³⁻
- (4) N³⁻ < F⁻ < O²⁻ < Mg²⁺ < Na⁺ < Al³⁺

Official Ans. by NTA (3)

Sol. Correct order of size for isoelectronic species.

$$\text{Al}^{3+} < \text{Mg}^{2+} < \text{Na}^+ < \text{F}^- < \text{O}^{2-} < \text{N}^{3-}$$


18. Consider the complex ions,

trans-[Co(en)₂Cl₂]⁺ (A) and

cis-[Co(en)₂Cl₂]⁺ (B). The correct statement regarding them is :


- (1) both (A) and (B) can be optically active
- (2) both (A) and (B) cannot be optically active
- (3) (A) can be optically active, but (B) cannot be optically active
- (4) (A) cannot be optically active, but (B) can be optically active

Sol. (A) $trans-[Co(en)_2Cl_2]^+$







\Rightarrow (A) is trans form and shows plane of symmetry which is optically inactive (not optically active)

(B) $cis-[Co(en)_2Cl_2]^+$




\Rightarrow (B) is cis form and does not show plane of symmetry, hence it is optically active.

19. Adsorption of a gas follows Freundlich adsorption isotherm. If x is the mass of the gas adsorbed on mass m of the adsorbent, the correct plot of $\frac{x}{m}$ versus p is :



Sol. $\frac{x}{m} = K.P^{1/n}$



20. The major product formed in the following reaction is :




- (1) $CH_3CH_2CH_2C(Br)(CH_3)_2$
- (2) $Br(CH_2)_3CH(CH_3)_2$
- (3) $CH_3CH_2CH(Br)CH(CH_3)_2$
- (4) $CH_3CH(Br)CH_2CH(CH_3)_2$

Official Ans. by NTA (1)

Official Ans. by (4)

Sol. $CH_3-CH=CH-CH-CH_3$




Addition of HBr according to M.R.

21. The number of chiral carbons present in sucrose is _____.


Official Ans. by NTA (9)

Sol.



Total no. of chiral carbon in sucrose = 9

22. For a dimerization reaction,



at 298 K, $\Delta U^\ominus = -20 \text{ kJ mol}^{-1}$, $\Delta S^\ominus = -30 \text{ J K}^{-1} \text{ mol}^{-1}$, then the ΔG^\ominus will be _____ J.

Official Ans. by NTA (-13538.00)

Sol. $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$
 $= (\Delta U^\circ + \Delta n_g RT) - T\Delta S^\circ$
 $= \left[\left\{ -20 + (-1) \frac{8.314}{1000} \times 298 \right\} - \frac{298}{1000} \times (-30) \right] \text{kJ}$
 $= -13.537572 \text{ kJ}$
 $= -13537.57 \text{ Joule}$

23. For a reaction $X + Y \rightleftharpoons 2Z$, 1.0 mol of X, 1.5 mol of Y and 0.5 mol of Z were taken in a 1 L vessel and allowed to react. At equilibrium, the concentration of Z was 1.0 mol L⁻¹. The equilibrium constant of the reaction is

_____ $\frac{x}{15}$. The value of x is _____.

Official Ans. by NTA (16)

Sol.


	X	+	Y	=	2Z	
t = 0	1		1.5		0.5	;
At eq.	0.75		1.25		1	

$$K_{\text{eq.}} = \frac{1^2}{\frac{3}{4} \times \frac{5}{4}} = \frac{16}{15}$$

24. The volume, in mL, of 0.02 M K₂Cr₂O₇ solution required to react with 0.288 g of ferrous oxalate in acidic medium is _____.

(Molar mass of Fe = 56 g mol⁻¹)

Official Ans. by NTA (50.00)




$$\frac{0.02 \times 6 \times V(\text{mL})}{1000} = \frac{0.288}{144} \times 3$$

$$\Rightarrow \boxed{V = 50\text{mL}}$$

25. Considering that $\Delta_0 > P$, the magnetic moment (in BM) of [Ru(H₂O)₆]²⁺ would be _____.

Official Ans. by NTA (00)

Sol. Magnetic moment (in B.M.) of [Ru(H₂O)₆]²⁺ would be; while considering that $\Delta_0 > P$,
 Ru₍₄₄₎; [Kr]4d⁷5s¹ (in ground state)
 \Rightarrow In Ru²⁺ $\Rightarrow 4d^6 \Rightarrow (t_2g)^6(eg)^0$



\Rightarrow Here number of unpaired electrons in

Ru²⁺ = (t₂g)⁶ (eg)⁰ = 0 and Hence

$$\mu_m = \sqrt{n(n+2)} \text{ B.M.} = \boxed{0 \text{ B.M.}}$$