

**FINAL JEE–MAIN EXAMINATION – MARCH, 2021**

(Held On Thursday 18<sup>th</sup> March, 2021) TIME : 3 : 00 PM to 6 : 00 PM

**CHEMISTRY**

**TEST PAPER WITH ANSWER & SOLUTION**


**SECTION-A**

1. The oxidation states of nitrogen in NO, NO<sub>2</sub>, N<sub>2</sub>O and NO<sub>3</sub><sup>-</sup> are in the order of :

- (1) NO<sub>3</sub><sup>-</sup> > NO<sub>2</sub> > NO > N<sub>2</sub>O
- (2) NO<sub>2</sub> > NO<sub>3</sub><sup>-</sup> > NO > N<sub>2</sub>O
- (3) N<sub>2</sub>O > NO<sub>2</sub> > NO > NO<sub>3</sub><sup>-</sup>
- (4) NO > NO<sub>2</sub> > N<sub>2</sub>O > NO<sub>3</sub><sup>-</sup>

**Official Ans. by NTA (1)**

Sol. The oxidation states of Nitrogen in following molecules are as follows



2. In basic medium, H<sub>2</sub>O<sub>2</sub> exhibits which of the following reactions ?

- (A) Mn<sup>2+</sup> → Mn<sup>4+</sup>
- (B) I<sub>2</sub> → I<sup>-</sup>
- (C) PbS → PbSO<sub>4</sub>


Choose the most appropriate answer from the options given below :

- (1) (A), (C) only
- (2) (A) only
- (3) (B) only
- (4) (A), (B) only


**Official Ans. by NTA (4)**

Sol. In basic medium, oxidising action of H<sub>2</sub>O<sub>2</sub>.  
Mn<sup>2+</sup> + H<sub>2</sub>O<sub>2</sub> → Mn<sup>4+</sup> + 2OH<sup>-</sup>

In basic medium, reducing action of H<sub>2</sub>O<sub>2</sub>




In acidic medium, oxidising action of H<sub>2</sub>O<sub>2</sub>.




Hence correct option (4)

3. In the reaction of hypobromite with amide, the carbonyl carbon is lost as :

- (1) CO<sub>3</sub><sup>2-</sup>
- (2) HCO<sub>3</sub><sup>-</sup>
- (3) CO<sub>2</sub>
- (4) CO



**Mechanism**







4. The oxide that shows magnetic property is :


- (1) SiO<sub>2</sub>
- (2) Mn<sub>3</sub>O<sub>4</sub>
- (3) Na<sub>2</sub>O
- (4) MgO

**Official Ans. by NTA (2)**


Sol. Mn<sub>3</sub>O<sub>4</sub> shows magnetic properties.

5. Main Products formed during a reaction of 1-methoxy naphthalene with hydroiodic acid are:





**Mechanism**



6. Deficiency of vitamin K causes :
- (1) Increase in blood clotting time
  - (2) Increase in fragility of RBC's
  - (3) Cheilosis
  - (4) Decrease in blood clotting time

**Official Ans. by NTA (1)**


Sol. Due to deficiency of Vitamin K causes increases in blood clotting time.

**Note :** Vitamin K related to blood factor.

7. An organic compound "A" on treatment with benzene sulphonyl chloride gives compound B. B is soluble in dil. NaOH solution. Compound A is :
- (1)  $C_6H_5-N-(CH_3)_2$
  - (2)  $C_6H_5-NHCH_2CH_3$
  - (3)  $C_6H_5-CH_2NHCH_3$
  - (4)  $C_6H_5-\underset{\substack{| \\ CH_3}}{CH}-NH_2$

**Official Ans. by NTA (4)**

Sol. Hinsberg reagent (Benzene sulphonyl chloride) gives reaction product with  $1^\circ$  amine and it is soluble in dil. NaOH.



8. The first ionization energy of magnesium is smaller as compared to that of elements X and Y, but higher than that of Z. The elements X, Y and Z, respectively, are :

- (1) chlorine, lithium and sodium
- (2) argon, lithium and sodium
- (3) argon, chlorine and sodium
- (4) neon, sodium and chlorine


**Official Ans. by NTA (3)**

Sol. The 1<sup>st</sup> IE order of 3<sup>rd</sup> period is  $Na < Al < Mg < Si < S < P < Cl < Ar$   
X & Y are Ar & Cl  
Z is sodium (Na).

9. The secondary valency and the number of hydrogen bonded water molecule(s) in  $CuSO_4 \cdot 5H_2O$ , respectively, are :

- (1) 6 and 4
- (2) 4 and 1
- (3) 6 and 5
- (4) 5 and 1

**Official Ans. by NTA (2)**



10. Given below are two statements :

Statement I : Bohr's theory accounts for the stability and line spectrum of  $Li^+$  ion.


Statement II : Bohr's theory was unable to explain the splitting of spectral lines in the presence of a magnetic field.

In the light of the above statements, choose the most appropriate answer from the options given below :

- (1) Both statement I and statement II are true.
- (2) Statement I is false but statement II is true.
- (3) Both statement I and statement II are false.
- (4) Statement I is true but statement II is false.

**Sol.** Statement-I is false since Bohr's theory accounts for the stability and spectrum of single electronic species (eg :  $\text{He}^+$ ,  $\text{Li}^{2+}$  etc)


Statement II is true.



Consider the given reaction, percentage yield of:

- (1)  $C > A > B$                       (2)  $B > C > A$   
 (3)  $A > C > B$                       (4)  $C > B > A$

**Official Ans. by NTA (4)**



% yield order  $\Rightarrow C > B > A$

12. The charges on the colloidal  $\text{CdS}$  sol and  $\text{TiO}_2$  sol are, respectively :

- (1) positive and positive  
 (2) positive and negative  
 (3) negative and negative  
 (4) negative and positive

**Official Ans. by NTA (4)**

**Sol.**  $\text{CdS}$  sol  $\rightarrow$  -ve sol

$\text{TiO}_2$  sol  $\rightarrow$  +ve sol

13. Match List - I with List - II :

List - I


List - II

- |  |                     |
|--|---------------------|
| (Class of Chemicals)                       | (Example)           |
| (a) Antifertility drug                     | (i) Meprobamate     |
| (b) Antibiotic                             | (ii) Alitame        |
| (c) Tranquilizer                           | (iii) Norethindrone |
| (d) Artificial Sweetener                   | (iv) Salvarsan      |
| (1) (a)-(ii), (b)-(iii), (c)-(iv), (d)-(i) |                     |
| (2) (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i) |                     |
| (3) (a)-(iii), (b)-(iv), (c)-(i), (d)-(ii) |                     |





**Official Ans. by NTA (3)**

- Sol.** (A) Antifertility drug  $\rightarrow$  (iii) Nor ethindrone  
 (B) Antibiotic  $\rightarrow$  (iv) Salvarsan  
 (C) Tranquilizer  $\rightarrow$  (i) Meprobamate  
 (D) Artificial sweetener  $\rightarrow$  (ii) Alitame


**Ans.** A-iii, B-iv, C-i, D-ii



Consider the above reaction, the product 'X' and 'Y' respectively are :

- (1) 
- (2) 
- (3) 
- (4) 

**Official Ans. by NTA (3)**



15. Match list-I with list-II :

List-I	List-II
(a) Be	(i) Treatment of cancer
(b) Mg	(ii) Extraction of metals
(c) Ca	(iii) Incendiary bombs and signals
(d) Ra	(iv) Windows of X-ray tubes
	(v) Bearings for motor engines.

Choose the most appropriate answer the option given below :

- (1) a-iv, b-iii, c-i, d-ii
- (2) a-iv, b-iii, c-ii, d-i
- (3) a-iii, b-iv, c-v, d-ii
- (4) a-iii, b-iv, c-ii, d-v

**Official Ans. by NTA (2)**

- Sol.** (a) Be → it is used in the Windows of X-ray tubes  
 (b) Mg → it is used in the Incendiary bombs and signals  
 (c) Ca → it is used in the Extraction of metals  
 (d) Ra → it is used in the Treatment of cancer

16. Given below are two statements :

**Statement I :**  $C_2H_5OH$  and  $AgCN$  both can generate nucleophile.

**Statement II :**  $KCN$  and  $AgCN$  both will generate nitrile nucleophile with all reaction conditions.

Choose the most appropriate option :

- (1) Statement I is true but statement II is false
- (2) Both statement I and statement II are true
- (3) Statement I is false but statement II is true
- (4) Both statement I and statement II are false

**Official Ans. by NTA (1)**

17. Given below are two statements :

**Statement I :** Non-biodegradable wastes are generated by the thermal power plants.

**Statement II :** Bio-degradable detergents leads to eutrophication.

In the light of the above statements, choose the most appropriate answer from the option given

- (1) Both statement I and statement II are false
- (2) Statement I is true but statement II is false
- (3) Statement I is false but statement II is true
- (4) Both statement I and statement II are true.

**Official Ans. by NTA (4)**

- Sol.** Non-biodegradable wastes are generated by the thermal power plants which produces fly ash. Detergents which are biodegradable causes problem called eutrophication which kills animal life by depriving it of oxygen.

18. Match list-I with list-II :

List-I	List-II
(a) Mercury	(i) Vapour phase refining
(b) Copper	(ii) Distillation refining
(c) Silicon	(iii) Electrolytic refining
(d) Nickel	(iv) Zone refining


Choose the most appropriate answer from the option given below :

- (1) a-i, b-iv, c-ii, d-iii
- (2) a-ii, b-iii, c-i, d-iv
- (3) a-ii, b-iii, c-iv, d-i
- (4) a-ii, b-iv, c-iii, d-i

**Official Ans. by NTA (3)**

- Sol.** (a) Mercury → Distillation refining  
 (b) Copper → Electrolytic refining  
 (c) Silicon → Zone refining  
 (d) Nickel → Vapour phase refining


19. In the following molecules,



Hybridisation of carbon a, b and c respectively are :

- (1)  $sp^3$ ,  $sp$ ,  $sp$
- (2)  $sp^3$ ,  $sp^2$ ,  $sp$
- (3)  $sp^3$ ,  $sp^2$ ,  $sp^2$
- (4)  $sp^3$ ,  $sp$ ,  $sp^2$

**Official Ans. by NTA (3)**



20. A hard substance melts at high temperature and is an insulator in both solid and in molten state.

This solid is most likely to be a / an :

- (1) Ionic solid
- (2) Molecular solid
- (3) Metallic solid
- (4) Covalent solid

**Official Ans. by NTA (4)**

**Sol.** Covalent or network solid have very high melting point and they are insulators in their solid and molten form.

### SECTION-B

1. A reaction has a half life of 1 min. The time required for 99.9% completion of the reaction is \_\_\_\_\_ min. (Round off to the Nearest integer)

[Use :  $\ln 2 = 0.69$ ,  $\ln 10 = 2.3$ ]

**Official Ans. by NTA (10)**

**Sol.**

$$\frac{t_{99.9\%}}{t_{50\%}} = \frac{\frac{1}{K} \ln \frac{100}{0.1}}{\frac{1}{K} \ln 2}$$

$$= \frac{\ln 1000}{\ln 2} \times t_{50\%}$$

$$= \frac{3 \ln 10}{\ln 2} \times 1$$

$$= \frac{3 \times 2.3}{0.69} = 10$$

2. The molar conductivities at infinite dilution of barium chloride, sulphuric acid and hydrochloric acid are 280, 860 and 426  $\text{Scm}^2 \text{mol}^{-1}$  respectively. The molar conductivity at infinite dilution of barium sulphate is \_\_\_\_\_  $\text{S cm}^2 \text{mol}^{-1}$  (Round off to the Nearest Integer).

**Official Ans. by NTA (288)**

**Sol.** From Kohlrausch's law

$$\Lambda_m^\infty(\text{BaSO}_4) = \lambda_m^\infty(\text{Ba}^{2+}) + \lambda_m^\infty(\text{SO}_4^{2-})$$

$$\Lambda_m^\infty(\text{BaSO}_4) = \Lambda_m^\infty(\text{BaCl}_2) + \Lambda_m^\infty(\text{H}_2\text{SO}_4) - 2 \Lambda_m^\infty(\text{HCl})$$





$$= 280 + 860 - 2(426)$$

$$= 288 \text{ Scm}^2 \text{mol}^{-1}$$

3. The number of species below that have two lone pairs of electrons in their central atom is \_\_\_\_\_ (Round off to the Nearest integer)

$\text{SF}_4$ ,  $\text{BF}_4^-$ ,  $\text{ClF}_3$ ,  $\text{AsF}_3$ ,  $\text{PCl}_5$ ,  $\text{BrF}_5$ ,  $\text{XeF}_4$ ,  $\text{SF}_6$

**Official Ans. by NTA (2)**



Two l.p. on central atom is =  $\text{ClF}_3$ ,  $\text{XeF}_4$


4. A xenon compound 'A' upon partial hydrolysis gives  $\text{XeO}_2\text{F}_2$ . The number of lone pair of electrons present in compound A is \_\_\_\_\_ (Round off to the Nearest integer)

**Official Ans. by NTA (19)**

**Sol.**  $\text{XeF}_6 + 2\text{H}_2\text{O} \longrightarrow \text{XeO}_2\text{F}_2 + 4\text{HF}$


(A) (Limited water)

Structure of 'A'



Total l.p. on (A) = 19

5. The gas phase reaction



at 400 K has  $\Delta G^\circ = + 25.2 \text{ kJ mol}^{-1}$ .

The equilibrium constant  $K_C$  for this reaction is \_\_\_\_\_  $\times 10^{-2}$ . (Round off to the Nearest integer)

[Use :  $R = 8.3 \text{ J mol}^{-1}\text{K}^{-1}$ ,  $\ln 10 = 2.3$

$\log_{10} 2 = 0.30$ ,  $1 \text{ atm} = 1 \text{ bar}$

[antilog (-0.3) = 0.501]

**Official Ans. by NTA (166)**

**Official Ans. by** (2)

**Sol.** Using formula

$$\Delta_r G^0 = -RT \ln K_p$$

$$25200 = -2.3 \times 8.3 \times 400 \log(K_p)$$

$$K_p = 10^{-3.3} = 10^{-3} \times 0.501$$

$$= 5.01 \times 10^{-4} \text{ Bar}^{-1}$$

$$= 5.01 \times 10^{-9} \text{ Pa}^{-1}$$

$$= \frac{K_c}{8.3 \times 400}$$




$$K_c = 1.66 \times 10^{-5} \text{ m}^3/\text{mole}$$

$$= 1.66 \times 10^{-2} \text{ L/mol}$$

$$\text{Ans} = 2$$

6. In Tollen's test for aldehyde, the overall number of electron(s) transferred to the Tollen's reagent formula  $[\text{Ag}(\text{NH}_3)_2]^+$  per aldehyde group to form silver mirror is \_\_\_\_\_. (Round off to the Nearest integer)

**Official Ans. by NTA (2)**



Total  $2e^-$  transfer to Tollen's reagent

7. The solubility of  $\text{CdSO}_4$  in water is  $8.0 \times 10^{-4} \text{ mol L}^{-1}$ . Its solubility in  $0.01 \text{ M H}_2\text{SO}_4$  solution is \_\_\_\_\_  $\times 10^{-6} \text{ mol L}^{-1}$ . (Round off to the Nearest integer) (Assume that solubility is much less than  $0.01 \text{ M}$ )



**Official Ans. by NTA (64)**

**Sol.** In pure water,

$$K_{sp} = S^2 = (8 \times 10^{-4})^2$$

$$= 64 \times 10^{-8}$$

In  $0.01 \text{ M H}_2\text{SO}_4$



$$x \qquad \qquad \qquad x \qquad \qquad (x + 0.01)$$

$$K_{sp} = x(x + 0.01)$$

$$= 64 \times 10^{-8}$$

$$x + 0.01 \cong 0.01 \text{ M}$$

$$\text{So, } x(0.01) = 64 \times 10^{-8}$$

$$x = 64 \times 10^{-6} \text{ M}$$

8. A solute dimerizes in water. The boiling point of a 2 molar solution of A is  $100.52^\circ\text{C}$ . The percentage association of A is \_\_\_\_\_. (Round off to the Nearest integer)

(Round off to the Nearest integer)

[Use :  $K_b$  for water =  $0.52 \text{ K kg mol}^{-1}$

Boiling point of water =  $100^\circ\text{C}$ ]

**Official Ans. by NTA (100)**

**Sol.**  $\Delta T_b = T_b - T_b^0$

$$100.52 - 100$$

$$= 0.52^\circ\text{C}$$

$$i = \left(1 - \frac{\alpha}{2}\right)$$

$$\therefore \Delta T_b = i K_b \times m$$

$$0.52 = \left(1 - \frac{\alpha}{2}\right) \times 0.52 \times 2$$

$$\alpha = 1$$

So, percentage association = 100%.


9.  $10.0 \text{ ml}$  of  $\text{Na}_2\text{CO}_3$  solution is titrated against  $0.2 \text{ M HCl}$  solution. The following titre values were obtained in 5 readings.

$4.8 \text{ ml}$ ,  $4.9 \text{ ml}$ ,  $5.0 \text{ ml}$ ,  $5.0 \text{ ml}$  and  $5.0 \text{ ml}$

Based on these readings, and convention of titrimetric estimation of concentration of  $\text{Na}_2\text{CO}_3$  solution is \_\_\_\_\_  $\text{mM}$ .

(Round off to the Nearest integer)

**Sol.** Most precise volume of HCl = 5 ml  
at equivalence point  
Meq. of  $\text{Na}_2\text{CO}_3$  = meq. of HCl.  
Let molarity of  $\text{Na}_2\text{CO}_3$   
solution = M, then  
 $M \times 10 \times 2 = 0.2 \times 5 \times 1$   
 $M = 0.05 \text{ mol / L}$   
 $= 0.05 \times 1000$   
 $= 50 \text{ mM}$



Consider the above reaction where 6.1 g of benzoic acid is used to get 7.8 g of m-bromo benzoic acid. The percentage yield of the product is\_\_\_\_\_.

(Round off to the Nearest integer)

[Given : Atomic masses : C = 12.0u, H : 1.0u, O : 16.0u, Br = 80.0 u]

**Official Ans. by NTA (78)**

**Sol.** Moles of Benzoic acid =  $\frac{6.1}{122}$   
= moles of m-bromobenzoic acid  
So, weight of m-bromobenzoic acid  
 $= \frac{6.1}{122} \times 201 \text{ gm}$   
 $= 10.05 \text{ gm}$   
 $\% \text{ yield} = \frac{\text{Actual weight}}{\text{Theoretical weight}} \times 100$   
 $= \frac{7.8}{10.05} \times 100$   
 $= 77.61\%$