

2016

## Section-I Mathematics

1. Use the following figure to find  $x^\circ$  and  $y^\circ$



- (A)  $x = 50^\circ$ ,  $y = 30^\circ$  (B)  $x = 30^\circ$ ,  $y = 50^\circ$   
(C)  $x = 50^\circ$ ,  $y = 60^\circ$  (D)  $x = 55^\circ$ ,  $y = 65^\circ$

2.  $\frac{2}{x} + \frac{5}{y} = 1$  and  $\frac{60}{x} - \frac{20}{y} = 13$

For these equations, the value of  $x$  and  $y$  will be—

- (A)  $x = 4$ ,  $y = 10$   
(B)  $x = 10$ ,  $y = 4$   
(C)  $x = \frac{1}{4}$ ,  $y = 5$   
(D) None of these

3. In an election of Sarpanch, there were two candidates. A total of 9701 votes were polled. In which 116 votes were declared invalid. The successful candidate got 5 votes in the place of 4 votes of his opponent got. The margin of won of successful candidate is—

- (A) 2008  
(B) 775  
(C) 1075  
(D) None of these


4. The value of  $\log_2 (\log_2 (\log_2 16))$  is—

- (A)  $\frac{1}{2}$   
(B)  $\frac{1}{3}$   
(C)  $\frac{1}{4}$   
(D) None of these

5. In the given figure, MN is parallel to PR,  $\angle LBN = 70^\circ$  and  $AB = BC$ . The value of  $\angle ABC$  will be—

- (A)  $\frac{1}{2}BC$   
(B)  $\frac{1}{3}BC$   
(C)  $\frac{2}{3}BC$   
(D) None of these

8. If the line PQ is parallel to line BC of triangle ABC, then



15. The value of  $\sqrt{(\cosec A - \sin A)(\sec A - \cos A)(\tan A)}$  is—

- (A)  $\sqrt{3}$   
(B) 2  
(C)  $\sqrt{3}$   
(D)  $\frac{1}{2}$

16. A rocket is fired vertically upward from launching pad P. It first rises 40 km vertically upwards and then 40 km at  $60^\circ$  to the vertical. The height attained by the rocket from launching pad P is—

- (A) 80 km  
(B) 60 km  
(C) 65 km  
(D) 85 km

17. If the difference between the two sides of a right-angled triangle is 2 cm and the area of the triangle is 24 cm<sup>2</sup>. The perimeter of the triangle will be—

- (A) 20 cm  
(B) 24 cm  
(C) 30 cm  
(D) 15 cm

18. The line  $4x - 3y + 12 = 0$  meets x-axis at A. The co-ordinates of A are—

- (A) (4, 0)  
(B) (4, 3)  
(C) (3, 0)  
(D) (3, 12)

19. The value of  $7^{\circ} \times (25)^{\circ} 32' - 5^{\circ} 3' 0''$  is—

- (A) 0  
(B) 1  
(C) 2  
(D) 1

20. Two straight lines  $3x - 2y = 5$  and  $2x + ky + 7 = 0$  are perpendicular to each other. The value of  $k$  is—

- (A) 3  
(B)  $\frac{1}{3}$   
(C)  $-\frac{4}{3}$   
(D)  $-\frac{3}{2}$

21. The co-ordinates of A and C are (3, 6) and (-1, 2) in a rhombus ABCD. The equation of BD is—

- (A)  $2x + 4y = 5$   
(B)  $2x + 4y = 0$   
(C)  $x + 4y = 12$   
(D)  $x + y = 5$

22.  $\int_{-6}^{-6} \frac{dx}{x-3} = \frac{2-6}{2-3} (\infty - 3)$

- (A)  $616 \text{ cm}^3$   
(B)  $416 \text{ cm}^3$   
(C)  $308 \text{ cm}^3$   
(D)  $832 \text{ cm}^3$

$$x + y = -3$$

22. In the given diagram of semicircle, O is centre.



Weight in kg	57	58	59	60	61
No. of boys	10	13	8	12	6

The median is

- (A) 50 (B) 59  
(C) 60 (D) 78

The area of the shaded portion is—

- (A)  $6\pi \text{ cm}^2$  (B)  $10\pi \text{ cm}^2$   
(C)  $63\pi \text{ cm}^2$  (D)  $61\pi \text{ cm}^2$

23. In the given figure, PS is a diameter of a circle and is of length 10 cm. Q and R are points on the diameter such that PQ, QR and RS are equal. Segments are drawn with PQ and QS as diameter.



- (A) ₹ 15,768 (B) ₹ 14,000  
(C) ₹ 15,600 (D) ₹ 15,500

(A) 15 and 24	(B) 21 and 34
(C) 6 and 18 (D) 24 and 40	

The numbers are—

- (A) 14 (B)  $\frac{1}{4}$   
(C) 24 (D) None of these

The locus of point P is—

- (A) Circle with diameter AB  
(B) Straight line  
(C) Triangle  
(D) Right angled triangle

30. If price of a TV set inclusive of sales tax of 9% is ₹ 40,221, then the marked price is—

- (A) ₹ 41,840.89 (B) ₹ 36,198.90  
(C) ₹ 36,900 (D) ₹ 43,890

31. Two numbers are in the ratio 3 : 5. If 8 is added to each number, the ratio becomes 2 : 3. The numbers are—

- (A) 15 and 24 (B) 21 and 34  
(C) 6 and 18 (D) 24 and 40

where  $x \neq 0$  and  $y \neq 1$ , are

32. Numbers 4, 5, 16 and 26 will become proportional if we add number in each of them—

- (A) 4 (B) 2  
(C) 6 (D) None of these

33. The roots of equation  $a\alpha^2 + b\alpha + c = 0$  where  $a, b, c \in \mathbb{R}$  and  $a \neq 0$  is equal in magnitude and opposite in sign if—

- (A)  $b^2 - 4ac \geq 0$  (B)  $b^2 - 4ac = 0$   
(C)  $b = 0$  (D)  $a = 0$

34. The value of  $\cot^2 \theta (1 - \cos^2 \theta)$  is

- (A) 1 (B) 4  
(C) tan  $\theta$  (D)  $\cos^2 \theta$

35. If  $X = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ ,  $Y = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,

- then the value of  $XY^T$  is—

- (A)  $\begin{bmatrix} -2 \\ 4 \end{bmatrix}$  (B)  $\begin{bmatrix} 9 \\ 0 \end{bmatrix}$   
(C)  $\begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$  (D)  $\begin{bmatrix} 1 & 0 \\ 4 & 1 \end{bmatrix}$

36. A man borrows ₹ 5,100 at 12% compound interest per annum, interest payable every 6 months. He pays back ₹ 1,800 at the end of every six months. The total payment he has to make at the end of 18 months in order to clear the entire loan is—

- (A) ₹ 5,624.00 (B) ₹ 3,024.60  
(C) ₹ 3,824.60 (D) None of these

then the value of 'x' is—

- (A) 31 (B) 49  
(C) 12 (D) 18

42. Rohit has ₹ 9,500 more than Deep. If gives ₹ 2,000 to Rohit, the money with will be four times the money left with him initially with Rohit and me—

- (A) ₹ 13,500 and ₹ 4,000  
(B) ₹ 16,500 and ₹ 6,000  
(C) ₹ 17,500 and ₹ 6,500  
(D) None of the above

The money initially with Rohit and

- (A) Factor of expression  $2x^3 + 5x^2 - 14x - 1$   
(B)  $(2x + 7)(x + 1)(x - 2)$   
(C)  $(x + 7)(2x + 1)(x - 2)$   
(D) None of the above

The value of 'x' for equation

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

- (A)  $\frac{4}{9}$  (B)  $\frac{1}{3}$  (C)  $\frac{3}{4}$  (D)  $\frac{4}{3}$

48. The arithmetic mean of given data will be—  
 (A) 62 (B) 72  
 (C) 71 (D) 67, 65, 71, 57 and 45
49. In the given chart, the mode of the following frequency distribution is—  
 Number      8      9      10      11      12      13  
 Frequency      3      1      8      13      9      5
- (A) 8 (B) 10  
 (C) 13 (D) 11
50. In the given figure, O is the centre of the circle and  $\angle AOD = 180^\circ$ ,  $\angle DAO = 51^\circ$ . The values of angles x and y will be—  
 D  
  
 (A) x = 31.5°, y = 24° (B) x = 24°, y = 48°  
 (C) x = 21°, y = 48° (D) x = 50°, y = 24°

## Physics and Chemistry

51. The percentae of nitrogen in ammonium nitrate ( $\text{NH}_4\text{NO}_3$ ) is, where H = 1, N = 14,  
 $10 \div 16 \times 100$   
 (A) 35% (B) 28%  
 (C) 48% (D) 15%
52. The volume of 7.1 g of chlorine at STP is—  
 (A) 22.4 litres (B) 2.24 litres  
 (C) 4.48 litres (D) 1.12 litres
53. The compound which decomposes on passing electricity through its aqueous solution is—  
 (A) Sugar (B) Urea  
 (C) Copper sulphate (D) Ethyl alcohol
54. The flow of current in an electrolyte is due to the movement of—  
 (A) Electrons (B) Molecules  
 (C) Protons (D) Ions
55. The carbon content in stainless steel is—  
 (A) 0.1 – 0.25% (B) 0.5 – 1.0%  
 (C) 1 – 2.5% (D) More than 2.5%
56. Which of the following does not contain aluminum?  
 (A) Alumina (B) Dicalcium  
 (C) gain of electrons (D) gain or loss of electrons
57. The catalyst used for making ammonia ( $\text{NH}_3$ ) gas from nitrogen and hydrogen is—  
 (A) iron and molybdenum  
 (B) iron and platinum  
 (C) molybdenum and platinum
58. Sodium potassium sulphate ( $\text{NaKSO}_4$ ) is an—  
 (A) complex salt (B) mixed salt  
 (C) impure salt (D) None of these
59. The acid used in storage batteries is—  
 (A) sulphuric acid (B) nitric acid  
 (C) phosphoric acid (D) None of these
60. Catalyst  $\text{V}_2\text{O}_5$  is used in manufacturing of—  
 (A)  $\text{SO}_2$  from S  
 (B)  $\text{SO}_3$  from  $\text{SO}_2$   
 (C)  $\text{H}_2\text{SO}_4$  from  $\text{SO}_3$   
 (D) None of the above
61. Which of the following is unsaturated compound?—  
 (A)  $\text{C}_2\text{H}_4$  (B)  $\text{C}_2\text{H}_6$   
 (C)  $\text{C}_2\text{H}_5\text{Cl}$  (D)  $\text{CH}_3\text{OH}$
62. IUPAC name of the  $\text{C}_2\text{H}_9\text{Cl}^+$  is—  
 (A) Alkane (B) Alkyne  
 (C) Alkene (D) None of these
63. Halogenation is—  
 (A) addition reaction  
 (B) oxidation reaction  
 (C) reduction reaction  
 (D) None of the above
64. In an aqueous solution of a compound A, a reddish brown precipitate is obtained on adding  $\text{Na}_3\text{AlO}_2$  drop by drop. The compound A is—  
 (A) Zinc sulphate (B) Lead nitrate  
 (C) Ferric sulphate (D) Ferric chloride
65. 1 mole of compound contains 1 mole of carbon and 2 moles of oxygen. The molecular weight of the compound is—  
 (A) 3 (B) 12  
 (C) 32 (D) 44
66. Which of the following has water of crystallization?—  
 (A) Zinc chloride (B) Caustic soda  
 (C) Baking soda (D) Washing soda
67. Sodium potassium sulphate ( $\text{NaKSO}_4$ ) is—  
 (A) A (B) B  
 (C) C (D) D
68. Acids A, B, C and D have pH values 1, 2, 3, 4 respectively. Which one is stronger acid?—  
 (A) A (B) B  
 (C) C (D) D
69. Methyl orange indicator provides yellow colour in—  
 (A) acidic solution (B) alkaline solution  
 (C) neutral solution (D) None of these
70. In the preparation of  $\text{FeCl}_3$ , we use fused calcium chloride—  
 (A) as a catalyst  
 (C) to keep  $\text{FeCl}_3$  dry  
 (D) None of the above
71. In an aqueous solution of a compound A, a reddish brown precipitate is obtained on adding  $\text{Na}_3\text{AlO}_2$  drop by drop. The compound A is—  
 (A) Zinc sulphate (B) Lead nitrate  
 (C) Ferric sulphate (D) Ferric chloride
72. 1 mole of compound contains 1 mole of carbon and 2 moles of oxygen. The molecular weight of the compound is—  
 (A) 3 (B) 12  
 (C) 32 (D) 44
73. Mass of oxygen in 36 g of pure water is—  
 (A) 16 g (B) 64 g  
 (C) 70 g (D) 32 g
74. Power of a pump which lifts 100 kg of water to a water tank situated at 20 m height in 10 seconds is (Take  $g = 10 \text{ m/s}^2$ )—  
 (A)  $0.1 \text{ A to } 100 \text{ A}$   
 (B)  $100 \text{ A to } 4000 \text{ A}$
75. The lever for which the mechanical advantage is always less than one has—  
 (A) fulcrum between load and effort  
 (B) load between effort and fulcrum  
 (C) effort between fulcrum and load  
 (D) None of the above
76. Mechanical advantage (MA), load (L), effort (F) are related as—  
 (A)  $MA = L \times F$  (B)  $L = MA \times F$   
 (C)  $E = MA \times L$  (D) None of these
77. A ray of light is incident normally on surface of water. Its angle of refraction water is—  
 (A)  $90^\circ$  (B)  $180^\circ$   
 (C)  $0^\circ$  (D)  $45^\circ$
78. The apparent depth of a pond is 9 m. Whiile the real depth if refractive index of water is respect to air is—  
 (A) 12 m (B) 6 m  
 (C) 6.25 m (D) 9 m
79. The point through which a ray of light passes without suffering deviation is called—  
 (A) Pole (B) Focus  
 (C) Centre of curvature (D) Optical centre
80. Lens used for projecting the image on screen is—  
 (A) convex lens (B) concave lens  
 (C) convex and concave lens (D) None of the above
81. A convex lens of focal length 20 cm is placed in contact of with a concave lens of focal length 40 cm. The nature of resulted lens is—  
 (A) convex lens (B) concave lens  
 (C) plane-convex lens (D) None of the above

84. Between  $\lambda = 4000\text{ \AA}$  to  $8000\text{ \AA}$   
 (A)  $4000\text{ \AA}$  to  $8000\text{ \AA}$   
 (B)  $4000\text{ \AA}$  to  $800\text{ \AA}$   
 (C)  $4000\text{ \AA}$  to  $8000\text{ \AA}$   
 (D) Above  $8000\text{ \AA}$
85. Source of ultraviolet light is—  
 (A) electric bulb  
 (B) ~~red hot iron ball~~  
 (C) sodium vapor lamp  
 (D) carbon arc lamp
86. The frequency of electromagnetic wave which has wavelength  $30\text{ cm}$  and velocity  $3 \times 10^8\text{ m/s}$  is—  
 (A)  $3000\text{ MHz}$   
 (B)  $300\text{ MHz}$   
 (C)  $30\text{ MHz}$   
 (D) None of these
87. If the amplitude of a wave is doubled, then its loudness becomes—  
 (A) double  
 (B) three times  
 (C) four times  
 (D) unchanged
88. Intensity of sound is measured in—  
 (A) hertz (Hz)  
 (B) decibel (dB)  
 (C) second  
 (D) metre
89. The resistance between points X and Y in the given figure is—  
 X,  $v_1$ ,  $v_2$ , P, Q, R, S, Y  
 (A)  $v_1 + v_2$   
 (B)  $v_1 - v_2$   
 (C)  $v_1 + v_2 + v_3$   
 (D)  $v_1 - v_2 - v_3$
90. The resistance between points X and Y in the given figure is—  
 X,  $v_1$ ,  $v_2$ , P, Q, R, S, Y  
 (A)  $v_1 + v_2$   
 (B)  $v_1 - v_2$   
 (C)  $v_1 + v_2 + v_3$   
 (D)  $v_1 - v_2 - v_3$
91. Which of the following is not a greenhouse gas?  
 (A)  $\text{CO}_2$   
 (B)  $\text{CH}_4$   
 (C)  $\text{N}_2\text{O}$   
 (D)  $\text{CH}_4$
92. A liquid A of mass  $10\text{ g}$  and at  $120^\circ\text{C}$  is poured in a liquid B at  $20^\circ\text{C}$ . The final temperature recorded is  $40^\circ\text{C}$ . The initial mass of liquid B is (If initial capacity of liquid A is  $0.8\text{ J/g}^\circ\text{C}$  and of liquid B  $4.2\text{ J/g}^\circ\text{C}$ )—  
 (A)  $70\text{ g}$   
 (B)  $80.0\text{ g}$   
 (C)  $70\text{ g}$   
 (D)  $76.1\text{ g}$
93. Among the elements Li, Na, Mg and Al, the lowest value of ionization potential of certain will be—  
 (A) Li  
 (B) Na  
 (C) Mg  
 (D) Al
94. On moving from left to right in a period of periodic table, the number of shell—  
 (A) remains the same  
 (B) increases  
 (C) decreases  
 (D) first increases then decreases
95. 1.0 kilogram force (kgf) is equal to—  
 (A)  $8\text{ N}$   
 (B)  $2\text{ N}$   
 (C)  $15\text{ N}$   
 (D) None of these
96. Four cells each of emf  $2\text{ V}$  and internal resistance  $0.1\text{ }\Omega$  are connected in series. The combination is connected in series to an ammeter of negligible resistance, a  $1.6\text{ }\Omega$  resistor and an unknown resistor R. The current in the circuit is  $2\text{ A}$ . The value of R is—  
 (A)  $4\text{ }\Omega$   
 (B)  $2\text{ }\Omega$   
 (C)  $6\text{ }\Omega$   
 (D)  $8\text{ }\Omega$
97. Wheels of a moving cycle ~~have~~ have rotational and translational motion  
 (A) have translational motion only  
 (B) have rotational motion only  
 (C) have both motion only  
 (D) None of the above
98. ~~Margin~~ margin of unit of—  
 (A)  $\text{J}/\text{kg}$   
 (B) force  
 (C) power  
 (D) energy

90. Specific heat capacity of copper is  $0.1\text{ cal/gm}$ . Its value in  $\text{J/kg}^\circ\text{C}$  is—(A)  $0.84 \times 10^3$   
 (B)  $0.42 \times 10^4$   
 (C)  $0.24 \times 10^3$   
 (D)  $4.2 \times 10^3$ 91. Which of the following is not a greenhouse gas?  
 (A)  $\text{CO}_2$   
 (B)  $\text{CH}_4$   
 (C)  $\text{N}_2\text{O}$   
 (D)  $\text{CH}_4$ 92. A liquid A of mass  $10\text{ g}$  and at  $120^\circ\text{C}$  is poured in a liquid B at  $20^\circ\text{C}$ . The final temperature recorded is  $40^\circ\text{C}$ . The initial mass of liquid B is (If initial capacity of liquid A is  $0.8\text{ J/g}^\circ\text{C}$  and of liquid B  $4.2\text{ J/g}^\circ\text{C}$ )—  
 (A)  $70\text{ g}$   
 (B)  $80.0\text{ g}$   
 (C)  $70\text{ g}$   
 (D)  $76.1\text{ g}$ 93. Among the elements Li, Na, Mg and Al, the lowest value of ionization potential of certain will be—  
 (A) Li  
 (B) Na  
 (C) Mg  
 (D) Al94. On moving from left to right in a period of periodic table, the number of shell—  
 (A) remains the same  
 (B) increases  
 (C) decreases  
 (D) first increases then decreases95. 1.0 kilogram force (kgf) is equal to—  
 (A)  $8\text{ N}$   
 (B)  $2\text{ N}$   
 (C)  $15\text{ N}$   
 (D) None of these96. The force of friction on a body kept on the table top does not depend on—  
 (A) nature of force  
 (B) material of the body  
 (C) weight of the body  
 (D) area of contact97. Wheels of a moving cycle ~~have~~ have rotational and translational motion  
 (A) have translational motion only  
 (B) have rotational motion only  
 (C) have both motion only  
 (D) None of the above98. ~~Margin~~ margin of unit of—  
 (A)  $\text{J}/\text{kg}$   
 (B) force  
 (C) power  
 (D) energy90. Specific heat capacity of copper is  $0.1\text{ cal/gm}$ . Its value in  $\text{J/kg}^\circ\text{C}$  is—(A)  $0.84 \times 10^3$   
 (B)  $0.42 \times 10^4$   
 (C)  $0.24 \times 10^3$   
 (D)  $4.2 \times 10^3$ 91. Which one is not a fossil fuel?  
 (A) Coal  
 (B) Biomass  
 (C) Petroleum  
 (D) None of these92. If a force F is applied on a body and it moves with a velocity v, then the value of the power will be—  
 (A)  $F/v$   
 (B)  $F \times v$   
 (C)  $Fv^2$   
 (D)  $F \times v^2$ 93. A liquid A of mass  $10\text{ g}$  and at  $120^\circ\text{C}$  is poured in a liquid B at  $20^\circ\text{C}$ . The final temperature recorded is  $40^\circ\text{C}$ . The initial mass of liquid B is (If initial capacity of liquid A is  $0.8\text{ J/g}^\circ\text{C}$  and of liquid B  $4.2\text{ J/g}^\circ\text{C}$ )—  
 (A)  $70\text{ g}$   
 (B)  $80.0\text{ g}$   
 (C)  $70\text{ g}$   
 (D)  $76.1\text{ g}$ 94. Among the elements Li, Na, Mg and Al, the lowest value of ionization potential of certain will be—  
 (A) Li  
 (B) Na  
 (C) Mg  
 (D) Al95. On moving from left to right in a period of periodic table, the number of shell—  
 (A) remains the same  
 (B) increases  
 (C) decreases  
 (D) first increases then decreases96. The force of friction on a body kept on the table top does not depend on—  
 (A) nature of force  
 (B) material of the body  
 (C) weight of the body  
 (D) area of contact97. Wheels of a moving cycle ~~have~~ have rotational and translational motion  
 (A) have translational motion only  
 (B) have rotational motion only  
 (C) have both motion only  
 (D) None of the above98. ~~Margin~~ margin of unit of—  
 (A)  $\text{J}/\text{kg}$   
 (B) force  
 (C) power  
 (D) energy90. Specific heat capacity of copper is  $0.1\text{ cal/gm}$ . Its value in  $\text{J/kg}^\circ\text{C}$  is—(A)  $0.84 \times 10^3$   
 (B)  $0.42 \times 10^4$   
 (C)  $0.24 \times 10^3$   
 (D)  $4.2 \times 10^3$ 91. Which one is not a fossil fuel?  
 (A) Coal  
 (B) Biomass  
 (C) Petroleum  
 (D) None of these92. If a force F is applied on a body and it moves with a velocity v, then the value of the power will be—  
 (A)  $F/v$   
 (B)  $F \times v$   
 (C)  $Fv^2$   
 (D)  $F \times v^2$ 93. A liquid A of mass  $10\text{ g}$  and at  $120^\circ\text{C}$  is poured in a liquid B at  $20^\circ\text{C}$ . The final temperature recorded is  $40^\circ\text{C}$ . The initial mass of liquid B is (If initial capacity of liquid A is  $0.8\text{ J/g}^\circ\text{C}$  and of liquid B  $4.2\text{ J/g}^\circ\text{C}$ )—  
 (A)  $70\text{ g}$   
 (B)  $80.0\text{ g}$   
 (C)  $70\text{ g}$   
 (D)  $76.1\text{ g}$ 94. Among the elements Li, Na, Mg and Al, the lowest value of ionization potential of certain will be—  
 (A) Li  
 (B) Na  
 (C) Mg  
 (D) Al95. On moving from left to right in a period of periodic table, the number of shell—  
 (A) remains the same  
 (B) increases  
 (C) decreases  
 (D) first increases then decreases96. The force of friction on a body kept on the table top does not depend on—  
 (A) nature of force  
 (B) material of the body  
 (C) weight of the body  
 (D) area of contact97. Wheels of a moving cycle ~~have~~ have rotational and translational motion  
 (A) have translational motion only  
 (B) have rotational motion only  
 (C) have both motion only  
 (D) None of the above98. ~~Margin~~ margin of unit of—  
 (A)  $\text{J}/\text{kg}$   
 (B) force  
 (C) power  
 (D) energy90. Specific heat capacity of copper is  $0.1\text{ cal/gm}$ . Its value in  $\text{J/kg}^\circ\text{C}$  is—(A)  $0.84 \times 10^3$   
 (B)  $0.42 \times 10^4$   
 (C)  $0.24 \times 10^3$   
 (D)  $4.2 \times 10^3$ 91. Which one is not a fossil fuel?  
 (A) Coal  
 (B) Biomass  
 (C) Petroleum  
 (D) None of these92. If a force F is applied on a body and it moves with a velocity v, then the value of the power will be—  
 (A)  $F/v$   
 (B)  $F \times v$   
 (C)  $Fv^2$   
 (D)  $F \times v^2$ 93. A liquid A of mass  $10\text{ g}$  and at  $120^\circ\text{C}$  is poured in a liquid B at  $20^\circ\text{C}$ . The final temperature recorded is  $40^\circ\text{C}$ . The initial mass of liquid B is (If initial capacity of liquid A is  $0.8\text{ J/g}^\circ\text{C}$  and of liquid B  $4.2\text{ J/g}^\circ\text{C}$ )—  
 (A)  $70\text{ g}$   
 (B)  $80.0\text{ g}$   
 (C)  $70\text{ g}$   
 (D)  $76.1\text{ g}$ 94. Among the elements Li, Na, Mg and Al, the lowest value of ionization potential of certain will be—  
 (A) Li  
 (B) Na  
 (C) Mg  
 (D) Al95. On moving from left to right in a period of periodic table, the number of shell—  
 (A) remains the same  
 (B) increases  
 (C) decreases  
 (D) first increases then decreases96. The force of friction on a body kept on the table top does not depend on—  
 (A) nature of force  
 (B) material of the body  
 (C) weight of the body  
 (D) area of contact97. Wheels of a moving cycle ~~have~~ have rotational and translational motion  
 (A) have translational motion only  
 (B) have rotational motion only  
 (C) have both motion only  
 (D) None of the above98. ~~Margin~~ margin of unit of—  
 (A)  $\text{J}/\text{kg}$   
 (B) force  
 (C) power  
 (D) energy90. Specific heat capacity of copper is  $0.1\text{ cal/gm}$ . Its value in  $\text{J/kg}^\circ\text{C}$  is—(A)  $0.84 \times 10^3$   
 (B)  $0.42 \times 10^4$   
 (C)  $0.24 \times 10^3$   
 (D)  $4.2 \times 10^3$ 91. Which one is not a fossil fuel?  
 (A) Coal  
 (B) Biomass  
 (C) Petroleum  
 (D) None of these92. If a force F is applied on a body and it moves with a velocity v, then the value of the power will be—  
 (A)  $F/v$   
 (B)  $F \times v$   
 (C)  $Fv^2$   
 (D)  $F \times v^2$ 93. A liquid A of mass  $10\text{ g}$  and at  $120^\circ\text{C}$  is poured in a liquid B at  $20^\circ\text{C}$ . The final temperature recorded is  $40^\circ\text{C}$ . The initial mass of liquid B is (If initial capacity of liquid A is  $0.8\text{ J/g}^\circ\text{C}$  and of liquid B  $4.2\text{ J/g}^\circ\text{C}$ )—  
 (A)  $70\text{ g}$   
 (B)  $80.0\text{ g}$   
 (C)  $70\text{ g}$   
 (D)  $76.1\text{ g}$ 94. Among the elements Li, Na, Mg and Al, the lowest value of ionization potential of certain will be—  
 (A) Li  
 (B) Na  
 (C) Mg  
 (D) Al95. On moving from left to right in a period of periodic table, the number of shell—  
 (A) remains the same  
 (B) increases  
 (C) decreases  
 (D) first increases then decreases96. The force of friction on a body kept on the table top does not depend on—  
 (A) nature of force  
 (B) material of the body  
 (C) weight of the body  
 (D) area of contact97. Wheels of a moving cycle ~~have~~ have rotational and translational motion  
 (A) have translational motion only  
 (B) have rotational motion only  
 (C) have both motion only  
 (D) None of the above98. ~~Margin~~ margin of unit of—  
 (A)  $\text{J}/\text{kg}$   
 (B) force  
 (C) power  
 (D) energy90. Specific heat capacity of copper is  $0.1\text{ cal/gm}$ . Its value in  $\text{J/kg}^\circ\text{C}$  is—(A)  $0.84 \times 10^3$   
 (B)  $0.42 \times 10^4$   
 (C)  $0.24 \times 10^3$   
 (D)  $4.2 \times 10^3$ 91. Which one is not a fossil fuel?  
 (A) Coal  
 (B) Biomass  
 (C) Petroleum  
 (D) None of these92. If a force F is applied on a body and it moves with a velocity v, then the value of the power will be—  
 (A)  $F/v$   
 (B)  $F \times v$   
 (C)  $Fv^2$   
 (D)  $F \times v^2$ 93. A liquid A of mass  $10\text{ g}$  and at  $120^\circ\text{C}$  is poured in a liquid B at  $20^\circ\text{C}$ . The final temperature recorded is  $40^\circ\text{C}$ . The initial mass of liquid B is (If initial capacity of liquid A is  $0.8\text{ J/g}^\circ\text{C}$  and of liquid B  $4.2\text{ J/g}^\circ\text{C}$ )—  
 (A)  $70\text{ g}$   
 (B)  $80.0\text{ g}$   
 (C)  $70\text{ g}$   
 (D)  $76.1\text{ g}$ 94. Among the elements Li, Na, Mg and Al, the lowest value of ionization potential of certain will be—  
 (A) Li  
 (B) Na  
 (C) Mg  
 (D) Al95. On moving from left to right in a period of periodic table, the number of shell—  
 (A) remains the same  
 (B) increases  
 (C) decreases  
 (D) first increases then decreases

8. (A) If line PQ is parallel to line BC of triangle.



Then,

$$\begin{aligned} 4x - 3(0) + 12 &= 0 \\ 4x - 12 &= 0 \end{aligned}$$

$$\frac{AP}{PB} = \frac{AQ}{QC}$$

Then,

13. (A) When two lines are perpendicular,

Then

$m_{1}m_2 = -1$

$m_1 = \frac{3}{2}$

$m_2 = -\frac{2}{3}$

$K = 1$

$K = 3$

$K = 1$

Height of cylinder = 16 cm  
 $\frac{2\pi r}{2\pi} = 22$  cm  
 and

$$r = \frac{11}{2}$$

$$\text{Volume of cylinder} = \pi r^2 h$$

$$= \pi \times \left(\frac{11}{2}\right)^2 \times 16$$

$$= \frac{11 \times 11 \times 16}{4}$$

$$= 616 \text{ cm}^3$$

22. (B) Angle formed at semicircle is  $90^\circ$



$$25. (\text{A})$$

$$N = \left[ \begin{array}{cc} 1 & -2 \\ -3 & 4 \end{array} \right]_{2 \times 2}$$

$$26. (\text{D}) \text{ mean} = \frac{5+3+0.5+4.5+b+8.5+9.5}{10}$$

$$31. (\text{D}) \text{ If}$$

$$\frac{x}{y} = \frac{4}{3}$$

$$\text{Then } \frac{5x+8y}{6x-7y} = \frac{y(x+y+8)}{y(6x-7)}$$

$$b = 49 - 31$$

Radius = 10 cm

Area of the shaded portion

$$\begin{aligned} &= \frac{\pi}{2} \times 100 - \frac{1}{2} \times 12 \times 16 \\ &= 50\pi - 96 \\ &= 157.1 - 96 \\ &= 61.1 \text{ cm}^2 \end{aligned}$$

23. (B) PS = PQ = 6 cm,  
 $PQ = QR = RS = 2 \text{ cm}$

Perimeter of shaded portion

$$= \pi R + \pi r_1 + \pi r_2$$

$$= \pi [3+2+1]$$

$$\begin{cases} \pi R = \frac{PS}{2} = 3 \text{ cm} \\ r_1 = \frac{QS}{2} = 2 \text{ cm} \\ r_2 = \frac{PQ}{2} = \frac{3}{2} = 1.5 \text{ cm} \end{cases}$$

$$\begin{aligned} &\text{C.P. for B} = 12,500 \\ &\text{C.P. for C} = 15,300 \\ &\text{Cost of machine after profit of C} \\ &= 13,300 + 1300 \\ &= 14,600 \end{aligned}$$

$$\text{VAT} = \frac{14600 \times 8}{100}$$

24. (A)



Area of rectangular sheet =  $30 \times 10$   
 $= 300 \text{ cm}^2$

Diameter of disc = 1 cm

So, 300 discs can be made from sheet.



From pythagoras theorem  
 $AB^2 = AP^2 + PB^2$

i. Locus of point P is right angled triangle.

30. (C) Let the marked price be x

$$x = \frac{109}{100} = 109.22$$

$$x = \frac{40.221}{100}$$

$$x = ₹ 36.900$$

35. (A) Equation,  $x^2 - 3x - 28 = 0$   
 sum of roots =  $-\frac{b}{a} = \frac{3}{1} = 3$

36. (D) Let the number = x, y  
 $x+y = 3 \cdot 5$   
 $x+y = 15$

$$x+y = 2$$

$$5K+8 = 2$$

$$5K+24 = 10K+16$$

$$K = 8$$

$$x = 24, y = 40$$

37. (A) Let x be added in each number  
 then

$$\frac{4+x}{8+x} = \frac{16+x}{26+x}$$

$$\begin{aligned} &104 + 30x + x^2 = 128 + 24x + x^2 \\ &6x = 24 \\ &x = 4 \end{aligned}$$

32. (D)

$$A = 5000 \left(1 + \frac{6}{100}\right)^3$$

33. (C)

$$38. (\text{A})$$

39. (A)

$$\cot^2 \theta (1 - \cos^2 \theta) = \cot^2 \theta (\sin^2 \theta)$$

40. (A)

$$\begin{aligned} &(5x-3)(x^2+4+4x) \\ &= 5x^3+20x^2+20x^3-3x^2-12-1 \\ &= 5x^3+17x^2+8x-12 \\ &\text{coefficient of } x^2 = 17, \\ &\text{coefficient of } x = 8 \end{aligned}$$

41. (B)

$$\left(x - \frac{2}{x}\right) \left(x^2 + 2 + \frac{4}{x^2}\right)$$

42. (A)

$$= x^3 + 2x + \frac{4}{x} - 2x - \frac{4}{x}$$

43. (C)

$$\text{S.I.} = \frac{250 \times 6 \times 24}{100} = ₹ 360$$

44. (B)

$$= x^3 - \frac{8}{x^3}$$

34. (B) Let the number = x

$$\begin{aligned} &x + \frac{1}{x} = 17 \\ &\frac{x^2+1}{x} = 17 \\ &x^2+1 = 17x \end{aligned}$$

$$4x^2+4 = 17x$$

$$4x^2-17x+4 = 0$$

$$4x(x-4)-13x+4 = 0$$

$$(x-4)(4x-1) = 0$$


$$x = 4, x = \frac{1}{4}$$



85. (C) Intensity ( $I$ )  $\propto a^2$ .

86. (B) Intensity of sound is measured in decibel (dB).

87. (A)



Let  $R_1$  and  $R_2$  are in series, then

$$R' = R_1 + R_2 = 3 + 3 \\ = 6\Omega$$

When  $R'$  and  $R_3$  are in parallel, then

$$P_{PQ} = \frac{R' \times R_3}{R' + R_3} \\ = \frac{6 \times 3}{6 + 3} \\ = \frac{18}{9} = 2\Omega$$

$R_4$ ,  $R_{PQ}$ ,  $R_5$  will be in series

Resistance between X and Y

$$R = R_4 + R_{PQ} + R_5 \\ = 3 + 2 + 3 = 8\Omega$$

88. (B) No. of cells ( $n$ ) = 4

$$E = 2 \text{ volt}$$

$$r = 0.1 \Omega$$

$$I = 2A$$

When the resistances  $1.6 \Omega$  and  $R$  are connected to ammeter, they will be in series. Hence,

$$R' = R + 1.6 \Omega$$

$$\text{Current } I = \frac{nE}{nr + R'} \\ I = \frac{4 \times 2}{4 \times 0.1 + R + 1.6} \\ I = \frac{4}{R + 0.4 + 1.6} \\ I = \frac{4}{R + 2.0}$$

$$\text{or } R + 2.0 = 4$$

$$\text{Resistance } R = 4 - 2 = 2\Omega$$

89. (C) The power rated on heater  $3 \text{ kV}$  and  $250$  volt

$$\text{Namely Power (P)} = 3 \text{ kV} = 3 \times 10^3 \text{ V}$$

$$\text{Potential difference (V)}$$

$$= 250 \text{ volt}$$

$$t = 5 \text{ hrs.}$$

Consumed electrical energy

$$= \frac{\text{Total power (in watt)} \times \text{Total time (in hrs.)}}{1000}$$

$$= \frac{3 \times 1000 \times 5}{1000} = 15 \text{ joules}$$

The cost of running the heater continuously for 5 hours

$$= 15 \times 5 = ₹ 75$$

90. (B)  $1 \text{ cal/gm} = 4.2 \times 10^3 \text{ J/kg}^{\circ}\text{C}$   
 $0.1 \text{ cal/gm} = 0.42 \times 10^3 \text{ J/kg}^{\circ}\text{C}$

91. (B) Hydrogen is not a greenhouse gas.

92. (D) Let the mass of liquid B is  $m_2 \text{ gm}$ . As per the principle of colorimetry,

$$\text{Heat lost} = \text{Heat gained}$$

$$100 \times 0.8 \times (120 - 40) = m_2 \times 4.2 \times (40 - 20)$$

$$100 \times 0.8 \times 80 = m_2 \times 4.2 \times 20$$

$$\text{or } m_2 = \frac{100 \times 0.8 \times 80}{4.2 \times 20} \\ = 76.19 \text{ gm}$$

93. (B) Element : Li Na Mg Al  
 Ionization potential ( $\text{kJ mol}^{-1}$ ) : 520.3 495.8 737.6 577.4

94. (A) On moving from left to right in a period of the periodic table, the number of shells remain the same.

95. (D)

96. (D) The force of friction does not depend on the area of contact.

97. (A) The wheels of a running cycle have both rotational and translational motions.

98. (A,D) Erg is the unit of work and energy both.  
 $1 \text{ Erg} = 1 \times 10^{-4} \text{ joule}$

99. (B) Biomass is not a fossil fuel.

100. (A) Power ( $P$ ) =  $F \times v$