

MATHEMATICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer :

1. If $2x^2 + (\cos\theta)x - 1 = 0$, $\theta \in [0, 2\pi]$ has roots α and β . Then the sum of maximum and minimum value of $\alpha^4 + \beta^4$.

- (1) $\frac{25}{16}$
- (2) $\frac{9}{16}$
- (3) $\frac{41}{16}$
- (4) $\frac{8}{17}$

Answer (1)

Sol. $\alpha + \beta = \frac{-\cos\theta}{2}$

$\alpha\beta = \frac{-1}{2} \Rightarrow \alpha^2\beta^2 = \frac{1}{4}$

$\Rightarrow \alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$

$= \frac{\cos^2\theta + 1}{4}$

$\alpha^4 + \beta^4 = (\alpha^2 + \beta^2)^2 - 2\alpha^2\beta^2$

$= \left(\frac{\cos^2\theta + 1}{4}\right)^2 - \frac{1}{2}$

For minimum, $\cos\theta = 0$

For maximum, $\cos\theta = 1$

$\Rightarrow \text{Minimum} = 1 - \frac{1}{2} = \frac{1}{2}$

Maximum $= \frac{25}{16} - \frac{1}{2} = \frac{17}{16}$

$\Rightarrow \text{Sum} = \frac{1}{2} + \frac{17}{16} = \frac{25}{16}$

2. If $\theta \in [0, 2\pi]$ satisfying the system of equations $2\sin^2\theta = \cos 2\theta$ and $2\cos^2\theta = 3\sin\theta$. Then the sum of all real values of θ is

- (1) $\frac{3\pi}{2}$
- (2) π
- (3) $\frac{\pi}{2}$
- (4) $\frac{5\pi}{6}$

Answer (2)

Sol. $2\sin^2\theta = \cos 2\theta$

$2\cos^2\theta = 3\sin\theta$

\Rightarrow Adding,

$2 = 1 - 2\sin^2\theta + 3\sin\theta$

$\Rightarrow 2\sin^2\theta - 3\sin\theta + 1 = 0$

$2\sin^2\theta - 2\sin\theta - \sin\theta + 1 = 0$

$2\sin\theta(\sin\theta - 1) - 1(\sin\theta - 1) = 0$

$\sin\theta = 1, \frac{1}{2}$ but $2\sin^2\theta = \cos 2\theta = 2$

but not is not possible

$\Rightarrow \theta = \frac{\pi}{6}, \left(\pi - \frac{\pi}{6}\right)$

$\Rightarrow \text{Sum of all values} = \frac{\pi}{6} + \pi - \frac{\pi}{6} = \pi$

3. Let $A = \{1, 2, 3, 4\}$ and $B = \{1, 4, 9, 16\}$.

If $f: A \rightarrow B$, number of many-one functions from A to B are

- (1) 24
- (2) 232
- (3) 256
- (4) 252

Answer (2)

Delivering Champions Consistently

Aakash
Medical | IIT-JEE | Foundations

JEE (Advanced) 2024						JEE (Main) 2024		
 AIR 25 Rishi Shekher Shukla 2 Year Classroom	 AIR 67 Krishna Sai Shishir 2 Year Classroom	 AIR 78 Abhishek Jain 2 Year Classroom	 AIR 93 Hardik Aggarwal 2 Year Classroom	 AIR 95 Ujjwal Singh 2 Year Classroom	 AIR 98 Rishabh Aggarwal 2 Year Classroom	 Karnataka Topper 100 PERCENTILE Sanyal Jais 2 Year Classroom	 Telangana Topper 100 PERCENTILE M Sai Divya Teja Reddy 2 Year Classroom	 Telangana Topper 100 PERCENTILE Rishi Shekher Shukla 2 Year Classroom

Sol. $n(A) = 4$

$n(B) = 4$

Number of many one functions =

Total functions – Number of one-one function

$$= 4^4 - 4! = 232$$

4. 4 boys and 3 girls are to be seated in a row such that all girls seat together and two particular boys B_1 and B_2 are not adjacent to each other. Then the number of ways in which this arrangement can be done.

(1) 432

(2) 430

(3) 516

(4) 1002

Answer (1)

Sol. 4 boys and 3 girls.

All girls together $\Rightarrow \boxed{G_1 G_2 G_3}$

3 girls and 2 boys can be seated in $3! \cdot 3!$ ways

Now B_1 and B_2 go into spaces.

$$\Rightarrow {}^4C_2 \times 2! \times 3! \times 3! = 432$$

5. If the sum $\sum_{r=0}^{30} \frac{r^2 \binom{30}{r}}{\binom{30}{r-1}} = \alpha \cdot 2^{29}$, then α is equal

to

(1) 225

(2) 465

(3) 345

(4) 425

Answer (2)

Sol. $\frac{r^2 \cdot 30!}{(30-r)!r!} \cdot \frac{30!}{(30-r)!r!} \times \frac{(r-1)!(31-r)!}{30!}$

$$= \frac{30!(31-r)}{(r-1)!(30-r)!}$$

$$= \frac{30(31-r)29!}{(r-1)!(30-r)!}$$

$$\Rightarrow \sum_{r=0}^{30} \frac{r^2 \binom{30}{r}}{\binom{30}{r-1}} = 30 \sum_{r=0}^{30} (31-r)^{29} C_{r-1} = 30 \sum_{r=0}^{30} (31-r)^{29} C_{30-r}$$

$$= 30 \sum_{r=0}^{30} [(30-r)+1]^{29} C_{30-r}$$

$$= 30 \sum_{r=0}^{30} \frac{29}{(30-r)} (30-r)^{28} C_{29-r} + 30 \sum_{r=0}^{30} 29 C_{30-r}$$

$$= 30 \cdot 29 \cdot 2^{28} + 30 \cdot 2^{29}$$

$$= 30 \cdot 2^{28} (29 + 2) = (31 \times 15) \cdot 2^{29}$$

6. Consider a function $f(x) = \int_0^{x^2} \frac{t^2 - 8t + 15}{e^t} dt$. The

number of points of extrema are

(1) 3

(2) 5

(3) 7

(4) 9

Answer (2)

Sol. $\therefore f(x) = \int_0^{x^2} \frac{t^2 - 8t + 15}{e^t} dt$

$$f'(x) = \frac{2x(x^4 - 8x^2 + 15)}{e^{x^2}}$$

$$= \frac{2x(x^2 - 5)(x^2 - 3)}{e^{x^2}}$$

The extremum value of $f(x)$ are $x = 0, \pm\sqrt{5}, \pm\sqrt{3}$

\therefore Number of extremum points are 5.

7. Let A and B are two events such that

$P(A \cap B) = \frac{1}{10}$ and $P(A/B)$ and $P(B/A)$ are the

roots of the equation $12x^2 - 7x + 1 = 0$, then

$\frac{P(\bar{A} \cup \bar{B})}{P(\bar{A} \cap \bar{B})}$ is equal to

(1) $\frac{4}{9}$

(2) $\frac{9}{4}$

(3) $\frac{3}{2}$

(4) $\frac{2}{3}$

Answer (2)

Sol. $P(A \cap B) = \frac{1}{10}$

Delivering Champions Consistently

JEE (Advanced) 2024

- AIR 25: Rishi Shekher Shukla (2 Year Classroom)
- AIR 67: Krishna Sai Shishir (2 Year Classroom)
- AIR 78: Abhishek Jain (2 Year Classroom)
- AIR 93: Hardik Aggarwal (2 Year Classroom)
- AIR 95: Ujjwal Singh (2 Year Classroom)
- AIR 98: Rachit Aggarwal (2 Year Classroom)

JEE (Main) 2024

- Karnataka Topper: AIR 1: Sarvi Jais (1 Year Classroom)
- Telangana Topper: AIR 15: M Sai Divya Teja Reddy (2 Year Classroom)
- Telangana Topper: AIR 19: Rishi Shekher Shukla (2 Year Classroom)

$$P(A/B) + P(B/A) = \frac{7}{12}$$

$$\text{and } P(A/B) \cdot P(B/A) = \frac{1}{12}$$

$$\frac{P(A \cup B)}{P(B)} \cdot \frac{P(A \cap B)}{P(A)} = \frac{1}{12}$$

$$\Rightarrow P(A) \cdot P(B) = 12 \left(\frac{1}{12} \right)^2 = \frac{12}{100}$$

$$P(A \cap B) \left[\frac{1}{P(A)} + \frac{1}{P(B)} \right] = \frac{7}{12}$$

$$P(A) + P(B) = \frac{7}{10}$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{7}{10} - \frac{1}{10} = \frac{6}{10}$$

$$\frac{P(\bar{A} \cup \bar{B})}{P(\bar{A} \cap \bar{B})} = \frac{P(\overline{A \cap B})}{P(\overline{A \cup B})} = \frac{1 - P(A \cap B)}{1 - P(A \cup B)}$$

$$\frac{1 - \frac{1}{10}}{1 - \frac{6}{10}} = \frac{\frac{9}{10}}{\frac{4}{10}} = \frac{9}{4}$$

8. Number of terms in an arithmetic progression is $2n$. Sum of terms occurring at even places is 40 and sum of terms occurring at odd places is 55. If the first term exceeds the last term by 27, then n equals to

- (1) 3 (2) 5
(3) 7 (4) 4

Answer (2)

Sol. Let the AP be

$$a, a + d, a + 2d, \dots, a + (2n - 1)d$$

Now given that

$$(a + d) + (a + 3d) + \dots + (a + (2n - 1)d) = 40$$

$$na + n^2d = 40 \quad \dots(1)$$

$$\text{Also } a + (a + 2d) + (a + 4d) + \dots + (a + (2n - 2)d) = 55$$

$$na + dn(n - 1) = 55 \quad \dots(2)$$

$$\text{Also } a - (a + (2n - 1)d) = 27$$

$$-(2n - 1)d = 27 \quad \dots(3)$$

$$d = \frac{-27}{2n - 1}$$

$$(2) - (1)$$

$$dn(n - 1) - n^2d = 15$$

$$d[n^2 - n - n^2] = 15$$

$$\left(\frac{-27}{2n - 1} \right) (-n) = 15$$

$$27n = 30n - 15$$

$$15 = 3n$$

$$n = 5$$

9. The perpendicular distance of point $P(3, 4, 5)$ from the line

$$\vec{r} = 2\hat{i} - \hat{j} + \hat{k} + \lambda(4\hat{i} - \hat{j} + 5\hat{k}) \text{ is}$$

(1) $\sqrt{\frac{19}{42}}$ (2) $\sqrt{\frac{19}{21}}$

(3) $\sqrt{\frac{42}{19}}$ (4) $\sqrt{\frac{21}{19}}$

Answer (1)

Sol. $P(3, 4, 5)$

$$L: \vec{r} = \vec{r} = 2\hat{i} - \hat{j} + \hat{k} + \lambda(4\hat{i} - \hat{j} + 5\hat{k})$$

Any point on L can be. $A(2 + 4\lambda, -1 - \lambda, 1 + 5\lambda)$

$$\text{Now } \overline{AP} \cdot (4\hat{i} - \hat{j} + 5\hat{k}) = 0$$

$$((4\lambda - 1)\hat{i} - (\lambda + 5)\hat{j} + (5\lambda - 4)\hat{k}) \cdot (4\hat{i} - \hat{j} + 5\hat{k}) = 0$$

$$16\lambda + 4 + \lambda + 5 + 25\lambda - 20 = 0$$

$$42\lambda = 19$$

$$\lambda = \frac{19}{42}$$

$$\text{Now } |AP| = \sqrt{(4\lambda - 1)^2 + (\lambda + 5)^2 + (5\lambda + 4)^2}$$

$$= \sqrt{\frac{19}{42}}$$

Delivering Champions Consistently

Aakash
Medical IIT-JEE Foundations

JEE (Advanced) 2024

AIR 25 Rishj Shekher Shukla 2 Year Classroom	AIR 67 Krishna Sai Shishir 2 Year Classroom	AIR 78 Abhishek Jain 2 Year Classroom	AIR 93 Hardik Aggarwal 2 Year Classroom	AIR 95 Ujjwal Singh 2 Year Classroom	AIR 98 Rishit Aggarwal 2 Year Classroom
---	--	--	--	---	--

JEE (Main) 2024

Karnataka Topper AIR 1 Savitri Jais 2 Year Classroom	Telangana Topper AIR 15 M Sai Divya Teja Reddy 2 Year Classroom	Telangana Topper AIR 19 Rishi Shekher Shukla 2 Year Classroom
---	--	--

10. In the expansion of $(x + \sqrt{x^3 - 1})^5 + (x - \sqrt{x^3 - 1})^5$, where α, β, γ and δ are the coefficient of x, x^3, x^5 and x^7 respectively. If $\alpha u - \beta v = 18, \gamma u + \delta v = 20$ then $u + v$ equal to.

- (1) $\frac{-14}{15}$ (2) $\frac{-13}{15}$
 (3) $\frac{-3}{5}$ (4) $\frac{-2}{3}$

Answer (1)

Sol. $(x + \sqrt{x^3 - 1})^5 + (x - \sqrt{x^3 - 1})^5$

$$\left[\int_0^3 (x)^5 + \int_1^5 x^4 \sqrt{x^3 - 1} + \int_2^5 x^3 (\sqrt{x^3 - 1})^2 + \dots \right] +$$

$$\left[\int_0^5 x^5 - \int_1^5 x^4 \sqrt{x^3 - 1} + \int_2^5 x^3 (\sqrt{x^3 - 1})^2 + \dots \right]$$

$$= 2 \left[x^5 + \int_2^5 x^3 (x^3 - 1) + \int_4^5 x (x^3 - 1)^2 \right]$$

$$= 2[x^5 + 10x^6 - 10x^3 + 5x(x^6 + 1 - 2x^3)]$$

$$= 10x^7 + 20x^6 + 2x^5 - 20x^4 - 20x^3 + 10x$$

 Coefficient of $x = 10 = \alpha$
 Coefficient of $x^3 = -20 = \beta$
 Coefficient of $x^5 = 2 = \gamma$
 Coefficient of $x^7 = 10 = \delta$
 $10u + 20v = 18 \quad \dots(i)$
 $2u + 10v = 20$
 $u = \frac{-11}{3}$
 $v = \frac{41}{15}$
 $u + v = \frac{41}{15} - \frac{11}{3}$
 $= \frac{-14}{15}$

11. Let $A(6, 8), B(10 \cos \alpha, -10 \sin \alpha)$ and $C(-10 \sin \alpha, -10 \cos \alpha)$ be 3 points and if orthocentre of the triangle ABC is $(0, 9)$ then $100 \sin^2 \alpha$ is equal to

- (1) $\frac{25}{4}$ (2) 25
 (3) $\frac{15}{4}$ (4) $\frac{5}{4}$

Answer (1)

Sol. Notice, origin is equidistance form A, B and C

$\Rightarrow (0, 0)$ is circumcentre



Since centroid divides orthocentre and circumcentre in 2 : 1 ratio.

$$\Rightarrow \frac{6 + 10 \cos \alpha + (-10 \sin \alpha)}{3} = \frac{2(0) + 1(0)}{3} = 0$$

$$\Rightarrow 15 \sin \alpha - 10 \cos \alpha = 6$$

Alos $\frac{8 - 10 \sin \alpha - 10 \cos \alpha}{3} = \frac{2(0) + 1(9)}{3} = 3$

$$\Rightarrow 8 - 10 \sin \alpha - 10 \cos \alpha = 9$$

$$10(\sin \alpha + \cos \alpha) = -1$$

$$10(\sin \alpha - \cos \alpha) = -6$$

$$20 \sin \alpha = 5$$

$$10 \sin \alpha = \frac{5}{2}$$

$$100 \sin^2 \alpha = \frac{25}{4}$$

12. If z be a complex number such that $|z - 3| \leq 1$, then the equation of line with largest slope passing through origin and z

- (1) $x - 2\sqrt{2}y = 0$ (2) $x + 2\sqrt{2}y = 0$
 (3) $2\sqrt{2}x + y = 0$ (4) $2\sqrt{2}x - y = 0$


Answer (1)

Delivering Champions Consistently

JEE (Advanced) 2024
 AIR 25 Rishj Shekher Shukla (2 Year Classroom)
 AIR 67 Krishna Sai Shishir (2 Year Classroom)
 AIR 78 Abhishek Jain (2 Year Classroom)
 AIR 93 Hardik Aggarwal (2 Year Classroom)
 AIR 95 Ujjwal Singh (2 Year Classroom)
 AIR 98 Rachit Aggarwal (2 Year Classroom)

JEE (Main) 2024
 Karnataka Topper: AIR 1 Sarvi Jain (2 Year Classroom)
 Telangana Topper: AIR 15 M Sai Divya Teja Reddy (2 Year Classroom)
 Telangana Topper: AIR 19 Rishi Shekher Shukla (2 Year Classroom)

Sol.



$$\tan\theta = \frac{1}{\sqrt{8}} = \frac{1}{2\sqrt{2}}$$

Therefore, equation of line with maximum slope is

$$(y - 0) = \frac{1}{2\sqrt{2}}(x - 0)$$

$$\Rightarrow y = \frac{x}{2\sqrt{2}}$$

13. A relation R is defined on set A , $A = \{1, 2, 3\}$ and $R = \{(1, 2), (2, 3)\}$. Elements are added such that R becomes reflexive and transitive but not symmetric. Find the number of such relations.

- (1) 3
(2) 4
(3) 2
(4) 9

Answer (1)

Sol. Transitivity

$$(1, 2) \in R, (2, 3) \Rightarrow (1, 3) \in R$$

$$(1, 1), (2, 2), (3, 3) \in R$$

$$(2, 1) \quad (3, 2) \quad (3, 1)$$

(3, 1) cannot be taken.

- (2, 1) taken and (3, 2) not taken.
- (3, 2) taken and (2, 1) not taken.
- Both not taken.

Therefore 3 relations are possible.

14.
15.
16.
17.
18.

19.

20.

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

21. Let \vec{a} and \vec{b} be two unit vectors such that angle between \vec{a} and \vec{b} is $\frac{\pi}{3}$. If $\lambda\vec{a} + 3\vec{b}$ and $2\vec{a} + \lambda\vec{b}$ are perpendicular to each other, then the product of all possible values of λ is _____

Answer (6)

Sol. $|\vec{a}| = 1, |\vec{b}| = 1$

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \frac{\pi}{3} = \frac{1}{2}$$

$\lambda\vec{a} + 3\vec{b}$ and $2\vec{a} + \lambda\vec{b}$ are perpendicular

$$\Rightarrow (\lambda\vec{a} + 3\vec{b}) \cdot (2\vec{a} + \lambda\vec{b}) = 0$$

$$\Rightarrow 2\lambda + 3\lambda + (\lambda^2 + 6)\vec{a} \cdot \vec{b} = 0$$

$$\Rightarrow 5\lambda + (\lambda^2 + 6)\left(\frac{1}{2}\right) = 0$$

$$\Rightarrow 10\lambda + \lambda^2 + 6 = 0$$

$$\Rightarrow \lambda^2 + 10\lambda + 6 = 0$$

Product of possible values of $\lambda = 6$

22. If A is the 3×3 matrix of order 3×3 , such that $\det(A) = \frac{1}{2}$, $\text{tr}(A) = 10$ and B be another matrix of order 3×3 and defined as $B = \text{adj}(\text{adj}(2A))$, then $\det(B) + \text{tr}(B)$ is equal to (where $\text{tr}(A)$ denotes trace of matrix A)

Answer (336)

Sol. $B = \text{adj}(\text{adj}(2A))$

$$B = |2A|^{n-2} (2A), [\text{Using } \text{adj}(\text{adj } P) = |P|^{n-2} \cdot P],$$

for $n = 3$

$$= |2A|(2A)$$

$$= 2^3|A|(2A)$$



Delivering Champions Consistently

JEE (Advanced) 2024						JEE (Main) 2024		
 AIR 25 Rishi Shekher Shukla 2 Year Classroom	 AIR 67 Krishna Sai Shishir 2 Year Classroom	 AIR 78 Abhishek Jain 2 Year Classroom	 AIR 93 Hardik Aggarwal 2 Year Classroom	 AIR 95 Ujwal Singh 2 Year Classroom	 AIR 98 Rachit Aggarwal 2 Year Classroom	 AIR 1 Savitri Jais 2 Year Classroom	 AIR 15 M Sai Divya Teja Reddy 2 Year Classroom	 AIR 19 Rishi Shekher Shukla 2 Year Classroom

$$= 8 \times \frac{1}{2}(2A)$$

$$= 4(2A)$$

$$B = 8A$$

$$|B| = |8A|$$

$$= 8^3|A|$$

$$|B| = 8^3 \times \frac{1}{2} = 256$$

$$B = 8A$$

[each element is multiplied 8 times]

$$\text{tr}(B) = 8\text{tr}(A)$$

$$= 80$$

$$|B| + \text{tr}(B) = 256 + 80$$

$$= 336$$

23. Consider two curves $E_1: \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ with eccentricity e_1 and $E_2: \frac{x^2}{A^2} + \frac{y^2}{B^2} = 1$ with eccentricity e_2 . If $\frac{e_1}{e_2} = \frac{1}{3}$ and distance between foci of both curves is $2\sqrt{3}$ and $a - A = 4$, then the sum of lengths of latus rectum of both curves is

Answer (12)

Sol. Since, distance between foci

for $E_1 = 2ae_1$, for $E_2 = 2Ae_2$

$$\Rightarrow 2ae_1 = 2\sqrt{3} = 2Ae_2$$

$$\Rightarrow ae_1 = Ae_2$$

$$\Rightarrow \frac{e_1}{e_2} = \frac{A}{a} = \frac{1}{3} \Rightarrow a = 3A$$

$$\text{Also } a - A = 4 \Rightarrow A = 2, a = 6$$

$$\text{Now, } 2(6)e_1 = 2\sqrt{3} \Rightarrow e_1 = \frac{\sqrt{3}}{6}$$

$$2(2)e_2 = 2\sqrt{3} \Rightarrow e_2 = \frac{\sqrt{3}}{2}$$

$$e_1^2 = \frac{1-b^2}{a^2} = \frac{3}{36} = \frac{1-b^2}{36} \Rightarrow \frac{b^2}{36} = \frac{33}{36}$$

$$b^2 = 33$$

$$e_2^2 = \frac{1-B^2}{A^2} = \frac{3}{4} = \frac{1-B^2}{4} \Rightarrow B^2 = 1$$

$$\text{Length of latus rectum of } E_1 = \frac{2b^2}{a}$$

$$\text{Length of latus rectum of } E_2 = \frac{2B^2}{A}$$


\Rightarrow Sum of lengths of latus rectum

$$= \frac{2(33)}{6} + \frac{2(1)}{2} = 11 + 1 = 12$$

24. The number of maximum number of common tangents to the curves $y = (x - 2)^2$ and $y^2 = 16 - 8x$ is

Answer (1)

Sol.



Clearly, one tangent is possible. Based on the graphs of these parabola.

25. Let $P(10, -2, -1)$ and Q be the point of perpendicular drawn from point $R(1, 7, 6)$ on the line joining the points $(2, -5, 11)$ and $(-6, 7, -5)$. Then the length PQ is

Answer (13)

$$\text{Sol. } L_1: \frac{x-2}{8} = \frac{y+5}{-12} = \frac{z-11}{16}$$

$$L_1: \frac{x-2}{2} = \frac{y+5}{-3} = \frac{z-11}{4}$$

Delivering Champions Consistently

JEE (Advanced) 2024						JEE (Main) 2024		
AIR 25	AIR 67	AIR 78	AIR 93	AIR 95	AIR 98	100 Percentile Rank	100 Percentile Rank	100 Percentile Rank
Rishi Shekher Shukla 2 Year Classroom	Krishna Sai Shishir 2 Year Classroom	Abhishek Jain 2 Year Classroom	Hardik Aggarwal 2 Year Classroom	Ujjwal Singh 2 Year Classroom	Rishabh Aggarwal 2 Year Classroom	Sanyu Jais 2 Year Classroom	M Sai Divya Tuja Reddy 2 Year Classroom	Rishi Shekher Shukla 2 Year Classroom

Avg point of L_1 be $A(2\lambda + 2, -3\lambda - 5, 4\lambda + 11)$

$R(1, 7, 6)$

$$RA \cdot (2\hat{i} - 3\hat{j} + 4\hat{k}) = 0$$

$$((2\lambda + 1)\hat{i} + (-3\lambda - 12)\hat{j} + (4\lambda + 5)\hat{k}) \cdot (2\hat{i} - 3\hat{j} + 4\hat{k}) = 0$$

$$4\lambda + 2 + 9\lambda + 36 + 16\lambda + 20 = 0$$

$$29\lambda = -58$$

$$\lambda = -2$$

\therefore Foot of perpendicular = $(-2, 1, 3)Q$

$$PQ = \sqrt{(10+2)^2 + (1+2)^2 + (3+1)^2}$$

$$= \sqrt{144 + 9 + 16}$$

$$= 13$$



Delivering Champions Consistently

JEE (Advanced) 2024

AIR 25 Rishi Shekher Shukla 2 Year Classroom	AIR 67 Krishna Sai Shishir 2 Year Classroom	AIR 78 Abhishek Jain 2 Year Classroom	AIR 93 Hardik Aggarwal 2 Year Classroom	AIR 95 Ujjwal Singh 2 Year Classroom	AIR 98 Rachit Aggarwal 2 Year Classroom
---	--	--	--	---	--

JEE (Main) 2024

1st AIR Sarvi Jais 2 Year Classroom	AIR 15 M Sai Divya Teja Reddy 2 Year Classroom	AIR 19 Rishi Shekher Shukla 2 Year Classroom
--	---	---