

**JEE-Main-28-01-2025 (Memory Based)**  
**[MORNING SHIFT]**  
**Maths**

**Question:** If  $\int_{-\pi/2}^{\frac{\pi}{2}} \frac{96(x^2 \cdot \cos x)}{1+e^x} dx = a\pi^3 + \beta$  (where  $\alpha + \beta$  are positive integers), then  $\alpha + \beta$  equal to

**Options:**

- (a) 144
- (b) 100
- (c) 64
- (d) 196

**Answer: (b)**

$$I = 96 \int_{-\pi/2}^{\pi/2} \frac{x^2 + \cos x}{1 + e^x} dx$$

$$I = 96 \int_{-\pi/2}^{\pi/2} \frac{x^2 + \cos x}{1 + e^{-x}} dx$$

$$2I = 96 \int_{-\pi/2}^{\pi/2} (x^2 + \cos x) dx$$

$$\begin{aligned} I &= 96 \left( \frac{x^3}{3} + \sin x \right) \Big|_0^{\pi/2} = 96 \left( \frac{\pi^3}{24} + 1 \right) \\ &= 4\pi^3 + 96 \\ \alpha + \beta &= 100 \end{aligned}$$

**Question:** Number of ways to form 5 digit numbers greater than 50000 with the use of digits 0, 1, 2, 3, 4, 5, 6, 7 such that sum of first and last digit is not more than 8, equal to

**Options:**

- (a) 5119
- (b) 5120
- (c) 4607
- (d) 4068

**Answer: (c)**

Available digit 0, 1, 2, 3, 4, 5, 6, 7

Middle three spaces can be filled in

1st place 5, then last place can be 0, 1, 2, 3 as sum is less than or equal to 8

So total no.'s =  $1 \times 512 \times 4 = 2048$

1st place 6, then last place can be 0, 1, 2

So total no. of ways =  $1 \times 512 \times 3 = 1536$

1st place 7, then last place can be 0, 1

So total no. of ways =  $1 \times 512 \times 2 = 1024$

So total no. of ways =  $2048 + 1536 + 1024 = 4608$

Excluding 50000,  $4608 - 1 = 4607$

**Question:** If  $f(x) = \frac{2^x}{2^x + \sqrt{2}}, x \in R$ , then  $\sum_{k=1}^{81} f\left(\frac{k}{82}\right)$  is equal to

**Options:**

- (a)  $81\sqrt{2}$
- (b) 82
- (c)  $\frac{81}{2}$
- (d) 42

**Answer: (c)**

$$f(x) = \frac{2^x}{2^x + \sqrt{2}} \Rightarrow f(x) + f(1-x) = 1$$

$$s = \sum_{n=1}^{81} f\left(\frac{k}{82}\right)$$

$$s = \sum_{n=1}^{81} f\left(\frac{82-k}{82}\right) = \sum_{n=1}^{81} f\left(1 - \frac{k}{82}\right)$$


$$2s = \sum_{n=1}^{81} f\left(\frac{k}{82}\right) + f\left(1 - \frac{k}{82}\right)$$

$$= \sum_{n=1}^{81} 1 = 81$$

$$s = \frac{81}{2}$$

**Question:**  $z_1 = \sqrt{3} + 2\sqrt{2}i$  &  $\sqrt{3}|Z_1| = |Z_2|$  and  $\arg(z_2) = \arg(z_1) + \frac{\pi}{6}$  then area, of triangle with vertices  $z_1, z_2$  and origin

**Solution:**



$$\begin{aligned}
 \text{Area} &= \frac{1}{2} \sqrt{11} \cdot \sqrt{33} \sin \frac{\pi}{6} \\
 &= \frac{1}{2} \cdot 11\sqrt{3} \cdot \frac{1}{2} \\
 &= \frac{11\sqrt{3}}{4}
 \end{aligned}$$

**Question:** The relation  $R = \{(x,y) | x, y \in z, x + y = \text{even}\}$  then  $R$  is

**Options:**

- (a) Equivalence
- (b) Reflexive & Transitive but - not Symmetric
- (c) Symmetric & Transitive but not reflexive
- (d) Reflexive & symmetric but not transitive

**Answer: (a)**

$$R = \{(x, y) : x + y = \text{even}, x, y \in z\}$$

Reflexive as  $x + x = \text{even}$

Symmetric as  $x + y = y + x$

Transitive as  $x + y = \text{even}$   $y + z = \text{even}$

$x + 2y + z = \text{even}$

$x + z = \text{even}$

*Equivalence.*

**Question:**  $\cos\left(\sin^{-1} \frac{3}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{33}{65}\right)$  is equal to

**Options:**

- (a) 0
- (b) 1
- (c)  $\frac{32}{65}$
- (d)  $\frac{33}{65}$

**Answer: (a)**

$$\cos\left(\sin^{-1} \frac{3}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{33}{65}\right)$$

$$\cos\left(\tan^{-1} \frac{3}{4} + \tan^{-1} \frac{5}{12} + \sin^{-1} \frac{33}{65}\right)$$

$$\cos\left(\tan^{-1} \frac{\frac{3}{4} + \frac{5}{12}}{1 - \frac{3}{4} \times \frac{5}{12}} + \sin^{-1} \frac{33}{65}\right)$$

$$\cos\left(\tan^{-1} \frac{14 \times 4}{33} + \cot^{-1} \frac{56}{33}\right)$$

$$\cos \frac{\pi}{2} = 0$$

**Question:**  $\int_0^x tf(t)dt = x^2 f(x)f(2)3, f(6) = ?$

**Solution :**

$$\int_0^x tf(x)dt = x^2 f(x)$$

$$xf(x) = 2xf(x) + x^2f'(x)$$

$$\Rightarrow xf'(x) = -f(x)$$

$$\Rightarrow \frac{f'(x)}{f(x)} = \frac{-1}{x}$$

$$\Rightarrow \ln f(x) + \ln x = \ln c$$

$$\Rightarrow xf(x) = c$$

$$2f(2) = 6f(6)$$

$$\Rightarrow f(6) = 1$$

**Question:** Area of region  $\{(x, y) : 0 \leq y \leq 2|x| + 1, 0 \leq y \leq x^2 + 1, |x| \leq 3\}$

**Options:**


(a)  $\frac{17}{3}$

(b)  $\frac{32}{3}$

(c)  $\frac{64}{3}$

(d)  $\frac{80}{3}$

**Answer: (c)**




$$\begin{aligned}
 A &= 2 \left[ \int_0^2 (x^2 + 1) dx + \int_2^3 (2x + 1) dx \right] \\
 &= 2 \left[ \left( \frac{x^3}{3} + x \right)_0^2 + (x^2 + x)_2^3 \right] \\
 &= 2 \left( \frac{8}{3} + 2 + 9 + 3 - 4 - 2 \right) = 2 \left( 8 + \frac{8}{3} \right) \\
 &= 2 \times \frac{32}{3} = \frac{64}{3}
 \end{aligned}$$

**Question:**  $x^2 - \alpha x + \beta y - \gamma + y^2 = 0$  if touches x axis at a point A(a, 0) & cut intercept of b v.

find value of a and b on y axis

**Solution :**



$$r = \sqrt{\frac{\alpha^2}{4} + \frac{\beta^2}{4} - \gamma} = \frac{\beta}{2}$$

$$\frac{\alpha^2}{4} - \gamma = 0$$

$$\alpha^2 = 4\gamma, \frac{\alpha}{2} = a$$

$$\alpha = 2a$$

$$2\sqrt{\frac{13^2}{4} - \gamma} = b$$

$$\frac{\beta^2}{4} = \gamma = \frac{b^2}{4}$$

$$b^2 = \beta^2 - 4\gamma$$

$$(2a, b^2) = (\alpha, \beta^2 - 4\gamma)$$

**Question:**  $x^2 + |2x - 3| - 4 = 0$  has sum of squares of roots

**Solution :**

$$x^2 + |2x - 3| - 4 = 0$$

$$\text{If } x \geq \frac{3}{2}, x^2 + 2x - 7 = 0$$

$$x = \frac{-2 \pm \sqrt{4+28}}{2} = \frac{-24\sqrt{2}}{2}$$

$$= -1 \pm 2\sqrt{2}$$

$$x = 2\sqrt{2} - 1$$

$$\text{If } x < \frac{3}{2}, x^2 - 2x - 1 = 0$$

$$x = \frac{2 \pm 2\sqrt{2}}{2} = 1 + \sqrt{2}$$

$$x = 1 - \sqrt{2}$$

$$\text{sum of squares of roots} = (2\sqrt{2} - 1)^2 + (1 - \sqrt{2})^2$$

$$= 12 - 6\sqrt{2}$$

$$= \frac{x-1}{2} = \frac{y-2}{1} = \frac{z-1}{3}$$

**Question:** The image of point  $(4, 4, 3)$  in the plane  $\alpha + \beta + \gamma$ .

**Solution :**

Given line is  $x = 1 + 2t, y = 2 + t, z = 1 + 3t$

Vector joining  $(4, 4, 3)$  to the general point on line

$$(1 + 2t, 2 + t, 1 + 3t) \text{ is } \vec{v} (-3 + 2t, -2 + t, -2 + 3t)$$

Orthogonal projection  $\vec{v} \cdot \vec{d} = 0$   $d$  is direction

$$\text{Vector of the line as } \vec{d} = 2i + j + 3k$$

$$\text{So } (-3 + 2t)2 + (-2 + t) \cdot 1 + (-2 + 3t) \cdot 3 = 0$$

$$-14 + 14t = 0 \Rightarrow t = 1$$

Point of projection is  $(3, 3, 4)$

Vector joining the projection point  $(3, 3, 4)$  and  $(4, 4, 3)$  -  $(1, 1, 1)$

$$\text{So image point} = (3, 3, 4) + (1, 1, -1) = (2, 2, 5)$$

$$\text{Sum of coordinates} = 2 + 2 + 5 = 9$$

**Question:**  $a_0 = 1, a_1 = \frac{1}{2}, 2a_n = 5a_{n-1} - 3a_{n-2}$  find  $\sum_{i=1}^{100} a_i$ .

**Options:**

(a)  $3a_{100} + 197$

- (b)  $2a_{100} - 1000$   
 (c)  $3a_{99} - 100$   
 (d)  $2a_{99} - 100$

**Answer: (a)**

$$a_0 = 1, a_1 = \frac{1}{2}, 2a_n - 3a_{n-1} = 2a_{n-1} - 3a_{n-2}$$

$$\Rightarrow 2a_n - 3a_{n-1} = 2a_1 - 3a_0$$

$$= 1 - 3 = -2$$

$$2 \sum_{n=1}^{100} a_n - 3 \sum_{n=1}^{100} a_{n-1} = -200$$


$$2s = 3[a_0 + s - a_{100}] = -200$$

$$2s = 3s - 3 + 3a_{100} = -200$$

$$s = 3a_{100} + 197$$

**Question:** Vertices of a trapezium lies on a parabola  $y^2 = 4x$  one of its diagonal passes through  $(1, 0)$ , and its length is  $\frac{25}{4}$ . Parallel sides are parallel to y. Find its area of trapezium.

**Solution :**



$$t_1 t_2 = -1$$

$$\sqrt{(t_1^2 + t_2)^2 + (2t_1 - 2t_2)^2} = \frac{25}{4}$$

$$|t_1 - t_2| \sqrt{(t_1 + t_2)^2 - 4t_1 t_2} = \frac{25}{4}$$

$$(t_1 - t_2)^2 = \frac{25}{4}$$

$$t_1 - t_2 = \frac{5}{2}$$

$$r_1 + \frac{1}{t_1} = \frac{5}{2}$$

$$t_1 = 2, t_2 = -\frac{1}{2}$$

$$A = \frac{1}{2}(8 + 2) \times \left(4 - \frac{1}{4}\right) = \frac{1}{2} \times 10 \times \frac{15}{4} = \frac{75}{4}$$

**Question:** A bag has 7 good and 3 defected oranges. 2 oranges are chosen randomly. Find the variance of oranges being defected.

**Solution:**

7 Good, 3 Defective


x	f
0	${}^7C_2 / {}^{10}C_2 = \frac{7}{15}$
1	${}^3C_1 \cdot {}^7C_1 / {}^{10}C_2 = \frac{7}{15}$
2	${}^3C_2 / {}^{10}C_2 = \frac{1}{15}$
$\bar{x}$	$0 + \frac{7}{15} + \frac{2}{15} = \frac{9}{15} = \frac{3}{5}$

$$\begin{aligned} Var &= (0 + \frac{7}{15} + \frac{4}{15}) - (\frac{3}{5})^2 = \frac{11}{15} - \frac{9}{25} \\ &= \frac{110-54}{150} = \frac{56}{150} = \frac{28}{75} \end{aligned}$$

**Question:** The sum of all local minimum values of the function

$$\begin{cases} 1 - 2x & x < -1 \\ \frac{1}{3}(7 + 2|x|) & -1 \leq x \leq 2 \text{ is} \\ \frac{11}{18}(x-4)(x-5) & x > 2 \end{cases}$$

**Solution :**



$$\text{Sum of local minimum values} = \frac{7}{3} - \frac{11}{72}$$

$$= \frac{168-11}{72} = \frac{157}{72}$$