

JEE-Main-28-01-2025 (Memory Based)
[EVENING SHIFT]
Maths

Question: Evaluate $\sum_{r=1}^{13} \frac{1}{\sin\left[\frac{\pi}{4}+(r-1)\frac{\pi}{6}\right]\sin\left[\frac{\pi}{4}+\frac{r\pi}{6}\right]}$

Options:

- (a) $2\sqrt{3} + 2$
- (b) $2\sqrt{3} - 2$
- (c) $3\sqrt{2} + 2$
- (d) $3\sqrt{2} - 4$

Answer: (b)

$$\sum_{r=1}^{13} \frac{1}{\sin\left[\frac{\pi}{4}+(r-1)\frac{\pi}{6}\right]\sin\left[\frac{\pi}{4}+\frac{r\pi}{6}\right]}$$

$$\sum_{r=1}^{13} \frac{\sin\left[\frac{\pi}{4}+\frac{r\pi}{6}\right]-\left[\frac{\pi}{4}+(r-1)\frac{\pi}{6}\right]}{\left(\sin\frac{\pi}{6}\right)\times\sin\left(\frac{\pi}{4}+(r-1)\frac{\pi}{6}\right)\sin\left(\frac{\pi}{4}+\frac{r\pi}{6}\right)}$$

$$\sum_{r=1}^{13} 2\left(\cot\left(\frac{\pi}{4}+(r-1)\frac{\pi}{6}\right)-\cot\left(\frac{\pi}{4}+\frac{r\pi}{6}\right)\right)$$

$$2\left[\cot\frac{\pi}{4}-\cot\left(\frac{\pi}{4}+\frac{13\pi}{6}\right)\right]$$

$$2\left[1-\cot\left(\frac{29\pi}{12}\right)\right]$$

$$2\left[1-\cot\left(2\pi+\frac{5\pi}{12}\right)\right]$$

$$2\left(1-\cot\frac{5\pi}{12}\right)$$

$$2\left(1-2+\sqrt{3}\right)\Rightarrow 2\left(\sqrt{3}-2\right)$$

Question: $f\left(\frac{dx}{x^{\frac{1}{4}}\left(x^{\frac{1}{4}}+1\right)}\right)$, $f(0) = -6$, $f(2) = ?$

Solution :

$$I = \int \frac{dx}{x^{1/4}(x^{1/4}+1)}, f(0) = -6, f(2)$$

$$I = \int \frac{4t^3 dt}{t(t+1)}$$

$$= 4 \int \frac{t^2}{t+1} dt$$

$$= 4 \int \frac{t^2 - 1 + 1}{t+1}$$

$$= 4 \int (t-1) + 4 \int \frac{1}{t+1}$$

$$\Rightarrow 4\left(\frac{t^2}{2} - t\right) + 4 \ln(t+1) + c$$

$$f(x) = 2\left(x^{\frac{1}{2}}\right) - 4x^{\frac{1}{4}} + 4 \ln\left(1+x^{\frac{1}{4}}\right) + c$$

$$C = -6$$

$$f(2) = 2\left(\sqrt{2}\right) - 4\sqrt[4]{2} + 4 \ln\left(1+2^{\frac{1}{4}}\right)$$

Question: Bags B_1, B_2, B_3 contains 4 Blue, 6 white balls, 5 white 5 blue balls and 6 Blue 4 white balls respectively. A bag is randomly selected and a ball is drawn. If the drawn ball is white then find the probability that B_2 bag was selected.

Solution :

$$B_1, B_2, B_3$$

$$B_1 = +4 Bl \ \& \ 6 w$$

$$B_2 = -5 wl \ \& \ 5 Bl$$

$$B_3 = 5 Bl \ \& \ 4 Wl$$

$$P(B_2/A) \Rightarrow \frac{1/3 \times 5/10}{1/3 \times 5/10 + 1/3 \times 6/10 + 1/3 \times 4/10}$$

$$\Rightarrow \frac{\frac{5}{3 \times 10}}{\frac{15}{3 \times 10}} \Rightarrow \frac{1}{3}$$

Question: Find domain of $\sec^{-1}(2[x] + 1)$, where $[.]$ denotes GIF.

Solution :

$$\text{Domain of } \sec^{-1}(2[x] + 1)$$

$$2[x] + 1 \geq 1 \text{ on } 2[x] + 1 \leq -1$$

$$2[x] \geq 0 \quad [x] \leq \frac{-2}{2}$$

$$x \geq 0 \quad [x] \leq -1$$

$$[x] \leq -1$$

$$\therefore x \in (-\infty, -1] \cup [0, \infty)$$

Question: If $x^2 - (3 - 2i)x - (2i - 2) = 0$ has roots $\alpha + i\beta$ and $\gamma + i\delta$ find the value of $\alpha\gamma + \beta\delta$.

Solution :

$$x^2 - (3 - 2i)x - (2i - 2) = 0$$

$$x = \frac{3 - 2i \pm \sqrt{9 - 4 - 12i + 4(2i - 2)}}{2}$$

$$x = \frac{(3 - 2i) \pm \sqrt{-4i - 3}}{2}$$

$$= \left[\left(\frac{1}{\sqrt{2}} \right)^{10} - 2^{10} \right] = \sqrt{-4i - 3} = \alpha - i\beta$$

$$-4i - 3 = \alpha^2 - \beta^2 + i2\alpha\beta$$

$$x^2 - y^2 = -3 \quad 2 \times y = 4$$

$$x^2 - y^2 = -3 \quad \alpha y = -2$$

$$x^2 + y^2 = \sqrt{\alpha^2 - y^2 + 4x^2y^2} = 5$$

$$x^2 + y^2 = 5$$

$$x^2 - y^2 = -3$$

$$2x^2 = 2 \rightarrow x^2 = 1 \Rightarrow \alpha = 1, \beta = -2$$

$$\alpha = -1, \beta = +2$$

$$n = \frac{(3 - 2i) \pm (1 - 2i)}{2} - 2i + 2$$

$$\alpha = 1, \beta = 0, \gamma = 2, \delta = -2$$

$$\alpha\gamma + \beta\delta = 2$$

Question: $A = \begin{bmatrix} \frac{1}{\sqrt{2}} & -2 \\ 0 & 1 \end{bmatrix} P = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} B = PAP^T ; X = PB^{10}P^T$ Find $X = ?$

Solution :

$$A = \begin{bmatrix} \frac{1}{\sqrt{2}} & -2 \\ 0 & 1 \end{bmatrix} P = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$PP^T = I \quad B = PAP^T \quad A^2 = \begin{bmatrix} \frac{1}{\sqrt{2}} & -2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & -2 \\ 0 & 1 \end{bmatrix}$$

$$B^2 = PAP^T(PAP^T) \quad A^2 = \left[\left(\frac{1}{2}\right)^2 I^2\right]$$

$$= PA(P^T P)AP^T$$

$$B^2 = PA^2 P^T \quad X = P^T(PA^{10} P^T)P$$

$$B^{10} = PA^{10} P^T \quad = IA^{10} I = A^{10}$$

$$x = P^T B^{10} P \quad \therefore x = A^{10} = \begin{bmatrix} \left(\frac{1}{\sqrt{2}}\right)^{10} & -2^{10} \\ 0 & 1^{10} \end{bmatrix}$$

Question: 212, 213,.....,999 find no. of numbers in the sequence above whose sum of digits is 15.

Solution :

212, 213,, 999

$\therefore \text{Sum} = 15$

$$x + y + 2 = 15$$

$$x \in \{2, 3, \dots, 9\}$$

$$y \in \{0, 1, 2, \dots, 9\}$$

$$\text{Coefficient } x^{15} : (x^2 + x^3 + \dots + x^9)(x^{11} + x^1 + \dots x^9)^2$$

$$(1 + x + x^2 + \dots + x^2) \left(\frac{1-x^{10}}{1-x}\right)^2$$

$$\left(\frac{1-x^8}{1-x}\right) \left(\frac{1-x^{10}}{1-x}\right)^2$$

$$(1 - x^8)(1 - x^{10})(1 - x)^{-3}$$

$$(1 - x^8)(1 - 2x^{10} + x^{20})(1 - x)^{-3}$$

$$(1 - x^8 - 2x^{10})(1 - x)^{-3}$$

$$\text{Coefficient } x^{12} = {}^{3+13-1}C_{13} - {}^{3+5-1}C_5 - 2^{3+3-1}C_3$$

$$= 64$$

Question: Words are formed using all letters of Word GARDEN. Find the probability in the word vowels are not together.

Solution :

GARDEN=6!

$$\times G \times R \times D \times N \times$$

$$= {}^5C_2 \times 2! \times 4!$$

$$P(\varepsilon) = \frac{{}^5C_2 \times 2 \times 4!}{6!}$$

$$= \frac{20 \times 4!}{6 \cdot 5 \cdot 4!}$$

$$= \frac{2}{3}$$

Question: Area bounded between the curves $C_1 : x(1+y^2)-1=0$ and $C_2 : y^2-2x=0$ is (in sq. unit)

Options:


(a) $\frac{\pi}{2} - \frac{1}{3}$

(b) $\frac{\pi}{4} - \frac{1}{6}$

(c) $2\left(\frac{\pi}{2} - \frac{1}{6}\right)$

(d) $\frac{\pi}{6} + \frac{1}{2}$

Answer: (a)



$$C_x = x(1 + r^2) - 1 \text{ and}$$

$$C_y = y^2 - 2x = 0$$

$$\int_{-1}^1 \left(\frac{1}{1+y^2} - \frac{y^2}{2} \right) dy$$


$$\left(\tan^{-1}y - \frac{y^3}{6} \right)_{-1}^1$$

$$\left(\frac{\pi}{4} - \frac{1}{6} \right) - \left(-\frac{\pi}{4} + \frac{1}{6} \right)$$

$$\frac{\pi}{2} - \frac{1}{3}$$

Question: If $f(x)$ is polynomial satisfying $f(x) \cdot f\left(\frac{1}{x}\right) = f(x) + f\left(\frac{1}{x}\right)$ and $f(2) = 129$; then find real values of “k” satisfying $f(k) = -2k$.

Solution :



$$f(x) \cdot f\left(\frac{1}{x}\right) = f(x) + f\left(\frac{1}{x}\right)$$

$$f(x) = \pm x^4 + 1$$

$$f(x) = \pm 2^4 + 1 = 129$$

$$x = 7$$

$$f(x) = x^7 + 1$$


$$f(x) = k^7 + 1 = -2k$$

$$k^7 + 2k + 1 = 0$$

$$f'(k) = 7k^6 + 2$$

Question: An isosceles triangle formed by 3 lines. Equation of equal sides of triangle are $-x + 2y = 4$ & $x + 3y = 6$. Find sum of possible values of slope of 3rd side.

Solution :



$$\tan \theta = \left| \frac{m - \frac{1}{2}}{1 + \frac{m}{2}} \right| = \left| \frac{\frac{-1}{2} - m}{1 + \frac{m}{3}} \right|$$

$$\frac{2m-1}{m+2} = \pm \left(\frac{-1-3m}{3-m} \right)$$

$$\frac{2m-1}{m+2} = \frac{-(1+3m)}{3-m}$$

$$\frac{2m-1}{m+2} = \frac{3m+1}{3-m}$$