

PHYSICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer:

- An ideal gas with an adiabatic exponent 1.5, initially at 27°C is compressed adiabatically from 800 cc to 200 cc. The final temperature of the gas is:
 - 600 K
 - 300 K
 - 450 K
 - 273 K

Answer (1)

Sol. $T_i = 27^\circ\text{C}$ or 300 K

$$T_f = (300\text{ K}) \left(\frac{800\text{ cc}}{200\text{ cc}} \right)^{1.5-1}$$

$$T_f V_f^{\gamma-1} = T_i V_i^{\gamma-1}$$

$$T_f = T_i \left(\frac{V_i}{V_f} \right)^{\gamma-1}$$

$$T_f = 300\text{ K} (4)^{0.5}$$

$$T_f = 600\text{ K}$$

- In YDSE, light of intensity of $4I$ and $9I$ passes through two slits respectively. Difference of maximum and minimum intensity of interference pattern is
 - $5I$
 - $10I$
 - $24I$
 - $26I$


Answer (3)

Sol. $I_{\max} = (\sqrt{4I} + \sqrt{9I})^2 = 25I$

$$I_{\min} = (\sqrt{4I} - \sqrt{9I})^2 = I$$

$$\Delta I = 24I$$

- An ammeter having resistance $240\ \Omega$ is connected in the given circuit as shown. Find current through the ammeter.



- 1 mA
- 5 mA
- 100 mA
- 2.5 mA

Answer (2)

Sol. $R_{\text{eq}} = 150.4 + \frac{10 \times 240}{250}$

$$= 160\ \Omega$$

$$I = \frac{20}{160} = \frac{1}{8}$$

$$I_A = \frac{10}{250} I = \frac{1}{25 \times 8} = \frac{1}{200}$$

$$I_A = 5\text{ mA}$$

- A thin uniform wire of length 25 m and area of cross-section 5mm^2 has resistivity $2 \times 10^{-8}\ \Omega\text{-m}$. If the wire is bent to form a circle, the resistance across diametrically opposite points is
 - 5 Ω
 - 2.5 Ω
 - 10 Ω
 - 12.5 Ω

Answer (2)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 stars

70+

1000+

4000+

OUR JEE CHAMPIONS



100% Rank



100% Rank



100% Rank



100% Rank




JEE Main
2020

JEE Main
2022

JEE Main
2024



Sol. $R = \rho \frac{l}{A} = \frac{(2 \times 10^{-6} \Omega \text{ m})(25 \text{ m})}{5 \times 10^{-6} \text{ m}^2}$



$R = 10 \Omega$

Resistance across diametrically point is $\frac{R}{4} = 2.5 \Omega$

5. Capacitors with dielectric are shown in figure (symmetric situation).



Find $\frac{C_1}{C_2}$

(1) $\frac{4\epsilon_1\epsilon_2}{(\epsilon_1 + \epsilon_2)^2}$ (2) $\frac{4\epsilon_1\epsilon_2}{\epsilon_1 + \epsilon_2}$

(3) $\frac{2\epsilon_1\epsilon_2}{(\epsilon_1 + \epsilon_2)^2}$ (4) $\frac{(\epsilon_1\epsilon_2)^2}{(\epsilon_1 + \epsilon_2)^2}$

Answer (1)

Sol. $\frac{1}{C_1} = \frac{d}{2A\epsilon_1} + \frac{d}{2A\epsilon_2}$


$\Rightarrow C_1 = \frac{2A}{d} \frac{\epsilon_1\epsilon_2}{(\epsilon_1 + \epsilon_2)^2}$

$C_2 = \frac{A\epsilon_1}{2d} + \frac{A\epsilon_2}{2d}$

$C_2 = \frac{A}{2d} (\epsilon_1 + \epsilon_2)$

$\frac{C_1}{C_2} = \frac{4\epsilon_1\epsilon_2}{(\epsilon_1 + \epsilon_2)^2}$

6. A sphere of mass 20 kg is pulled with force of 49 N as shown in diagram. Acceleration of sphere assuming no slipping.



(1) 2.5 m/s²

(2) 2.8 m/s²

(3) 1.4 m/s²

(4) 3.5 m/s²


Answer (4)

Sol. $\tau = F2R = \frac{7}{5} MR^2 \alpha = \frac{7}{5} MRa$

$2F = \frac{7}{5} Ma$

$a = \frac{10F}{7M} = \frac{10 \times 49}{7 \times 20} = 3.5 \text{ m/s}^2$

7. A current carrying wire is bent as shown in the figure. Find magnetic field at centre 'O' of the semi-circles. (take $R_1 = 4\pi$ and $R_2 = 6\pi$)



(1) $8.3I \times 10^{-6} \text{ T}$

(2) $8.3I \times 10^{-8} \text{ T}$

(3) $4\pi I \times 10^{-7} \text{ T}$

(4) $6I \times 10^{-8} \text{ T}$

Answer (2)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-I) 2025

4 **100+** AIR

70+ **100+** AIR

1000+ **100+** AIR

4000+ **100+** AIR

100
100



100
100



100
100



100
100



100
100

100
100

100
100

Sol. $B = \frac{\mu_0 I}{4R_1} - \frac{\mu_0 I}{4R_2}$
 $= \frac{4\pi \times 10^{-7}}{4} I \left(\frac{1}{4\pi} - \frac{1}{6\pi} \right)$
 $= \frac{1}{12} \times 10^{-7} I$
 $= 8.3 I \times 10^{-9} \text{ T}$


8. A biconvex lens is having the radius of curvature of 10 cm and 15 cm. If focal length of the lens is 12 cm find refractive index of material of the lens.

- (1) $\frac{3}{2}$ (2) $\frac{4}{3}$
 (3) 2 (4) $\sqrt{3}$

Answer (1)

Sol. $\frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$
 $\frac{1}{12} = (\mu - 1) \left(\frac{1}{10} + \frac{1}{15} \right)$
 $\mu - 1 = \frac{1}{2}$
 $\mu = \frac{3}{2}$

9. The figure below shows an oscillating system of two blocks and a spring. The horizontal surface is smooth and the contact between the blocks is rough with coefficient of static friction μ .



Considering that the blocks of mass m is always stationary relative to M , choose the correct option regarding the statements below:

- (A) Maximum frictional force between blocks is μmg .
 (B) Time period of oscillation is $2\pi \sqrt{\frac{m+M}{k}}$
 (C) Friction between the blocks at any instant is $\mu(m+M)g$

- (1) Only A is correct
 (2) Only B is correct
 (3) A, B and C all three are correct
 (4) Only C is correct

Answer (2)

Sol. $f \leq \mu N$

$\Rightarrow f \leq \mu mg$

$T = 2\pi \sqrt{\frac{m+M}{k}}$ for no slipping between the blocks.



10. A point source of power 450 W is emitting light in all direction. Radiation pressure at distance of 2m from the source is nearly

- (1) $3 \times 10^{-8} \text{ Pa}$ (2) $2 \times 10^{-8} \text{ Pa}$
 (3) $2 \times 10^{-9} \text{ Pa}$ (4) $4 \times 10^{-7} \text{ Pa}$

Answer (1)

Sol. $P = \frac{I}{C} = \frac{P}{4\pi r^2} = \frac{450}{4 \times 3.14 \times 2^2 \times 3 \times 10^8} = \frac{450}{48 \times 3.14} \times 10^{-8}$
 $\approx 3 \times 10^{-8} \text{ Pa}$

11. From a horizontal surface a particle is projected with a speed u . Which of the following graph correctly represent the variation of maximum height above the surface attained by the particle as the angle of projection is varied?



THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4
times attempt

70+
top scorers

1000+
rankings

4000+
candidates

100
times



Shreyas Lohiya
AIR 100 (2025)

100
times



Harsh Jha
AIR 100 (2025)

100
times



Devya Rautel
AIR 100 (2025)

100
times



Anshul Bansal
AIR 100 (2025)


OUR JEE CHAMPIONS





Shreyas Lohiya
AIR 100 (2025)



Devya Rautel
AIR 100 (2025)



Anshul Bansal
AIR 100 (2025)



Answer (1)

Sol. $h_{\max} = \frac{u^2 \sin^2 \theta}{2g}$

12. An object is dropped from height S . At a point its kinetic energy is three times its potential energy. Find its height from ground and speed at that point.

- (1) $\frac{3S}{4} \cdot \sqrt{\frac{3gS}{2}}$ (2) $\frac{S}{4} \cdot \sqrt{\frac{3gS}{2}}$
 (3) $\frac{S}{2} \cdot \sqrt{gS}$ (4) $\frac{S}{4} \cdot \sqrt{\frac{3gS}{4}}$

Answer (2)

Sol.

KE + PE = mgS
 3PE + PE = mgS
 4mgh = mgS
 $h = \frac{S}{4}$
 $v = \sqrt{2g \cdot \left(\frac{3S}{4}\right)}$
 $v = \sqrt{\frac{3gS}{2}}$

13. The electric potential at the surface of a shell of radius 10 cm is 120 V. Find the potential at its centre, at $r = 5$ cm from centre and at $r = 15$ cm from centre.

- (1) 0 V, 0 V, 80 V (2) 120 V, 120 V, 80 V
 (3) 120 V, 0 V, 80 V (4) 80 V, 0 V, 120 V


Answer (2)

Sol. $V_{\text{inside}} = V_{\text{surface}}$
 $\Rightarrow V_{\text{centre}} = V(r = 5 \text{ cm}) = 120 \text{ V}$
 $V_{\text{outside}} = \frac{(V_{\text{surface}})(R)}{r}$
 $\Rightarrow V(r = 15 \text{ cm}) = \frac{(120 \text{ V})(10 \text{ cm})}{(15 \text{ cm})}$
 $= 80 \text{ V}$

14. Truth table of logical circuit is given

A	B	Y
0	0	0
1	1	1
0	1	0
1	0	0

Then identify the correct circuit.



Answer (3)

Sol. Truth table is of AND gate

$A \cdot B = \overline{\overline{A} + \overline{B}}$

THE LEGACY OF SUCCESS CONTINUES

OUR JEE CHAMPIONS

JEE Main (C-Session-1) 2025

4 100% TOPPERS

70+ 100% AIR RANKS

1000+ 100% ADMITTANCE

4000+ 100% PLACEMENTS

Shreyas Lodha
100% AIR RANK

Harsh Jha
100% AIR RANK

Devya Rautagi
99.99 AIR RANK

Anagha Barwal
99.99 AIR RANK

Shreyas Lodha
100% AIR RANK
2020

Devya Rautagi
100% AIR RANK
2022

Anagha Barwal
100% AIR RANK
2024

