


PHYSICS

SECTION - A

Multiple Choice Questions: This section contains 20 multiple choice questions. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE** is correct.

Choose the correct answer:

1. Find output voltage in the given circuit.



- (1) +5 Volt (2) 0
(3) 10 Volt (4) -5 Volt

Answer (2)

Sol. D_2 is in forward biased.

$$\text{So, } V_0 = 0$$

2. A fractional errors in x , y and z are 0.1, 0.2 and 0.5 respectively. Find maximum fractional error in $x^{-2}y^{\frac{3}{2}}z^{-\frac{2}{5}}$.

- (1) 0.2
(2) 0.7
(3) 0.6
(4) 0.3

Answer (2)

$$\begin{aligned} \text{Sol. Max. fractional error} &= 2 \times 0.1 + \frac{3}{2} \times 0.2 + \frac{2}{5} \times 0.5 \\ &= 0.2 + 0.3 + 0.2 \\ &= 0.7 \end{aligned}$$

3. A convex lens ($f = 30$ cm) is in contact with concave lens ($f = 20$ cm). Object is placed on the left side at a distance of 20 cm. Find the image distance.

- (1) 10 cm (2) 20 cm
(3) 15 cm (4) 25 cm

Answer (3)

$$\text{Sol. } \frac{1}{F} = \frac{1}{f_1} + \frac{1}{f_2} = \frac{1}{30} + \frac{1}{-20} = \frac{2-3}{60} = \frac{-1}{60} \Rightarrow F = -60 \text{ cm}$$

$$v = \frac{Fu}{F+u} = \frac{(-60)(-20)}{-60-20} = \frac{60 \times 20}{-80} = -15 \text{ cm}$$

4. For a nucleus of mass number A and radius R , mass density is ρ . Then choose the correct option.

- (1) $\rho \propto A^{\frac{1}{3}}$ (2) ρ is independent of A
(3) $\rho \propto A$ (4) $\rho \propto A^3$

Answer (2)

Sol. Conceptual

5. There are two charged sphere of radius R and $3R$. When the sphere are made to touch each other and then separate, the surface charge density becomes r_1 and r_2 respectively. Find $\frac{r_1}{r_2}$

- (1) $\frac{1}{9}$ (2) $\frac{1}{3}$
(3) 3 (4) 9

Answer (3)

Sol. We know finally the potential will be same.

$$\text{So, } \delta_1 r_1 = \delta_2 r_2$$

$$\Rightarrow \delta_1 R = \delta_2 \cdot 3R$$

$$\Rightarrow \frac{\delta_1}{\delta_2} = 3$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS

1000+ 99
PERCENTILERS

4000+ 95
PERCENTILERS

100
Percentile
in Physics
& Maths



Shreyas Lohiya
PSID: 00003389699

100
Percentile
in Physics



Harsh Jha
PSID: 00014863322

100
Percentile
in Physics
& Chemistry



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



Chirag Falor
4 Year Classroom
AIR-16 CRL
JEE (Adv.)
2022



Tanishka Kabra
4 Year Classroom
AIR-16 CRL
JEE (Adv.)
2022



Sanvi Jain
4 Year Classroom
AIR-34 CRL
JEE (Main)
2024

6. Given $\lambda = \frac{2nC}{m}$ (linear charge density) is for Q wire which is passing through body diagonal of a closed cube of side length $\sqrt{3}$ cm. Find flux through the cube.

- (1) 1.44π
 (2) 0.72π
 (3) 2.16π
 (4) 6.84π

Answer (3)

Sol. Total length of the wire inside cube is 3 cm

$$q_{(\text{in})} = 2 \times 10^{-9} \times 3 \times 10^{-2}$$

$$\phi = 2 \times 10^{-9} \times 3 \times 10^{-2} \times 4\pi \times 9 \times 10^9$$

$$\Rightarrow \phi = 2.16\pi$$

7. A monoatomic gas is stored in a thermally insulated container. The gas is suddenly compressed to $\left(\frac{1}{8}\right)^{\text{th}}$ of its initial volume. Find ratio of final pressure to initial pressure.

- (1) 8
 (2) 16
 (3) 4
 (4) 32

Answer (4)

Sol. Process will be adiabatic

$$P_i V_i^{5/3} = P_f \left(\frac{V_i}{8}\right)^{5/3}$$

$$\frac{P_f}{P_i} = 32$$

8. Two balls are projected with same speed at different angles. If maximum height of 1st is 8 times maximum height of 2nd ball. Find the ratio of their time of flight

- (1) $1:2\sqrt{2}$ (2) $2\sqrt{2}:1$
 (3) $2:1$ (4) $4:1$

Answer (2)

$$\text{Sol. } \frac{H_1}{H_2} = 8 = \frac{u^2 \sin^2 \theta_1 / 2g}{u^2 \sin^2 \theta_2 / 2g} = \left(\frac{\sin \theta_1}{\sin \theta_2}\right)^2 \Rightarrow \frac{\sin \theta_1}{\sin \theta_2} = 2\sqrt{2}$$

$$\frac{T_1}{T_2} = \frac{2u \sin \theta_1 / g}{2u \sin \theta_2 / g} = \frac{\sin \theta_1}{\sin \theta_2} = 2\sqrt{2}$$

9. A uniform disc of radius r is rotating about a axis passing through diameter with angular speed 800 rpm. A torque of magnitude 25π Nm is applied on the disc for 40 sec. If final angular speed of disc is 2100 rpm. Find diameter of the disc if mass is 1 kg.

- (1) $\frac{40}{3}$ (2) $40\sqrt{\frac{3}{13}}$
 (3) $20\sqrt{\frac{2}{13}}$ (4) $10\sqrt{\frac{3}{2}}$

Answer (2)

$$\text{Sol. } \omega_f = \omega_i + \alpha t$$

$$\frac{2100 \times 2\pi}{60} = \frac{800 \times 2\pi}{60} + \alpha \times 40$$

$$\alpha = \frac{13}{12} \pi$$

$$\frac{25\pi}{13\pi} = \frac{mR^2}{4}$$

$$\sqrt{\frac{1200}{13}} = 4$$

$$R = 20\sqrt{\frac{3}{13}}, D = 40\sqrt{\frac{3}{13}}$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILERS

1000+ 99
PERCENTILERS

4000+ 95
PERCENTILERS

100
Percentile



Shreyas Lohiya
PSID: 00003389699

100
Percentile



Harsh Jha
PSID: 00014863322

100
Percentile



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



10. Water falls from 200 m height. What is increase in temperature when it touches the bottom. (Assume that all the heat goes into same amount of mass which was falling).

- (1) 0.7°C
(2) $\frac{10}{21}^{\circ}\text{C}$
(3) $\frac{20}{21}^{\circ}\text{C}$
(4) $\frac{11}{10}^{\circ}\text{C}$

Answer (2)

Sol. $mg4h = m \Delta T$

$$\Rightarrow \Delta T = \frac{g\Delta h}{s} = \frac{10 \times 200}{4200}$$

$$\Delta T \approx 05^{\circ}\text{C}$$

$$\Delta T = \frac{10}{21}^{\circ}\text{C}$$

11. An electron is released in the field generated by a non-conductivity sheet of uniform surface charge density σ . The rate of change of de-Broglie wavelength associated with electron varies inversely as n^{th} power of distance travelled. Find the value of n

- (1) $\frac{1}{2}$ (2) 2
(3) $\frac{1}{4}$ (4) 4

Answer (1)

Sol. $E = \frac{\delta}{2\epsilon_0}$

$$F = \frac{e\delta}{2\epsilon_0}$$

$$\frac{1}{2}mv^2 = \frac{e\delta}{2\epsilon_0}x$$

$$\Rightarrow (mv)^1 = \frac{2me\delta}{2\epsilon_0}x = \frac{me\delta}{\epsilon_0}x$$

$$\Rightarrow P = \sqrt{x \frac{me\delta}{\epsilon_0}}$$

$$l = \frac{h}{p} = h \sqrt{\frac{\epsilon_0}{me\delta \cdot x}}$$

$$\lambda \propto \frac{1}{(x)^{\frac{1}{2}}}$$

12. Bulk modulus of a liquid is 2×10^9 Pa initially and final pressure are 1 atm and 5 atm respectively. Find initial volume of the liquid if change in volume is 0.8 cm^3 .

- (1) $2 \times 10^3 \text{ cm}^3$ (2) $4 \times 10^3 \text{ cm}^3$
(3) $2 \times 10^{-4} \text{ cm}^3$ (4) $4 \times 10^{-3} \text{ cm}^3$

Answer (2)

Sol. $\beta = \frac{\Delta P}{\frac{\Delta V}{V}} \Rightarrow V = \frac{\Delta V \beta}{\Delta P}$

$$= \frac{0.8 \times 10^{-6} \times 2 \times 10^9}{4 \times 10^5} = 4 \times 10^{-3} \text{ m}^3 = 4 \times 10^3 \text{ cm}^3$$

13. The amplitude and phase of the wave when two travelling waves given as $y_1(x, t) = 4 \sin(\omega t - kx)$ & $y_2(x, t) = 2 \sin\left(\omega t - kx + \frac{2\pi}{3}\right)$ are superimposed.

- (1) $6, \frac{2\pi}{3}$ (2) $6, \frac{\pi}{3}$
(3) $2\sqrt{3}, \frac{\pi}{6}$ (4) $\sqrt{3}, \frac{\pi}{6}$

Answer (3)

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

**4 STATE
TOPPERS**

**70+ 100
PERCENTILERS**

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS

100
Percentile



Shreyas Lohiya
PSID: 00003389699

100
Percentile



Harsh Jha
PSID: 00014863322

100
Percentile



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



Chirag Falor
4 Year Classroom
1 AIR
JEE (Adv.)
2020



Tanishka Kabra
4 Year Classroom
1 AIR-16 CRL
JEE (Adv.)
2022



Sanvi Jain
4 Year Classroom
1 AIR-34 CRL
JEE (Main)
2024

Sol. $A^2 = A_1^2 + A_2^2 + 2A_1 A_2 \cos \phi$

$$= 4^2 + 2^2 + 2 \times 4 \times 2 \times -\frac{1}{2}$$

$$A = \sqrt{12} = 2\sqrt{3}$$

$$\phi = \frac{2\pi}{3}$$

14. A force $6\hat{k}$ is applied for $\frac{5}{3}$ seconds on a body of mass 2 kg. If initial velocity of body was $3\hat{i} + 4\hat{j}$. Then find final velocity of the body.

- (1) $3\hat{i} + \hat{j} + 5\hat{k}$ (2) $3\hat{i} + 4\hat{j} + 5\hat{k}$
 (3) $3\hat{i} + 2\hat{j} - 3\hat{k}$ (4) $3\hat{i} + 4\hat{j} - 5\hat{k}$

Answer (2)

Sol. Impulse $= 6 \times \frac{5}{3} \hat{k} = 10\hat{k}$

$$\vec{P}_f - \vec{P}_i = 10\hat{k}$$

$$2\vec{v}_f - 2(3\hat{i} + 4\hat{j}) = 10\hat{k} \quad \vec{v}_f = 3\hat{i} + 4\hat{j} + 5\hat{k}$$

15. A rod of linear mass density ' λ ' and length ' L ' is bent into the form of a ring of radius R . Moment of inertia of ring about any of its diameter is


- (1) $\frac{\lambda L^3}{12}$ (2) $\frac{\lambda L^3}{4\pi^2}$
 (3) $\frac{\lambda L^2}{12}$ (4) $\frac{\lambda L^3}{8\pi^2}$

Answer (4)

Sol. $I = \frac{MR^2}{2} = \frac{\lambda L}{2} R^2 \Rightarrow \frac{\lambda L L^2}{2 \times 4\pi^2} \quad \left[\because R = \frac{L}{2\pi} \right]$

$$= \frac{\lambda L^3}{8\pi^2}$$

16. A cube of side 10 cm is suspended from one end of a fine string of length 27 cm, and a mass of 200 gram is connected to the other end of the string. When the cube is half immersed in water, then the system remains in balance. Find density of cube.



- (1) 800 kg/m³
 (2) 500 kg/m³
 (3) 700 kg/m³
 (4) 600 kg/m³

Answer (3)

Sol. $\left(\frac{1}{10}\right)^3 \sigma g - \left(\frac{1}{10}\right)^3 \frac{\rho_0 g}{2} = 0.2 g$

$$\Rightarrow \left(\frac{1}{10}\right)^3 \left(\sigma - \frac{\rho_0}{2}\right) g = 0.2 g$$

$$\Rightarrow \sigma - \frac{\rho_0}{2} = 200$$

$$\Rightarrow \sigma = 200 + 500$$

$$\Rightarrow \sigma = 700 \text{ kg/m}^3$$

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

4 STATE
TOPPERS

70+ 100
PERCENTILES

1000+ 99 PERCENTILES

4000+ 95 PERCENTILES

100
Percentile



Shreyas Lohiya
PSID: 00003389699

100
Percentile



Harsh Jha
PSID: 00014863322

100
Percentile



Devya Rustagi
PSID: 00014768785

99.99
Percentile



Amogh Bansal
PSID: 00014769016

OUR JEE CHAMPIONS



Chirag Falor
4 Year Classroom
1 AIR-2020
JEE (Adv.)




Tanishka Kabra
4 Year Classroom
1 AIR-16 CRL
JEE (Adv.)
2022



Sanvi Jain
4 Year Classroom
1 AIR-34 CRL
JEE (Main)
2024

17. A cone made of conducting material is given a charge Q . $\sigma_1, \sigma_2, \sigma_3$ and σ_4 are charge densities at four points P, Q, R and S . P is at vertex of cone and Q, R and S are at periphery of the base. Choose correct option.



- (1) $\sigma_1 > \sigma_2 > \sigma_3 > \sigma_4$
(2) $\sigma_1 > \sigma_2 = \sigma_3 = \sigma_4$
(3) $\sigma_1 < \sigma_2 = \sigma_3 < \sigma_4$
(4) $\sigma_1 = \sigma_2 > \sigma_3 > \sigma_4$

Answer (2)

Sol. Charge density $\propto \frac{1}{\text{Radius of curvature}}$

18. **Assertion (A)** : Work done to move a charge between two points is zero inside a uniformly charged shell.

Reason (R) : Potential inside a uniformly charged shell is constant and equal to the potential at its surface.

- (1) Both (A) & (R) are correct, and Reason is the correct explanation of (A).
(2) Both (A) & (R) are correct but R is not correct explanation of A.
(3) A is correct and R is incorrect
(4) A is correct and R is correct

Answer (1)

Sol. $W = 0 = q \Delta V$

as $\Delta V = 0$

19.

20.

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

21. Two ropes of same material of radius R and $\frac{R}{2}$. What will

be the ratio of wave speed in second rope to first?

(They both are with same tension)

Answer (2)

Sol. $\mu = \frac{(\pi R^2 l \rho)}{l} = \pi R^2 \rho$

So, $v_2 = \sqrt{\frac{5\tau}{\pi \rho R^2}}$

$v_1 = \sqrt{\frac{\tau}{\pi \rho R^2}}$

$\therefore \frac{v_2}{v_1} = 2$

22.

23.

24.

25.

THE LEGACY OF SUCCESS CONTINUES

JEE Main (Session-1) 2025

**4 STATE
TOPPERS**

**70+ 100
PERCENTILERS**

1000+ 99 PERCENTILERS

4000+ 95 PERCENTILERS



OUR JEE CHAMPIONS

