

$$\int dy + \int \frac{dx}{1+x^2} - \int \frac{\tan^{-1} x}{1+x^2} dx = 0$$

Or $y + \tan^{-1} x - I = 0$, where

$$I = \int \frac{\tan^{-1} x}{1+x^2} dx$$

Let $\tan^{-1} x = t$

$$\frac{1}{1+x^2} dx = dt$$

$$\int t ds \Rightarrow \frac{t^2}{2} = \frac{(\tan^{-1} x)^2}{2}$$

$$\Rightarrow y + \tan^{-1} x - \frac{(\tan^{-1} x)^2}{2} + c = 0$$

$$\therefore y(0) = 1 \quad 1 + c = 0 \Rightarrow c = -1$$

$$\Rightarrow y = \frac{(\tan^{-1} x)^2}{2} - \tan^{-1} x + 1$$

$$\text{Now } y(1) = \frac{1}{2} \left(\frac{\pi}{4} \right)^2 - \frac{\pi}{4} + 1$$

4. The value of

$$\int_0^{\pi/2} |\sin x + \sin 2x + \sin 3x| dx \text{ is}$$

(1) $\frac{8}{3}$

(2) $\frac{7}{3}$

(3) $\frac{2}{3}$

(4) 3

Answer (2)

Sol. $\therefore \sin x + \sin 3x = \sin(2x - x) + \sin(2x + x)$

$$= 2\sin 2x \cos x$$


$$\therefore \sin 3x + \sin 2x + \sin x = \sin 2x (2\cos x + 1)$$

$$\therefore \int_0^{\pi/2} |\sin 2x (2\cos x + 1)| dx$$

$$= \int_0^{\pi/2} (\sin 2x) dx + \int_0^{\pi/2} 4\cos^2 x \sin x dx$$

$$= \frac{-\cos 2x}{2} \Big|_0^{\pi/2} + \left(\frac{-4}{3} (\cos^3 x) \right) \Big|_0^{\pi/2} = \frac{7}{3}$$

5. If L_1 and L_2 are two parallel lines and ΔABC is an equilateral triangle then area of triangle ABC is



(1) $7\sqrt{3}$


(2) $4\sqrt{3}$

(3) $21\sqrt{3}$

(4) 84

Answer (3)

Sol. Let a be the side of ΔABC



$$\sin \theta = \frac{3}{a}$$

$$\sin(60^\circ + \theta) = \frac{9}{a}$$

$$\Rightarrow \frac{\sqrt{3}}{2} \times \sqrt{1 - \frac{9}{a^2}} + \frac{3}{a} \times \frac{1}{2} = \frac{9}{a}$$

$$\Rightarrow \sqrt{3}(\sqrt{a^2 - 9}) + 3 = 18$$

$$\Rightarrow 3(a^2 - 9) = 15^2$$

$$\Rightarrow a^2 - 9 = 15 \times 5$$

$$\Rightarrow a^2 = 84$$

$$\text{Area} = \frac{\sqrt{3}}{4} a^2 = 21\sqrt{3}$$

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



6. Consider a set $S = \{a, b, c, d\}$. Then number of reflexive as well as symmetric relations from $S \rightarrow S$ are
 (1) 1024 (2) 256
 (3) 16 (4) 64

Answer (4)

Sol. Number of such relations:

From $A \rightarrow A$ such that

$n(A) = N$ is

$$\Rightarrow 2^{\binom{N^2-N}{2}}$$

for $N = 4 \Rightarrow 2^6 = 64$

Alter :


$$\begin{bmatrix} \Theta & - & - & - \\ \Theta & \Theta & - & - \\ \Theta & \Theta & \Theta & - \\ \Theta & \Theta & \Theta & \Theta \end{bmatrix} \Rightarrow \text{each } \Theta \text{ has two choices}$$

$\Rightarrow 2^6 = 64$

7. The locus of point of intersection of tangent drawn to the circle $(x - 2)^2 + (y - 3)^2 = 16$, which sub-tends an angle of 120° is
 (1) $3x^2 + 3y^2 - 12x - 18y - 25 = 0$
 (2) $x^2 + y^2 - 12x - 18y - 25 = 0$
 (3) $3x^2 + 3y^2 + 12x + 18y - 25 = 0$
 (4) $x^2 + y^2 + 12x + 18y - 25 = 0$

Answer (1)

Sol.



$x^2 + y^2 - 14x - 6y - 3 = 0$

$$\tan 60^\circ = \frac{4}{\sqrt{h^2 + k^2 - 4h - 6k - 3}}$$

squaring both side

$3(h^2 + k^2 - 4h - 6k - 3) = 16$

To get locus replace h, k by x and y ;

$3x^2 + 3y^2 - 12x - 18y - 9 - 16 = 0$

$3x^2 + 3y^2 - 12x - 18y - 25 = 0$

8. If a_1, a_2, a_3, \dots are the terms of an increasing geometric progression such that

$a_1 + a_3 + a_5 = 21,$

$a_1 a_3 a_5 = 64$

then $a_1 + a_2 + a_3$ is

- (1) 5 (2) 7
 (3) 10 (4) 15

Answer (2)

Sol. Let $a_3 = P, a_1 = \frac{P}{r^2}, a_5 = Pr^2$

$\Rightarrow P + \frac{P}{r^2} + Pr^2 = 21$

and $P \times \frac{P}{r^2} \times Pr^2 = 64 \Rightarrow P^3 = 64 = 4^3$

$\Rightarrow P = 4$

$\Rightarrow 1 + \frac{1}{r^2} + r^2 = \frac{21}{4} \Rightarrow r^4 - \frac{17}{4}r^2 + 1 = 0$

$r^2 = \frac{\frac{17}{4} \pm \sqrt{\frac{289}{16} - 4}}{2} = \frac{\frac{17}{4} \pm \frac{15}{4}}{2} = 4, \frac{1}{4}$

$\Rightarrow r = \pm 2, \pm \frac{1}{2}$

$\Rightarrow a_1 + a_2 + a_3 = \frac{P}{r^2} + \frac{P}{r} + P$

$= 4 \left(\frac{1}{4} + \frac{1}{2} + 1 \right)$

$= 1 + 2 + 4 = 7$

9. Ellipse $E: \frac{x^2}{36} + \frac{y^2}{16} = 1$, A hyperbola confocal with ellipse

E and eccentricity of hyperbola is equal to 5. The length of latus rectum of hyperbola is, if principle axis of hyperbola is x-axis?

- (1) $\frac{96}{\sqrt{5}}$ (2) $24\sqrt{5}$
 (3) $18\sqrt{5}$ (4) $12\sqrt{5}$

Answer (1)

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)				JEE (MAIN)			
 ADVAY MAYANK AIR 36	 RUJUL GARG AIR 41	 ARUSH ANAND AIR 64	 SHREYAS LOHIYA AIR 6 Uttar Pradesh Topper 100 Overall	 KUSHAGRA BAINGAHA AIR 7 Uttar Pradesh Topper 100 Overall	 HARSH A GUPTA AIR 15 Telangana Topper 100 Overall		

Sol. $E: \frac{x^2}{A^2} + \frac{y^2}{B^2} = 1$

$H: \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

For con-focal E and H

$\therefore A^2 - B^2 = a^2 + b^2$ (Given that $A^2 = 36$ and $B^2 = 16$)

$36 - 16 = a^2 + b^2$

$a^2 + b^2 = 20 \dots(1)$

$e_H = 5$

$\Rightarrow 25 = 1 + \frac{b^2}{a^2}$

$\Rightarrow 25a^2 = 20 \Rightarrow a^2 = \frac{4}{5} \Rightarrow a = \frac{2}{\sqrt{5}}$

Also $a^2 + b^2 = 20$

$b^2 = 20 - \frac{4}{5} \Rightarrow \frac{96}{5}$

$L(LR) = 2 \times \frac{b^2}{a} = \frac{2 \times \frac{96}{5}}{\frac{2}{\sqrt{5}}} = \frac{96}{\sqrt{5}}$

10. If the mean and variance of observations $x, y, 12, 14, 4, 10, 2$ is 8 and 16 respectively where $x > y$. Then, the value of $3x - y$ is

- (1) 18 (2) 20
(3) 22 (4) 24

Answer (1)

Sol. $\frac{x + y + 12 + 14 + 4 + 10 + 2}{7} = 8$

$\Rightarrow x + y = 14$

also var = 16

$\frac{x^2 + y^2 + 144 + 196 + 16 + 100 + 4}{7} - (8)^2 = 16$

$\Rightarrow x^2 + y^2 = 100$

By solving we get

$x = 8, y = 6$


$\Rightarrow 3x - y = 18$

11. If O is the vertex of the parabola $x^2 = 4y$, Q is the point on parabola. If C is the locus of point which divides OQ in ratio 2:3. The equation of chord of C which is bisected at point $(1, 2)$.

- (1) $5x + 4y + 3 = 0$ (2) $5x - 4y - 3 = 0$
(3) $5x - 4y + 3 = 0$ (4) $5x + 4y - 3 = 0$

Answer (C)

Sol.



$h = \frac{4t}{5}, k = \frac{2t^2}{5}$

$\Rightarrow \frac{2t^2}{5} = t, \frac{2t^2}{5} = t^2$

$\frac{5k}{2} = \left(\frac{5h}{4}\right)^2$

Replace (h, k) with (x, y)

$\frac{5y}{2} = \frac{25x^2}{16}$

$C: 8y = 5x^2$

chord with given middle point

$T = S_1$

$5xx_1 - 4(y + y_1) = 5x_1^2 - 8y_1$

$5x - 4(y + 2) = 5 - 16$

$5x - 4y - 8 = -11$

$5x - 4y + 3 = 0$

12. The value of $\int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \frac{\pi + 4x^{11}}{1 - \sin\left(|x| + \frac{\pi}{6}\right)} dx$

- (1) 3π (2) 4π
(3) 6π (4) 12π

Answer (2)

Our Problem Solvers shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSH
A GUPTA
AIR 15
Telangana Topper
100 Overall



Sol.
$$\int_0^{\frac{\pi}{6}} \frac{\pi + 4x^{11}}{1 - \sin\left(x + \frac{\pi}{6}\right)} + \frac{\pi - 4x^{11}}{1 - \sin\left(x + \frac{\pi}{6}\right)} dx$$

$$= 2\pi \int_0^{\frac{\pi}{6}} \frac{1}{1 - \sin\left(x + \frac{\pi}{6}\right)} dx$$

$$= 2\pi \int_0^{\frac{\pi}{6}} \frac{1 + \sin\left(x + \frac{\pi}{6}\right)}{\cos^2\left(x + \frac{\pi}{6}\right)} dx$$

$$= 2\pi \int_0^{\frac{\pi}{6}} \sec^2\left(x + \frac{\pi}{6}\right) + \tan\left(x + \frac{\pi}{6}\right) \sec\left(x + \frac{\pi}{6}\right) dx$$

$$\Rightarrow 2\pi \left[\tan\left(x + \frac{\pi}{6}\right) + \sec\left(x + \frac{\pi}{6}\right) \right]_0^{\frac{\pi}{6}}$$

$$\Rightarrow 4\pi$$

13. If $f(3) = 18$, $f'(3) = 0$ and $f''(3) = 4$. Then, the value of

$\lim_{x \rightarrow 1} \ln\left(\frac{f(x+2)}{f(3)}\right)^{\frac{18}{(x-1)^2}}$ is equal to

- (1) 2
- (2) 4
- (3) 6
- (4) 8

Answer (1)

Sol.
$$\ln\left(\lim_{x \rightarrow 1} \left(\frac{f(x+2)}{f(3)}\right)^{\frac{18}{(x-1)^2}}\right) (1^\infty)$$

$$\ln\left(e^{\lim_{x \rightarrow 1} \frac{18}{(x-1)^2} \left(\frac{f(x+2)}{f(3)} - 1\right)}\right)$$

$$\ln\left(e^{\lim_{x \rightarrow 1} \frac{f'(x+2)}{2(x-1)} \cdot 18}\right)$$

$$\ln\left(e^{\lim_{x \rightarrow 1} \frac{f''(x+2)}{2} \cdot 18}\right)$$

$$\lim_{x \rightarrow 1} \frac{f''(3)}{2} = \frac{4}{2} = 2$$

14. If the domain of the function

$\cos^{-1}\left(\frac{2x-5}{11x-7}\right) + \sin^{-1}(2x^2 - 3x + 1)$ is

$[0, a] \cup \left[\frac{12}{13}, b\right]$ then $\frac{1}{ab}$ is equal to

- (1) -3
- (2) 3
- (3) 2
- (4) 4

Answer (2)

Sol. $\cos^{-1}\left(\frac{2x-5}{11x-7}\right)$ is defined as

$\frac{2x-5}{11x-7} \leq 1, \frac{2x-5}{11x-7} \geq -1$

$\frac{-9x+2}{11x-7} \leq 0, \frac{13x-12}{11x-7} \geq 0$

$\Rightarrow \frac{x-\frac{2}{9}}{x-\frac{7}{11}} \geq 0, \frac{x-\frac{12}{13}}{x-\frac{7}{11}} \geq 0$



$\Rightarrow \left[\left(-\infty, \frac{2}{9}\right) \cup \left(\frac{7}{11}, \infty\right) \right] \cap \left[\left(-\infty, \frac{7}{11}\right) \cup \left[\frac{12}{13}, \infty\right) \right]$

$\Rightarrow x \in \left(-\infty, \frac{2}{9}\right) \cup \left[\frac{12}{13}, \infty\right)$

Similarly, $\sin^{-1}(2x^2 - 3x + 1)$ is defined

When $-1 \leq 2x^2 - 3x + 1 \leq 1$


$\Rightarrow 2x^2 - 3x \leq 0 \Rightarrow x\left(x - \frac{3}{2}\right) \leq 0$

$\Rightarrow x \in \left[0, \frac{3}{2}\right]$

and $2x^2 - 3x + 2 \geq 0 \Rightarrow x \in \mathbb{R} \Rightarrow x \in \left[0, \frac{3}{2}\right]$

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)				JEE (MAIN)			
 ADVAY MAYANK AIR 36	 RUJUL GARG AIR 41	 ARUSH ANAND AIR 64	 SHREYAS LOHIYA AIR 6 Uttar Pradesh Topper 100 Overall	 KUSHAGRA BAINGAHA AIR 7 Uttar Pradesh Topper 100 Overall	 HARSH A GUPTA AIR 15 Telangana Topper 100 Overall		



$$4 \int_0^{0.6} [\sqrt{4-4y^2} - (1+y)] dy$$

$$I_1 = 4 \int_0^{\frac{3}{5}} \sqrt{4-4y^2} dy, \text{ put } y = \sin\theta, dy = \cos\theta d\theta$$

$$\Rightarrow I_1 = \int_0^{\theta} 2\cos^2 \theta d\theta = \int_0^{\theta} (1 + \cos^2 \theta) d\theta = \left[\theta + \frac{\sin 2\theta}{2} \right]_0^{\theta}$$

$$\therefore \sin = \frac{3}{5}, \cos\theta = \frac{4}{5}, \sin 2\theta = \frac{24}{25}$$

$$\Rightarrow I_1 = \left(\sin^{-1} \frac{3}{5} + \frac{12}{25} \right), I_2 = \int_0^{\frac{3}{5}} (1+y) dy = \frac{39}{50}$$

$$\Rightarrow \text{Area} = 4 \sin^{-1} \left(\frac{3}{5} \right) - \frac{6}{5}$$

- 18.
- 19.
- 20.

SECTION - B

Numerical Value Type Questions: This section contains 5 Numerical based questions. The answer to each question should be rounded-off to the nearest integer.

21. If $x^2 + x + 1 = 0$, then

$$\left(x + \frac{1}{x} \right)^4 + \left(x^2 + \frac{1}{x^2} \right)^4 + \left(x^3 + \frac{1}{x^3} \right)^4 + \dots +$$

$$\left(x^{25} + \frac{1}{x^{25}} \right)^4 \text{ is}$$

Answer (145)

Sol. $x^2 + x + 1 = 0 \begin{cases} \leftarrow w \\ \leftarrow w^2 \end{cases}$

$$\left(w + \frac{1}{w} \right)^4 + \left(w^2 + \frac{1}{w^2} \right)^4 + \left(w^3 + \frac{1}{w^3} \right)^4 + \dots$$

$$\left(w^{25} + \frac{1}{w^{25}} \right)^4$$

$$\sum \left(w^k + \frac{1}{w^k} \right)^4$$

$$k = 3x \Rightarrow w^{3x} + \frac{1}{w^{3x}} = 2$$

$$k \neq 3x \Rightarrow w^k + \frac{1}{w^k} = -1$$

$$\sum_{k=1}^{25} \left(w^k + \frac{1}{w^k} \right)^4 \Rightarrow 8(1+1+2^4) + 1$$

$$= 145$$

22. The sum of roots of the equation $|x-1|^2 - 5|x-1| + 6 = 0$ is

Answer (4)

Sol. $|x-1|^2 - 5|x-1| + 6 = 0$

Let $|x-1| = t$

$$t^2 - 5t + 6 = 0$$

$$(t-3)(t-2) = 0$$

$$t = 2, 3$$

$$|x-1| = 2 \text{ or } |x-1| = 3$$

$$x-1 = \pm 2 \text{ or } x-1 = \pm 3$$

$$\Rightarrow x = 3, -1, 4, -2$$

$$\Rightarrow \text{Sum} = 3 - 1 + 4 - 2$$

$$= 4$$

23. If $A = \{1, 2, 3, 4, 5, 6\}$, $B = \{1, 2, 3, \dots, 8, 9\}$. Then the number of strictly increasing functions from $A \rightarrow B$ such that $f(i) \neq i \forall i = 1, 2, 3, 4, 5, 6$ are

Answer (28)

Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY MAYANK
AIR 36



RUJUL GARG
AIR 41



ARUSH ANAND
AIR 64



JEE (MAIN)

SHREYAS LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSH A GUPTA
AIR 15
Telangana Topper
100 Overall



Sol.

- 1
- 2
- 3
- 4
- 5
- 6

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9

Case (i) If $f(1) = 2$ then 7C_5 functions

Case (ii) $f(1) = 3$ then 6C_5

Case (iii) $f(1) = 4$ then 5C_5

$$\Rightarrow 21 + 6 + 1 = 28$$

24.

25.



Our Problem *Solvers* shine bright in **JEE 2025**

JEE (Advanced)

ADVAY
MAYANK
AIR 36



RUJUL
GARG
AIR 41



ARUSH
ANAND
AIR 64



JEE (MAIN)

SHREYAS
LOHIYA
AIR 6
Uttar Pradesh Topper
100 Overall



KUSHAGRA
BAINGAHA
AIR 7
Uttar Pradesh Topper
100 Overall



HARSH
A GUPTA
AIR 15
Telangana Topper
100 Overall

